This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

۳۱۹۹۱

1276-16684
Stanford University Library
APR 9



Old Series Vol. XVI. New Series Vol. XI, No. 1

The Wilson Bulletin

No 46

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

March, 1904

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.





CONTENTS

	PAGE
The Cerulean Warbler. Lynds Jones	3
Partial List of Summer Birds of Holderness, N. H. Arthur C. Comey	5
The Larks of Germany. W. F. Henninger.	9
The Marsh Wren's Midnight Song. Chreswell J. Hunt.	13
A Double Nest of Red-eyed Vireo. Lynds Jones.	15
The Clock Factory. Esther Craigmile.	15
The New Year Bird Census.	16
A Disastrous Trip. W. F. Henninger.	21
An Improvident Kingbird. Lynds Jones.	22
The May Horizon.	23
A Door-Yard List of Birds. Robert L. Baird.	23
Editorial.	25
Notes and Publications Received.	27
List of Members.	30

The Wilson Bulletin

is published on the fifteenth of March, June, September and December, by the Wilson Ornithological Club, at Oberlin, Ohio, edited by Lynds Jones.

Subscription: Fifty cents a year, including postage, *strictly in advance.* Single numbers fifteen cents, unless they are "Special" numbers, when a special price is fixed. The Bulletin, including all "Special" numbers, is free to all paid up members, either Active, Associate or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, 1794 Lanier Ave., N. W. Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for reviews, should be addressed to Lynds Jones, Oberlin, Ohio.

Articles of general interest relating to the bird life are solicited. They should be in the hands of the editor not later than the 20th of the month preceding publication.

THE WILSON BULLETIN

AN ILLUSTRATED QUARTERLY MAGAZINE DEVOTED
TO THE STUDY OF BIRDS IN THE FIELD

Edited by LYNDS JONES

OFFICIAL ORGAN OF THE WILSON ORNITHOLOGICAL CLUB

LIBRARY
UNIVERSITY

OLD SERIES, VOLUME XVI. NEW SERIES, VOLUME XI.
1904

Published by the Club at Oberlin, Ohio

115450

TYPICAL NEST OF CERULEAN WARBLER

Courtesy of Dawson's Birds of Ohio

Photo by Lynds Jones

THE

WILSON BULLETIN

NO. 46

A QUARTERLY JOURNAL OF ORNITHLOGY

VOL. XVI. MARCH, 1904. NO. 1

THE CERULEAN WARBLER (Dendroica rara).

LYNDS JONES.

It is only recently that this warbler's nests have been found anything like commonly. Indeed, until recently its eggs have been sold at a high price and its nests much prized. The accompanying illustration of a nest found the past season near Oberlin gives nearly the typical nest and its position on the branch.

About Oberlin this warbler is found in the deeper woods, preferably where there is considerable moisture, at least during the spring months. Swampy woods are not much in favor. The trees must be small and slender, with few or no branches within twenty feet of the ground. Among these high branches the male feeds and sings, during the brooding days making regular rounds to and from the nest. One may locate the tree in which the nest is hidden by patiently watching the male on his rounds, for the nest tree marks a halt in his otherwise somewhat regular progress of search for food. It is also the focus for all of the different excursions in different directions.

The nests which I have actually found have been in trees which grow near, but not in, a shallow, sluggish woods spring stream. Spring freshets have removed enough earth to make a depression, but not enough to make a ditch. The tree has not been one of several forming a thicket of branches, but rather one which stands somewhat by itself. The nests have invariably been placed upon branches at some distance from the bole of the tree, on the top of a more or less horizontal fork of the branch. I have never seen a nest built into a per-

pendicular crotch, after the manner of the Yellow Warbler, nor distinctly saddled into the crotch like the Wood Pewee.

The male has never shown much concern about the nest, but the female makes some feeble demonstration if the nest is closely approached. I have never seen the evidence of distraction exhibited by so many birds when their nests are threatened. A sharp, metallic chip,and restless flitting within two rods of the nest is the extent of the female's demonstration.

The eggs are decidedly warbler-like in every respect, but with a tendency to grayness in the markings. However, a word description of the eggs of many of the warblers is worthless; it is sufficiently difficult to distinguish between the eggs when one has them in hand.

It seems strange that the eggs and nests of this warbler have been so long rarities in collections when the birds are certainly not rare breeders in almost any part of Ohio where conditions are at all favorable. No doubt the height at which the nests are placed is the main cause for the scarcity.

A PARTIAL LIST OF THE SUMMER BIRDS OF HOLDERNESS, NEW HAMPSHIRE.

BY ARTHUR C. COMEY.

The town of Holderness lies on the west shore of Squam Lake (or Lake Asquam), in almost the geographical center of the state of New Hampshire. It is divided naturally by the Squam Range and Little Squam Lake into three sections. I made the following observations in the area between the ridge of the Squam Range (elevation 2100 ft.) and Squam Lake (elevation 510 ft.), with my headquarters at Camp Algonquin, on the shore of the lake. Notes were taken during the months of July and August, 1899, 1900, 1901, and 1903, but chiefly in the latter season, which also included the last five days of June and the first four days of September.

The shores of the lake are for the most part quite low, but rocky, with occasional narrow sand-beaches, and are covered with rather young woods of mixed growth, with here and there a grove of pine or hemlock. Nearly all of the farm-houses are situated on the "old road," which in most places is about a half-mile from the lake. Between these and the new road, not far from the shore, there are overgrown pastures and maple groves, with open fields near the houses. Back from the old road are mountain pastures and birch woods, stretching up to the almost bare top of Squam Range, about a mile away. Most of the marsh-birds were seen at Bennett's Creek, which meanders through a meadow into Bennett's Cove. Several years ago the dam at the outlet of Squam Lake was raised so that the lower end of this little plain was inundated and a little lagoon was formed, with a sand-beach along the shore. According to Mr. Faxon, in the introduction to his Holderness list (The Auk, Vol. V., p. 149), the locality lies "near the border of the Alleghanian and Canadian Avifaunæ, although decidedly Alleghanian in its general character."

The list includes six species recorded as migrants only, and eighty-seven summer residents. The sixty-five species recorded by Mr. Faxon (June 4–12, 1885, and June 4–11, 1886) are starred.

*1. **Gavia imber.** Loon. Rather common, chiefly in the northern part of the lake.

2. **Aix sponsa.** Wood Duck. Rare. One record, a female at Hoag Island, July 18, 1903.

3. **Botaurus lentiginosus.** Bittern. Local; found at Bennett's Creek only.

4. **Ardea herodias.** Great Blue Heron. Rare; recorded as a migrant only. One record, one at Camp Algonquin, August 23, 1903.

5. **Butorides virescens.** Green Heron. Local; found at Bennett's Creek only.

6. **Nycticorax nycticorax nævius.** Black-crowned Night Heron. Locally rather common.

7. **Helodromas solitarius.** Solitary Sandpiper. Uncommon migrant. In 1903, a pair at Bennett's Creek, August 9 and 10.

*8. **Actitis macularia.** Spotted Sandpiper. Common.

*9. **Bonasa umbellus.** Ruffed Grouse. Quite common, but rare in 1903, owing to the cold spring, which killed nearly all the young.

10. **Accipiter cooperii.** Cooper's Hawk. Rare migrant. One record two on Squam Ridge, August 24, 1903.

*11. **Buteo borealis.** Red-tailed Hawk. Rare; recorded as a migrant only. One record, one on Rattlesnake Mountains, August 28, 1903.

12. **Buteo lineatus.** Red-shouldered Hawk. Uncommon.

*13. **Haliæetus leucocephalus.** Bald Eagle. Rare. In 1903, found about Rattlesnake Cove only.

14. **Falco sparverius.** Sparrow Hawk. Rare. Two records in 1903.

15. **Coccyzus americanus.** Yellow-billed Cuckoo. Uncommon.

*16. **Coccyzus erythrophthalmus.** Black-billed Cuckoo. Rather common.

*17. **Ceryle alcyon.** Belted Kingfisher. Common.

*18. **Dryobates villosus.** Hairy Woodpecker. Not common.

19. **Dryobates pubescens medianus.** Downy Woodpecker. Common.

20. **Sphyrapicus varius.** Yellow-bellied Sapsucker. Uncommon. Not found below 1000 ft. elevation.

*21. **Ceophlœus pileatus abieticola.** Northern Pileated Woodpecker. Rare. Two records in 1900.

*22. **Colaptes auratus luteus.** Northern Flicker. Rather common.

*23. **Antrostomus vociferus.** Whip-poor-will. Quite common.

*24. **Chordeiles virginianus.** Nighthawk. Rather common. On August 28, 1903, at 11 A. M., I saw a flock of 100 or more between the two summits of Rattlesnake Mountains, circling about and slowly moving southward, very close to the ground.

*25. **Chætura pelagica.** Chimney Swift. Very common, but not seen the last two or three weeks.

26. **Trochilus colubris.** Ruby-throated Hummingbird. Uncommon.

*27. **Tyrannus tyrannus.** Kingbird. Abundant.

*28. **Myiarchus crinitus.** Crested Flycatcher. Rather common.

*29. **Sayornis phœbe.** Phœbe. Very common.

30. **Nuttallornis borealis.** Olive-sided Flycatcher. Rare migrant. One record, two at Camp Algonquin, August 26, 1903.

*31. **Contopus virens.** Wood Pewee. Abundant.

*32. **Empidonax minimus.** Least Flycatcher. Quite common.

*33. **Cyanocitta cristata.** Blue Jay. Quite common.

*34. **Corvus americanus.** American Crow. Common.

*35. **Dolichonyx oryzivorus.** Bobolink. Rather common summer resident. Quite common in August.

36 **Molothrus ater.** Cowbird. Rare; recorded in 1900 only.

*37. **Agelaius phœniceus.** Red-winged Blackbird. Rather common.

*38. **Icterus galbula.** Baltimore Oriole. Rather common. In 1903, last seen August 13.

39. **Quiscalus quiscula æneus.** Bronzed Grackle. Locally common.

*40. **Carpodacus purpureus.** Purple Finch. Rather common summer resident. Common in August.

*41. **Astragalinus tristis.** American Goldfinch. Abundant.

*42. **Pocœcetes gramineus.** Vesper Sparrow. Common.

*43. **Passerculus sandwichensis savanna.** Savanna Sparrow. Local; common at Bennett's Creek.

*44. **Zonotrichia albicollis.** White-throated Sparrow. Uncommon; found on the Squam Range (elevation 2000 ft.). Also one singing bird, not 100 ft. above the lake, July 30, 1903.

*45. **Spizella socialis.** Chipping Sparrow. Common.

*46. **Spizella pusilla.** Field Sparrow, Abundant.

*47. **Junco hyemalis.** Junco. Common above 1000 ft. elevation. Not found below 700 ft.

*48. **Melospiza cinerea melodia.** Song Sparrow. Common, especially along the shores of the lake.

*49. **Melospiza georgiana.** Swamp Sparrow. Local. Probably regular at Rattlesnake Cove, where a singing bird was found, July 17, 1903.

*50. **Pipilo erythrophthalmus.** Towhee. Common in second growth woods, but not in overgrown pastures.

*51. **Zamelodia ludoviciana.** Rose-breasted Grosbeak. Uncommon summer resident. Common in August.

*52. **Cyanospiza cyanea.** Indigo Bunting. Abundant.

*53. **Piranga erythromelas.** Scarlet Tanager. Common.

*54. **Progne subis.** Purple Martin. Uncommon.

*55. **Petrochelidon lunifrons.** Cliff Swallow. Uncommon. In 1903, last seen August 12.

*56. **Hirundo erythrogaster.** Barn Swallow. Abundant.

*57. **Iridoprocne bicolor.** Tree Swallow. Common the first three weeks of July.

58. **Riparia riparia.** Bank Swallow. Rather common. Not seen after July 20.

*59. **Ampelis cedrorum.** Cedar Waxwing. Abundant.

*60. **Vireo olivaceus.** Red-eyed Vireo. Abundant. One nest had two young birds about five days old, August 1, 1903.

*61. **Vireo gilvus.** Warbling Vireo. Rare; recorded in 1900 only.

62. **Vireo flavifrons.** Yellow-throated Vireo. Rare.

63. **Vireo solitarius.** Blue-headed or Solitary Vireo. Uncommon.

*64. **Mniotilta varia.** Black and White Warbler. Uncommon.

*65. **Heliminthophila rubricapilla.** Nashville Warbler. Uncommon.

*66. **Compsothlypis americana usneæ.** Northern Parula Warbler. Local; Common at Hoag Island, July 18, 1903.

*67. **Dendroica æstiva.** Yellow Warbler. Uncommon.

*68. **Dendroica cærulescens.** Black-throated Blue Warbler. Rare.

69 **Dendroica coronata.** Myrtle Warbler. In July, one recorded at 700 ft elevation. Common at Hoag Island on July 18, 1903. In 1903, after August 6, a very common migrant.

70. **Dendroica maculosa.** Magnolia Warbler. Uncommon migrant. First seen, August 12, 1903.

*71. **Dendroica pensylvanica.** Chestnut-sided Warbler. Quite common.

72. **Dendroica striata.** Black-poll Warbler. Rare; recorded in 1900 only.

*73. **Dendroica blackburniæ.** Blackburnian Warbler. Rare summer resident. In 1903, after August 6, a rather common migrant.

*74. **Dendroica virens.** Black-throated Green Warbler. Quite common.

75. **Dendroica vigorsii.** Pine Warbler. Common in all red pine groves.

*76. **Seiurus aurocapillus.** Oven-bird. Common.

*77. **Seiurus noveboracensis.** Water-Thrush. Rare. In July, one record, one at Hoag Island, July 18, 1903. In 1903, the only migrant was seen at Camp Algonquin, August 9.

*78. **Geothlypis trichas brachidactyla.** Northern Yellow-throat. Common.

*79. **Wilsonia canadensis.** Canadian Warbler. Rare migrant. On record, one at Camp Algonquin, August 23, 1903.

*80. **Setophaga ruticilla.** American Redstart. Abundant.

*81. **Galeoscoptes carolinensis.** Catbird. Common.

*82. **Taxostoma rufum.** Brown Thrasher. Common.

83. **Troglodytes aedon.** House Wren. Rather common.

84. **Certhia familiaris americana.** Brown Creeper. Uncommon on the Squam Range. Not found below 1500 ft.

85. **Sitta carolinensis.** White-breasted Nuthatch. Rather common.

86. **Sitta canadensis.** Red-breasted Nuthatch. Rather common summer resident. In August, common migrant in the flocks of Warblers

*87. **Parus atricapillus.** Chickadee. Abundant.

88. **Regulus satrapa.** Golden-crowned Kinglet. Rather common above 1500 ft. elevation. Not found below 700 ft.

89. **Regulus calendula.** Ruby-crowned Kinglet. Rare. One record, one on Squam Range (elevation 1800 ft.), July 15, 1903.

*90. **Hylocichla fuscescens.** Wilson's Thrush or Veery. Abundant, but became uncommon towards the end of August.

*91. **Hylocichla guttata pallasii.** Hermit Thrush. Common.

*92. **Merula migratoria.** Robin. Quite common. Rather common on the bare top of Squam Range.

*93. **Sialia sialis.** Bluebird. Common.

94. **Passer domesticus.** English Sparrow. A few around some of the farm houses.

THE LARKS OF GERMANY.

W. F. HENNINGER.

Characteristic of the Larks is the bill, longer than that of the finches and buntings, longer and broader wings, and the earth-colored plumage, commonly called "lark gray." Only three species are ordinarily found in Germany, the others being little more than mere stragglers. Largest of these is the Crested or Tufted Lark, easily recognized by its conspicuous crest, then the far-famed Skylark, well known to all, and the smallest, the Woodlark, distinguished from the others by its size and the white wing-bands or spots.

A little home-body is the Tufted Lark. The dusty roadside, fence-posts, deserted wagons and ploughs, stone walls, and roofs are her home, where we can listen to her low, but not uninteresting song, or see her skipping through the air with lisping notes or running to and fro with raised hood, as she snatches up an insect or a grain. Having entered Germany but lately, she has become familiar in many a region. She loves the slopes of the mountains, the grain-fields, and is more common in sandy places than her cousins. There she makes her nest on the ground in the fields, meadows and gardens; four to six eggs it holds, white or cream-colored, with gray and brownish spots. Both parents brood and take care of the young in turn, looking for their food. Even winter's chills do not drive them away. They run ahead of us in the streets with Yellowhammers and Sparrows, or visit our back yards and barns to look for offal, dungheaps, spilled grain and similar tid-bits, being very tame and modest, easily contented at all times, liked by young and old for their confidence in man, till spring comes and the humble creature is forgotten for her more brilliantly singing relatives.

In golden splendor the morning sun appears on the eastern horizon, still tinted with the rosy hue of early dawn, like mil-

lions of liquid diamonds the dew is sparkling in the fragrant grass of the meadows, and from the leafy arches overhead the morning concert of our friends is sounding forth in honor of their Maker's praise. Look! From yonder field of grain, in ever-widening spirals, higher and higher rises a plain-colored bird, floating on the balmy breeze of the bright spring morn. The little throat is swelled with exuberant joy and the sweetest warbling reaches our ear, now loud and clear, as the spiral nears us, now soft and distant as it turns the other way. How insignificant is the song of our Horned Lark in comparison with this music of the Skylark! With delightful ecstasy we follow it with our eyes till it soars as a mere speck in the azure sky, and now is lost to our sight. From every field and meadow they seem to rise, in glorious blending their notes ring out till some bold Sparrow Hawk sweeps past us and buries his cruel talons in the merry songster's breast, slowly dragging it to some secluded spot for lunch. For a few moments all is quiet, but soon they rise again on all sides. The Skylark's lyric nature accompanies its movements with the singing life of its soul. With its slow rising it creates the beautiful warbling trill, and in the invisible realms of the ether the flute-notes of its nature melt away in circling waves, whirling they descend earthward again, till, like a dart of Cupid, it swoops into the surging sea of grain. There she seeks her food—humble fare—grain, insects and worms. There she builds her excuse of a nest—a mere hollow, lined with rootlets. There she watches her four or five white, brown-speckled treasures, there she raises her broods in safety from the haunts of man, but not always of the reaper's scythe.

But come again with me to the meadows, when the twilight falls and the sun has gone to sleep. Softer now the tints of heaven, hushed the voices of the joyous spring, murmuring lowly are the forests' trees, slowly homeward turneth man and beast. Far, far away the Lark's melodious voice is heard; but no! there she sits behind a clod in the furrow near by. More like that insolent ruffian, the Sparrow, she seems in her plain garb, and I understand your look of disappointment. But 'tis often so; in plain garb is hidden many a jewel. And as she runs over the clods and through the grain with heaving breast, graceful neck, alert and free, you wilt soon see, 'tis

not the Lord of ruffians, but a star of heaven come down to earth. Now she stops! Sweetly, tenderly falls the song upon thine ear. It speaks to thee of by-gone days, of love's first dream, of childhood's play, of monher's tender care, of the old home by the wayside, of the brook's clear flowing waters, of all that is dear and sweet to memory and heart. The German nation's "Gemuet" is in the song. A yearning and a longing for yonder world comes over the soul as the dream-like love notes call 'till the last sound vanishes in the solemn darkness of the night. And Shakespeare called the Lark "shrillgorg'd"! Had the great poet no ear for music? Or is it because there is such a brutal shrillness in all British efforts of philanthropy and civilization from the time of Richard Coeur de Lion down to the days of Lord-butcher Kitchener in the Transvaal, that e'en the Lark to them was "shrillgorg'd"? For once, immortal Briton, thou wert mistaken! 'Tis not a "shrillgorg'd" monster, but nature's best, its own harmonious melody that reaches us in the Lark's divine lullaby and lets us feel the poet's truth:

> "And I so ravished with her heavenly note
> I stood entranced and had no room for thought."

Did you ever wander homeward through the woods, when the dark-winged angel of the night has kissed the fields and hamlets and breathed down to the roots of everything that lives? Come with me then, where the bushy beeches whisper softly, where the gloomy firs stand out in silent awe, broken here and there by the ghost-like drooping branches of the birch, whose bark reflects the moonlight's silvery rays,

> "Als waere dran in stiller Nacht
> Das Mondlicht blieben hangen."

where the blooming heath's carpet, in shining red and white, sends out its delicious odor. Only the distant croaking of the frogs in the treacherous shimmer of the marsh, the howling "boohoo" of the Horned Owl, that robber-knight of the winged world, the spinning of the Goatsucker is heard,—all else is quiet in the moor and heath. The Robins sleep in safety, the Mavis and the Skylark have ceased their song, the Nightingale's bosquets and man's abode are far away, only the elfs dance over the meadows' fog, inviting you to join

their merry throng, and erlking's will-'o-the wisp lures you into unknown depths. Hurriedly we pass along the barren fallow waste, close by the sombre forest. But lo! Up from the sterile barren goes a song, clear, bright and cheery. Is it a new spirit form that wants to mock us? Everywhere it seems to carry peace and comfort by its virile master-melody—a lovely greeting to the weary, wandering man! High up into the air they soar, those earnest, flute-like notes. Forgotten are the hobgoblins of the night, sweet thoughts of rest and joy enter thy heart, trust to God is poured into thy soul! Yes, praised be thou, O woodlark, thou living voice of mountain-heath and solitude! Though not crowned with the Nightingale's laurels, not made immortal by the poet's strain, to me thou art dearer in thy modesty. Not only in the dazzling light of day thou cheerest us, the dwellers on the earth, but in the night, "which is no mortal's friend," thou bringest with thy heavenly flute a welcome to the lonely pilgrim!

'Tis a song simpler than the Skylark's, but still more pleasing to the ear, a true ring of the forest's rustling leaves. It is a slow verse of some ten to twelve deep flute-notes, followed in a few seconds by a higher trill. Each spiral of its flight brings out a new verse, clear, powerful and melodious, full of masculine strength and not so much of feminine sentimentality as the Skylark's song. And even in the fall good singers let their voices be heard, though softer then and more subdued. Her flight is not so wide, more swerving than the Skylark's, and on the ground she is more modest in her agile movements than the latter. Twice a year she makes her pretty, well-built home on the ground in the heath or by the elder-bushes, to hold her five white, gray-speckled eggs. Carefully running on the ground for quite a distance, she brings the food to her loudly clamoring young. One of the latest birds to leave us in the fall, she is one of spring's earliest messengers, with her heavenly song, though snow may still cover the heath and forest, and we rejoice to have her with us again.

THE MARSH WRENS' MIDNIGHT SONG.

CHRESWELL. J. HUNT.

The Pensauken Creek forms a part of the boundary line between Burlington and Camden Counties, New Jersey. It twists and turns in a very irregular course, and at last finds its way into the Delaware River.

As with all these tide-water creeks, the banks are high and wooded on one side, while on the other side, for the most part, lie low stretches of alder swamps, covered in the late autumn with a rank growth of wild rice, spatter-dock and pickerel-weed, with here and there a clump of rose-mallow or a gorgeous cardinal flower.

It is here, among this almost impregnable growth, that countless numbers of Long-billed Marsh Wrens find a congenial home, building their globular nests in the alder bushes just above the water, and when within the swamp one is never beyond hearing of their rippling song.

They seem to be fairly overflowing with music—a bubbling, gushing song that seems rather to have had its birth above some rushing mountain stream, than above these sluggish waters. Before one has stopped singing another takes up the strain, hence all day long these marshes are merry with music, nor does the coming of night silence them.

On the moonlight nights of July and August, these happy little songsters make merry the midnight hours with their cheery warble.

They mount into the air, singing, and then dive back again among the sheltering reeds. The song is no doubt the same as that sung in the daylight, but the night gives to it a certain charm. One must hear it, mingled with the quivering call of a Screech Owl and the "quauk, quauk" of Night Herons, to fully appreciate it.

These concerts are not restricted to moonlight nights. I spent the night of August 8, 1903, in a boat among these swamps. It was cloudy, and now and then a light shower fell, but the Wrens were in song. Could they be otherwise? To the bird lover who has never witnessed this night performance, there awaits a very pleasant experience indeed.

DOUBLE NEST OF RED-EYED VIREO
Courtesy of Dawson's Birds of Ohio
Photo by Lynds Jones

A DOUBLE NEST OF RED-EYED VIREO.

LYNDS JONES.

The accompanying illustration represents a nest of the Red-eyed Vireo, taken more than a dozen years ago at my old home in Iowa. As shown in the illustration, there were two eggs in one side and one in the other. When the nest was found the mother bird was sitting upon the two egg side, but there were evidences showing that she had also sat upon the one egg side. To all appearances the two nests are exactly alike, and each is practically complete, only a small amount of the material being common to both nests. It could rarely happen that a main branch would have two twigs growing from it on opposite sides at almost the same level, and at practically the same angle. It is hardly conceivable that two females or two pairs of birds could have built these nests. At any rate, there was no evidence of it, for only one pair of birds was to be found in the vicinity. Neither is it likely that the male built one nest while the female built the other. The evidence is strongly in favor of the same architect for both nests. It looks like a case of indecision between two equally favorable nesting sites. We are not surprised at evidences of indecision in the Long-billed Marsh Wren—if that is indecision rather than a method of working off superfluous energy—but the Vireos have not been supposed to do such things.

THE CLOCK FACTORY.

ESTHER CRAIGMILE.

Our little party was eager with expectancy, for it stood on the edge of a pine grove from which proceeded no small commotion. It was eleven o'clock in the morning and the colony was just retiring for the day. We were nearer now, and could see they were Black-crowned Night Herons (*Nycticorax nycticorax nævius*). Some parent birds sat silently at the top of the dead pines, while some were feeding the young. The larger trees contained three and four nests. Three or

four eager heads were outstretched from these nests, while some were making awkward attempts at walking on the naked pine branches. There were hundreds of nests in an area of about two acres.

The young kept up a constant ticking, fast and slow, high and low, while an occasional *squawk* was heard from one of the old birds. It was a typical clock factory, where the *squawk* played the part of the cuckoo clock, and the young ticked off the seconds. They seemed little disturbed by our party of six walking below them. It was little wonder that the flock had killed the beautiful pine grove, when the ground underneath was carpeted with twigs. The whitewashed undergrowth was developing ruggedly in spite of the environment. Pieces of pale blue shell, dead fish and one dead bird, half grown, were found below the nests. Barring all sensations of the olfactory tract, it was a delightful experience.

THE NEW YEAR BIRD CENSUS.

This, our second annual bird census, shows an increase of near 50 per cent. in the number of reports sent in, and that in the face of more wintry weather and general difficulty in getting into the fields. The reports show that even under the most adverse conditions the birds are certain to be present in some numbers.

Vicinity of Alma, Lincoln County, Maine. Time 9:25 to 11:55 a. m. Weather fine, no wind. Temperature 6 to 22.

Redpoll, one flock of 30; Red-breasted Nuthatch, one flock of about 8; Chickadee, 7; Golden-crowned Kinglet, 2.

Total, 4 species about 47 individuals.

HAROLD E. PORTER, CHARLES H. ROGERS.

Bristol, Conn. Time 7:30 a. m. to 4:30 p. m. Weather, cloudy at first, clearing by 8:30 a. m., clear the rest of the day. Six inches of snow, powdered over by a light fall the day before.

Bob-white, 6, and fresh tracks; Ruffed Grouse, 4; Red-tailed Hawk, 1; Hairy Woodpecker, 4; Downy Woodpecker, 1; Blue Jay, 16; Crow, 205†; American Goldfinch, 250†; Tree Sparrow, 102†; Junco, 15; Song Sparrow, 1; Winter Wren, 1; White-breasted Nuthatch, 9; Chickadee, 14; Golden-crowned Kinglet, 3; Hermit Thrush, 1.

Total, 16 species, over 630 individuals.

ELBERT E. SMITH, FRANK BRUEN.

Boston, Mass., including the Parkway from Longwood through Jamaica Park and the Arnold Arboretum; also the Arnold estate and Hall's pond, Brookline. Time 8:50 a. m. to 1:15 p. m.

Bob-white, 15; Sparrow Hawk, 1; Barred Owl, 1; Flicker, 15; Blue Jay, 13; Crow, 19; Canadian Pine Grosbeak, 1; Goldfinch, 9; White-throated Sparrow, 10; Tree Sparrow 11; Junco, 1; Song Sparrow 6; Northern Shrike, 1; Brown Creeper, 2; White-breasted Nuthatch, 3; Chickadee, 23; Golden-crowned Kinglet, 8; Hermit Thrush, 1.

Moon Island and Squantum, Mass. Time 2:10 to 5:25 p. m.

Black-backed Gull, 2; Herring Gull, 600 (estimated); Red-legged Black Duck, 9; American Scaup Duck, 600 (estimated); American Golden-eye, 450 (estimated); Buffle-head, 6; Horned Lark, 10; Crow, 137; Meadowlark, 4; American Goldfinch, 60 (approximate); Snowflake, 22; Tree Sparrow, 6 (one singing); Song Sparrow, 8; Chickadee, 4.

Total, 27 species, 2059 individuals.

FRANCIS G. AND MAURICE C. BLAKE.

Cambridge, Mass., through the Fresh Pond Marshes, over Arlington Heights to Waverly; eight miles. Time 8:45 to 12:15. Clear, modate N. W. wind. Temperature 24 to 35. Five inches of snow on the ground.

Bob-white, 8; Red-shouldered Hawk, 1; Hairy Woodpecker, 1; Downy Woodpecker, 5; Northern Flicker, 1; Blue Jay, 2; Crow, 10; Red-winged Blackbird, 2; Canadian Pine Grosbeak, 2; Goldfinch, 2; Tree Sparrow, 30; Song Sparrow, 6; Swamp Sparrow, 4; Myrtle Warbler, 1; Brown Creeper, 5; White-breasted Nuthatch, 3; Chickadee, 15; Golden-crowned Kinglet, 10.

Total, 18 species, 108 individuals. ARTHUR C. COMEY.

Philadelphia, Pa. Time 3 to 5 p. m. Place, a cemetery two miles from the city hall. Weather, clear, ground with a little snow in some places.

Red-tailed Hawk, American Crow, 1; American Goldfinch, 2; Junco (estimated) 40; White-throated Sparrow, 6; Song Sparrow, 5; Cardinal, 1; Carolina Wren, 1; Winter Wren, 1; White-breasted Nuthatch, 2.

Total, 10 species, 60 individuals. LOUIS T. PARKE.

Cupola, Chester County, Pa. Time 11 a. m. to 4:30 p. m. Weather, clear, ground scantily covered with snow here and there. Wind, N.W. to N.. brisk. Temperature 43 to 38. About 250 acres gone over thoroughly.

Downy Woodpecker, 4; Northern Flicker, 1; American Crow, several hundred; Tree Sparrow, 35; Junco, 60; Song Sparrow, 20; White-breasted Nuthatch, 3; Chickadee, 30; Tufted Titmouse, 1; Golden-crowned Kinglet, 1; Hermit Thrush, 1.

Total, 12 species, 157 individuals (excluding Crows).

CHRESWELL J. HUNT.

Central Park, New York, January 1, 1904. Time 10:15 a. m. to 12:45 p. m. and 3 to 4 p. m. Clear, wind slight. Temperature 38.

Herring Gull, 3; Hairy Woodpecker, 2; Downy Woodpecker, 2; Starling, (about) 30; European Goldfinch, 4; White-throated Sparrow, (about) 30; Junco, 1; Song Sparrow, 1; Cardinal, 4; Brown Creeper, 1

White-breasted Nuthatch, 2; Red-breasted Nuthatch, 1; Chickadee, (about) 10; Robin, 1.

Total, 14 specimens, 92 individuals.

CLINTON G. ABBOTT AND MAUNSELL S. CROSBY.

Leonia, Nordhoff, Teaneck and Englewood, N. J. Time 9:15 a. m. to 4:15 p. m. Weather, fine. Wind, light west. Ground snow-covered.

Herring Gull, about 200 (in Hudson River); Red-tailed Hawk, 5; Snowy Owl, 1; American Crow, 19; White-throated Sparrow, 7; Tree Sparrow, 93; Junco, 67; Song Sparrow, 23; Myrtle Warbler, 5; Brown Creeper, 1; White-breasted Nuthatch, 1; Chickadee, 18; Golden-crowned Kinglet, 3.

Total, 13 species, 443 individuals. GEORGE E. HIX.

Cadiz, O. Time 9:30 to 12-30 and 1:30 to 4:30. Clear until 1:30 p. m. Wind, N. W. to N. E. Temperature at starting, 24, at noon, 31, on return, 27. Distance walked, by pedometer, registered 14 miles. Five inches of snow.

Bob-white, 15; Ruffed Grouse, 1; Red-tailed Hawk, 1; Great Horned Owl, 1; Hairy Woodpecker, 3; Downy Woodpecker, 15; Red-bellied Woodpecker, 4; Northern Flicker, 1; Horned Lark and Prairie Horned Lark, 65; Blue Jay, 3; Tree Sparrow, 34; Junco, 22; Song Sparrow, 9; Cardinal, 16; Carolina Wren, 4; White-breasted Nuthatch, 17; Chickadee, 23; Tufted Titmouse, 25.

Total, 19 species, 259 individuals. HARRY B. MCCONNELL.

Jefferson, O. Time 8 to 11:30 a. m. and 1 to 4:30 p. m. Temperature, 25. Slight west wind or none; mostly cloudy. Snow 8 inches, drifts in some places.

Ruffed Grouse, 1; Red-tailed Hawk, 1; Barred Owl, 2; Hairy Woodpecker, 2; Downy Woodpecker, 8; Northern Pileated Woodpecker, 2; Red-bellied Woodpecker, 2; Northern Flicker, 1; Blue Jay, 8; Tree Sparrow, 2; Junco, 5; Carolina Wren, 1; White-breasted Nuthatch, 9; Red-breasted Nuthatch, 1; Tufted Titmouse, 2; Chickadee, 13; Golden-crowned Kinglet, 1.

Total, 17 species, 61 individuals. ROBT. J. SIMM.

McZena, O. Time 7:50 to 11:50 and 1 to 4:30. Partly cloudy, partly clear. Wind S. W., light in the morning, N., brisk in the afternoon. Temperature 14 to 32. Distance walked, about 8 miles. Alone in the morning, with Mr. Zeno Metcalf in the afternoon.

Bob-white, 15; Ruffed Grouse, 2, Mourning Dove, 40; Red-tailed Hawk, 3; Sparrow Hawk, 1; Belted Kingfisher, heard several times; Hairy Woodpecker, 1; Downy Woodpecker, 2; Red-headed Woopecker, 4; Red-bellied Woodpecker, 2; Northern Flicker, 1; Prairie Horned Lark, 30; Blue Jay, 16; American Goldfinch, 27; Tree Sparrow, 60; Junco, 40; Song Sparrow, 6; Cardinal, 6; Brown Creeper, heard; Carolina Wren, heard singing; White-breasted Nuthatch, 20; Chickadee, heard; Tufted Titmouse, 13; Golden-crowned Kinglet, 1; Bluebird, 6.

Total, 25 species, 300 individuals. C. L. METCALF.

McConnelsville, O. Time 8 to 12 a. m. Cloudy and snowing. Temperature, 26. Wind north. Two inches of snow. Distance traveled, two miles.

Red-tailed Hawk, 1; Hairy Woodpecker, 2; Downy Woodpecker, 12; Red-bellied Woodpecker, 8; Northern Flicker, 1; American Goldfinch, 1. Tree Sparrow, 5; Junco, 30; Song Sparrow, 4; Cardinal, 45; Carolina Wren, 9; White-breasted Nuthatch, 8; Tufted Titmouse, 13, Chickadee, 12.

Total 14 species, 151 individuals.

C. H. Morris and E. J. Arrick.

Oberlin, O. Temperature 24 to 28. Cloudy. Wind north, light to brisk. Oberlin and immediate vicinity.

Bob-white, 1; Sparrow Hawk, 1; Screech Owl, 1; Hairy Woodpecker, 5; Downy Woodpecker, 7; Northern Flicker, 5; Blue Jay, 7; Meadowlark, 6; American Goldfinch, 3; Tree Sparrow, 5; Junco, 60; Song Sparrow, 2; Cardinal, 4; White-breasted Nuthatch, common; Tufted Titmouse, 6; Chickadee, common; Bronzed Grackle, probable.

Total, 16 species, over 60 individuals. Francis M. Root.

Oberlin, O. Weather as above. Trip of 22 miles to Lake Erie, at Oak Point, and west via Vermilion river and Chance Creek and the old quarry region.

Herring Gull, 1; Bob-white, 14; Mourning Dove, 1; Sharp-shinned Hawk, 1; Red-tailed Hawk, 1; Golden Eagle, 1; Long-eared Owl, 1; Barred Owl, 1; Screech Owl, 2; Hairy Woodpecker, 6; Downy Woodpecker, 12; Red-bellied Woodpecker, 4: Northern Flicker, 2; Blue Jay, 20; American Crow, 1; Goldfinch, 3; Vesper Sparrow, 1; Tree Sparrow, 61; Song Sparrow, 18; Junco, 39; Cardinal, 29; Northern Shrike, 1; Carolina Wren, 2; Brown Creeper, 1; White-breasted Nuthatch, 21; Chickadee, 19; Tufted Titmouse, 18; Golden-crowned Kinglet, 2.

Total, 28 species, 283 individuals.

Lynds Jones, D. E. Nye, F. B. Wolfe.

Belle Isle, Detroit River, Mich., January 1, 1904. Time 10 a. m. to 2 p. m. Four inches of snow on ground, a fine, mild day for this time of the year.

Cardinal, (male and female); Tufted Titmouse, 25; Chickadee, 50; White-breasted Nuthatch, 10; Red-breasted Nuthatch, 2; Brown Creeper, 4; Blue Jay, 1.

Total, 8 species, about 90 individuals.

This same day Norman A. Wood saw a Cardinal at Ann Arbor, Mich. Raymond E. Miller captured an American Bittern at Waterford, Mich., and J. Claire Wood saw a Northern Shirke at Wyandotte, Wayne County, Mich. Alexander W. Blaine, Jr.

Hinsdale, Ill. Time 8 a. m. to 1 p. m. Cloudy; north wind. Temperature 15. Two inches of snow, with drifts. A five-mile tramp through low prairie, east of town, and woods skirting east and north sides.

Hairy Woodpecker, 1; Horned Lark, 25; Blue Jay, common; American Crow, common; Evening Grosbeak, 20; Purple Finch, 15; Junco, 50; Brown Creeper, 3; White-breasted Nuthatch, 5; Chickadee, 25; Golden-crowned Kinglet, 4; Tree Sparrow, 100.

Total, 12 species, over 250 individuals. ESTHER A. CRAIGMILE.

Rock Island, Ill. Time 8:50 a. m. to 12:50 p. m. Mostly clear. Wind N. E., strong. Ground partly covered with snow. Temperature 22 to 24. A 12-mile tramp through woods and fields.

Bob-white, fresh tracks; Downy Woodpecker, 2; Red-headed Woodpecker, 2; Prairie Horned Lark, 8; Blue Jay, 15; American Crow, 4; Tree Sparrow, 30; Junco, 7; Brown Creeper, 1; White-breasted Nuthatch, 2; Chickadee, 14.

Total, 11 species, 85 individuals. BURTIS H. WILSON.

Mt. Carmel, Ill. Time 7:30 to 11 a. m. Distance traveled, about seven miles. Cloudy, threatening rain: light west wind; ground bare. Temperature 40.

Canada Goose, 18; Hawk (Coopers ?) 1; Red-tailed Hawk, 1; American Sparrow Hawk, 1; Hairy Woodpecker, 5: Downy Woodpecker, 4; Yellow-bellied Sapsucker, 2: Red-headed Woodpecker, 12. Red-bellied Woodpecker, 7; Flicker, 8; Blue Jay, about 20; Crow, about 25; Meadowlark, 16; Goldfinch, about 30; Tree Sparrow, about 50; Junco, about 100; Song Sparrow, 6; Cardinal, 15; Loggerhead Shrike, 1; Carolina Wren, 3; Bewick's Wren, 1; Winter Wren, 1: Brown Creeper, 1; White-breasted Nuthatch, 4; Tufted Titmouse, about 50; Chickadee, about 20; Golden-crowned Kinglet, 4; Bluebird, 10.

Total. 28 species, about 416 individuals. CHAS. F. BRENNAN.

Washington, D. C., Dec. 30, 1903. Ground just covered with a light fall of snow which came during the preceding night. Sky clear. Sharp northwest wind blew during most of the day, making it difficult to catch the low notes of birds, but also tending to keep the birds bunched in the sheltered spots. Temperature not known, but just warm enough in middle of day to thaw the snow in the sunshine. Time spent in the fields, 9 a. m. to 11:30 a. m., along Piney Branch and Rock Creek, in in the northwestern outskirts of the city; 12 m. to 4:30 p. m., along the eastern branch of of Potomac, just east of the city.

Species given in the order in which first observed, and numbers of each species seen, accurately counted except in the following cases, where accurate count was impossible and numbers only estimated, namely: Junco, Crow, Turkey Vulture, White-throated Sparrow, Fish Crow, Tree Sparrow.

Junco, 40; American Crow, 1200; Song Sparrow, 11. White-breasted Nuthatch, 5: Cardinal, 14; Tufted Titmouse, 11; Downy Woodpecker, 2; Hairy Woodpecker, 1; Chickadee, 6; Winter Wren, 2: Yellow-bellied Sapsucker, 1; Turkey Vulture, 11; White-throated Sparrow, 18: Goldfinch, 4; Carolina Wren, 4; Red-breasted Nuthatch, 1; Hermit Thrush, 1; Cedarbird, 16; Fish Crow, a few among other Crows; Golden-crowned Kinglet, 1; Meadowlark, 2; Tree Sparrow, 20; Brown Creeper, 2; Duck, one of the larger wild ducks, but species not recognized; Sparrow Hawk, 1. THOS. H. LEVERING.

A DISASTROUS TRIP.

W. F. HENNINGER.

For several years it had been the desire of the writer to visit the famous Port Clinton (Ottawa County, O.) marshes to explore the bird world at the same place where Dr. Langdon had been so fortunate in 1880. The afternoon of June 1st found me at Port Clinton, in a terrible rain and wind storm. Early the next morning found me out on the Portage River exploring the marshes for miles, then in the afternoon out on the Lake Erie waters. June 3rd, and 4th, on which day I was joined by Rev. W. Leon Dawson of Columbus, found me on Sandusky Bay, on the grounds of the Portage Gun Club and the Wynous Point Shooting Club. It was the same scene everywhere. The storm had carried the water higher inland than for the last eighteen years, and everything had been flooded. The only birds that had escaped destruction of their nests were the Red-winged Blackbird and the Long-billed Marsh Wren. All our searching was in vain. Not a Grebe, not a Least Bittern were seen, but few Coots and Gallinules heard. On Friday, Brother Dawson ascended the dizzy height of the water works tower at Sandusky, but as far as the eye could see, the waters spread over the Sandusky marshes. Under these condition it was a wonder that any birds had escaped, and our record of sixty-six species noted during our three days' stay will still compare favorably with Langdon's ninety, as seen in 1880.

Of interest were only a troop of five Bonaparte Gulls and four Semi-palmated Sandpipers on June 3rd, several Black Terns and sixteen Turnstones, seen on June 4th on Sandusky Bay. The Turnstones were found on a newly planted cornfield, and it was a pretty sight to watch them turning over the clods and catching their prey. It was in the club house of the Wynous Point Shooting Club that we found the most interesting things, stored away in the collection of birds, and enabling Brother Dawson and myself to bring home at least a few noteworthy records from this disastrous trip.

The first was a specimen (sex unknown) of Chen hyperborea nivalis, shot in the fall of 1886.

2. Trumpeter Swan 1877 and White-fronted Goose, shot in the fall of 1868.

3. White-winged Scoter, shot in fall of 1881.

4. Peregrine Falcon, shot in fall of 1882, by Colonel E. A. Scoville.

5. A Hybrid, between Anas obscura and Anas boschas, killed in the fall of 1878, by Judge E. B. Sadler.

6. A pure Albino Redhead, killed in fall of 1880.

7. A partial Albino Coot, and

8. A partial Albino Wilson s Snipe, both killed in the fall of 1881, by C. J. Clark.

The Coot has many white feathers on the head and neck, also smaller white feathers on various parts of the body. The Wilson's Snipe has the upper part of both wings almost entirely white.

9. A Snowy Owl, shot in fall of 1881.

All of the Ohio Ducks were represented in this collection, among them the rare Gadwall in several specimens.

AN IMPROVIDENT KINGBIRD.

LYNDS JONES.

One of the Oberlin public school teachers has given me a Kingbird's nest which a pupil of hers found and preserved. The nest is normally made of strings, rags, weeds, hair, twigs and rootlets. It was built in an apple tree in no usual manner, except that the birds failed to notice that they had built into one side of the nest a fertile apple bud. In the natural course of growth the apple had to have room, and soon began to disarrange the side of the nest When completely grown the apple was fully two inches in diameter, and since the inside diameter of the nest is only a half-inch larger, there was little left of the nest cavity. No doubt the young had left the nest before the apple had attained much size, but probably not before it began to encroach upon the space intended for the young birds. I have heard of growing apples in bottles and other fanciful receptacles, but never heard of the use of a bird's nest before.

THE MAY HORIZON.

So much interest was shown in the May Horizon last year, and such good results grew out of it, that we are ready to undertake it again the coming May. If possible, it would be the best plan to spend several whole days with the birds during May; but if that is not possible, then plan to spend some one whole day with them, in the definite expectation of making your list for that day reach the hundred mark. Woods, fields, marshes, and bodies of water should be included in the field of operations if possible. It is always best to give the region as much study as possible the day previous to the day set for the supreme effort, so that you may be able to find the less common species with the least expenditure of time. You cannot hope for a very large list unless the study begins as soon as the morning light is strong enough to make large birds visible. I expect to get the owls during the morning twilight, and the Whipporwill during the evening twilight. If you must sleep, do it at high noon. You will need four lunches at least, and if you eat them while you watch for birds, you will be surprised at the number of new records that are made during the meal in the woods or beside the lake or pond. Lists sent to the editor not later than the 12th of May can be printed in the Bulletin.

A DOOR-YARD LIST OF BIRDS.

BY ROBERT L. BAIRD.

Many local lists are monthly published in our bird magazines. These are valuable and show the increased interest in bird study. Many of these lists are from favored localities and few are from such limited areas as one's own door-yard. It is with the purpose of showing some of the possibilities in this time that I contribute the following account from my notes and check-book for September and the first twenty days of October, 1902:

At this time my home was with Dr. Chas. Hancock, of Denmark, Lee County, Iowa. Dr. Hancock's place is situated at the southeast edge of town. With the exception of one house across the road, there is no other within a hundred and fifty yards. The lot is about a hundred and fifty yards long by sixty to seventy wide. About the house are six or seven evergreen trees, Norway Spruce, Cedar and Pine, several fruit trees and an Osage Orange tree. At the farther end of the lot is a row of Locust trees. Part of the place was devoted to a garden, but a large part was simply in grass.

Within these limits I saw or heard forty different species of birds in the seven weeks of the fall above mentioned. Not

the most favorable time of the year, by any means, for finding birds, yet I was surprised almost every day by some new visitor. I wish I might have kept a list for a whole year there. I would be willing to wager that in the course of twelve consecutive months, ninety or one hundred different species of birds could be seen or heard in this yard or flying over it. The list which I have, I am sure could have been made much larger, but my school duties kept me away from eight o'clock in the morning to four in the afternoon.

The best find of the period, for myself at least, was Bewick's Wren. He came on one of the last days of the period, a cool, cloudy day. I had a good look at him, for he did not seem to notice my presence much He was investigating everything around, especially through the large wood pile.

It was just a few rods down the road that I found three Leconte's Sparrows one week, but I found none right in the yard.

One wet, foggy morning, just before leaving for school, I happened to look out of the window to a cedar tree, a few feet from the house. It was fairly swarming with little birds and warblers. Among them were Kinglets, an Ovenbird, Black-throated Green Warblers, Yellow Warblers, a Myrtle Warbler, and two or three which I could not identify.

One evening just before twilight, I could hardly believe my ears when, from a little clump of bushes, came an occasional "chewink" of a retiring Towhee. I investigated more closely and watched for some time a female Towhe from a distance of only six feet.

Taking it all in all, I found the numbers of individuals in this region of southeastern Iowa far larger than here about Oberlin. My first of January all day horizon, published in the first Bulletin of 1903, was a good example of this fact. Especially was this true of the Larks, Blue Jays, Warblers and Sparrows.

Following is the complete list of birds found in the yard I have described:

Mourning Dove, few; Screech Owl, few; Hairy Woodpecker, few: Downy Woodpecker, few; Red-headed Woodpecker, tolerably common; Red-bellied Woodpecker. few; Northern Flicker, tolerably common; Chimney Swift, common; Nighthawk, few; Crested Flycatcher, few; Wood Pewee, few; Prairie Horned Lark, 2; Blue Jay, common; American Crow, few; Meadowlark, tolerably common; Baltimore Oriole, few; Bronzed Grackle, common; English Sparrow, common; American Goldfinch. tolerably common; Whitethroated Sparrow, few; Field Sparrow, few; Slate-colored Junco, few; Towhee, 1; Barn Swallow, few; Warbling Viero, few; Yellow Warbler, few; Myrtle Varbler, 1; Chestnut-sided Warbler, 1; Black throated Green Warbler, 2; Oven-bird, 1; Catbird, few; Brown Thrasher, few; Bewick's Wren, 1; House Wren, few; Whitebreasted Nuthatch, few; Chickadee, tolerably common; Golden-crowned Kinglet, few; Ruby-crowned Kinglet, few; American Robin, few; Bluebird, common, migrating.

THE WILSON BULLETIN

A Quarterly Magazine Devoted to the Study of Living Birds
Official Organ of the Wilson Ornithological Club

Edited by LYNDS JONES
PUBLISHED BY THE CLUB, AT OBERLIN, OHIO

SUBSCRIPTION RATES

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number, postage paid.

Price in all counries in the International Postal Union, 65 cents a year, 20 cents a number.

Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr. Frank L. Burns, Berwyn, Penn., or to Mr. John W. Daniel, Jr., Q street, N. W., Washington, D. C.

EDITORIAL.

Spring has begun when the first birds arrive from the south. We recognize in the appearance of the birds the beginning of new conditions of both weather and landscape. To the true ornithologist spring covers the time during which the birds are passing north, summer the time of nesting and molting, after the movement north has ceased and before the breeding birds have ceased to remain in hiding, autumn is the season of southward migration, and winter the remainder of the year. We are just now entering upon spring, with its predictions of coming birds and coming mild weather and have little difficulty in spurring ourselves to more effort in bird study. We do well when we pause to take stock of what we know and plan to accomplish the most in this delightful study with the time at our disposal. There are two things which will be of great service in this study. The first is: Plan your study to suit your own convenience. The second: faithfully keep a note-book record of what you see and learn. By the record I mean what you actually see and learn, not what you think about the birds and hope to learn. You are after facts first, after the facts it will be time to draw conclusions.

It is the purpose of this Bulletin and of the Club which it represents to help those who need help in their study of birds. The editor is always ready to answer questions through these pages, if the question is of general interest and application, or privately, if that seems the better course. Questions which would naturally lead to controversy cannot be given a place for the very obvious reason that our knowledge of the birds would not be increased thereby. We are learners direct from Nature.

Mr. Frank S. Daggett is now in Chicago, and will probably remain there for a year. Any of our members who may be in the city would meet a welcome from him at 241-2 Rialto Building.

Mr. Clarence F. Stone of Brockport, N. Y., has had remarkable success in quest of the nests of the Cerulean, Canadian, Black-throated Blue, and other rare warblers, and if he is blessed with the necessary leisure hours, we may hope for an illustrated article on the subject.

Mr. W. H. Brownson, who is news editor of the Portland, Maine, Daily Advertiser, writes a very attractive bird column once a week for that paper. In his long walks he sees and writes of many interesting species.

Mr. Harry B. McConnell, of Cadiz (Ohio) Republican, has occasional interesting and instructive articles about the birds, giving special attention to the reasons why we should do all in our power to protect them.

Mr. Walter F. Webb, now in the nursery business, is just as much of a bird enthusiastic as ever. He is planning to build a fine house with plenty of room for cabinets for his collections. He would be greatly pleased to meet any of our members who may find themselves in Rochester, New York.

Advance proof sheets of Gleanings From Nature No. III, The Haunts of the Golden-winged Warbler, by J. Warren Jacobs, give promise of another treat of this interesting series. The makeup is similar to the Martin Colony booklet. There will be three full page plates showing the haunts of the Golden-wing, three smaller plates with nests in natural position, and two on one page showing series of eggs, with parents, natural size. There will also be a color chart, and notes on migration, nest building, song, food, young, eggs, etc., thirty pages in all. Mr. Jacobs is a careful observer and knows how to record what he sees.

NOTES.

FROM BOULDER, COLORADO.

Our experience with House Finches (Carpodacus mexicanus frontalis) and Say Phoebes (Sayornis saya) may be of interest to your readers. Early in May a pair of the finches began building a nest of roots and mud in a 4x5 dry plate box placed under our front porch for their benefit. A few days later another pair began work on the same nest, which finally resulted in a pitched battle lasting for several hours, with brief intermissions. One pair then left and the other continued work. In a day or two a Phoebe put in an appearance and began work on the same nest, apparently working in harmony with the finches, for we never caught them in any altercation, though unquestionably both families worked on the nest. In two or three days the finches left Then the Phoebes built a fine symmetrical nest of hair and wool over the finches' rough foundation. In due time four Phoebe eggs appeared, then after three or four days incubation the Phoebes abandoned the nest, having apparently been kept from the eggs by visitors on the porch so long that the eggs got chilled. Nest and eggs are now in the Museum of the University of Colorado. JUNIUS HENDERSON.

HOPE, DICKINSON CO., KAS., January 25, 1904.

I am glad to note some new birds in this locality in the last two years. All first noticed in 1902 and again in 1903. The first was the Phoebe, three nests found in 1902 and twelve in 1903. Wood Thrush, three nests in 1902 and one in 1903, in park in town. Chimney Swift one pair nested in school house chimney in 1902 and three birds seen in spring of 1903, but did not nest as they only staid about a week and then left. The Phoebe was quite common last summer along the creeks, but only a few in 1902. I am sure they have not nested here before 1902, as I have been under the bridges where they nested every year since they were built, and never saw them or their nest until 1902. The Chimney Swifts are also the first pair ever noted in town; pointed them out to old eastern people and they said, "they were the first they ever had seen here but are quite common in eastern part of state" On January 5, 1904, as I was driving in the country I saw a Brown Thrasher hopping along a hedge fence, the first one of them I ever noticed in the winter here. O. H. PEASE.

PUBLICATIONS RECEIVED.

A Revision of the American Great Horned Owls. By Harry C. Oberholser. From the Proceedings of the United States National Museum, Vol. XXVII, pages 177-192. No. 1352.

A Review of the Wrens of the Genus Troglodytes. By Harry C. Oberholser. From the Proceedings of the U. S. National Museum, Vol. XXVII, pages 197-210. No. 1354.

In these two papers Mr. Oberholser has reviewed all American forms, not simply those which we call North American. We regard this broad-

ening of view which has been evident recently, a distinct gain in classification methods. It is certain to throw light upon many problems of distribution and relationship hitherto only suspected. The use of *Asio* instead of *Bubo* for the the genus of the Great Horned Owls will cause temporary confusion, but ultimately make for stability, we trust. Seventeen forms of the Great Horned Owl are recognized, six of which are new species. That does not seem excessive from the whole of the Americas, but we trust that the naming of forms may rest permanently here. In treating the wrens of the genus *Troglodytes*, Mr. Oberholser has made a new genus in which to place that curious form, *Thryorchilus browni* from Panama. Of the 37 forms of *Troglodytes* recognized there are 14 independent species showing no subspecific affinities. The South American form *musculus* is split into 14 forms, while our North American form *aedon*, remains triple. In the whole genus but three new subspecies are eloborated, none of which affect our fauna. We congratulate Mr. Oberholser upon this work, and trust that the whole field of American ornithology may be gone over as carefully in the near future. L. J.

THE BIRDS OF LICKING COUNTY, OHIO. By I. A. Field. Reprinted from The Bulletin Scientific Laboratories of Denison University, Vol. XII, December, 1903.

Mr. Field prefaces the annotated part of his catalogue with remarks upon the topography of the county and the general distribution of the birds found there. The annotations consist of statements concerning the times of appearance of the birds and the regions in which they may be found. Of the 203 species found in the county during the three years of his study, 27 are permanent residents, 79 summer residents, 9 winter residents, 80 transient visitants, and 8 accidental visitants. Of the accidental species the most interesting are the European Widgeon and Cinnamon Teal, both captured on Licking Reservoir. We welcome this additional faunal list as a contribution to the literature of distribution. L. J.

BOLL WEEVILS AND BIRDS. Address by Prof. H. P. Attwater, industrial agent Southern Pacific, at the Second Annual Convention of the Texas Cotton Growers' Association, Dallas, Texas, November 6th, 1903.

In this paper Prof. Attwater shows clearly that one of the greatest enemies of the Cotton Boll Weevil is the host of birds. He pleads for the protection and encouragement of the birds that they may go about their beneficent work of destruction of insect pests unhindered. It is not too much to expect that if the birds are allowed to increase as they would normally they will keep in check insect depredations, for they will eat such insects as are most easily secured, other things being equal. L. J.

POSTGLACIAL ORIGIN AND MIGRATION OF THE LIFE OF THE NORTHEASTERN UNITED STATES. By Charles C. Adams. Reprinted from

Journal of Geography, Vol. I, No. 7, September, 1902, pages 303-310, 352-357.

In this paper the author endeavors to trace the redistribution of Life in that part of North America which was covered with glacial ice. The first migration was by the arctic types, which pushed up against the border of the retreating ice; second by the subarctic life, following close upon the heels of the arctic, and the third the temperate, the last migration, represented by the forms now found in Ohio, Indiana and Illinois. Only the third class are typically American. The northward migrations were marked not by river courses so much as by forest and plains regions. We welcome this paper as throwing light upon the post-glacial origin of our flora and fauna. L. J.

SOUTHEASTERN UNITED STATES AS A CENTER OF GEOGRAPHICAL DISTRIBUTION OF FLORA AND FAUNA. By Charles C. Adams. Reprinted from Biological Bulletin, Vol. III, No. 3, July, 1902. Pages 116-131.

The author here shows that so far as the eastern United States is concerned, the post-glacial life has been distributed from the southeast, except the distinctly boreal forms, and still remains as a center of dispersal. He recognizes, also, a southwestern center of dispersal in the arid region of northern Mexico and the southwestern United States. LJ

Amateur Sportsman, Vol. XXX, Nos. 2, 3, 4.
American Ornithology, Vol. IV, Nos. 1, 2, 3.
Bird-Lore, Vol. VI, Nos. 1, 2.
Boll Weevels and Birds.
Cassinia, 1903.
Condor, The, Vol. IV, No. 1.
Journal of Applied Microscopy, Vol. VI, Nos. 9, 10.
Maine Sportsman, The, Vol. II, Nos. 125, 126.
Naturaliste Canadien, Le, Vol. XXX, Nos. 11, 12; Vol. XXXI, No. 1.
Nature Notes, Vol. XV, No. 170.

LIST OF MEMBERS.

Officers for 1904.

President—Lynds Jones, Oberlin, Ohio.
Vice-President—W. L. Dawson, Columbus, Ohio.
Secretary—John W. Daniel, Jr., 1794 Lanier Ave., N. W., Washington, D. C.
Treasurer—Frank L. Burns, Berwyn, Pa.
Executive Committee—Harry C. Oberholser, John H. Sage, Benj. T. Gault.

Active Members.

R. L. Baird, Oberlin, O.
A. C. Bent, Taunton, Mass.
Homer L. Bigelow, 511 Washington St , Boston, Mass.
Dr. Louis B. Bishop, 356 Orange St., New Haven, Conn.
A. W. Blain, Jr., 131 Elmwood Ave., Detroit, Mich.
Charles F. Brennan, Mt. Carmel, Ill.
W. H. Brownson, Advertiser office, Portland, Me.
Clarence Morrison Brooks, West Point, N. Y.
Frank Bruen, 84 Spring St., Bristol, Conn.
Henry R. Buck, Wethersfield, Conn.
G. M. Burdick, Milton, Wis.
Frank L. Burns, Berwyn, Pa.
Mrs. Agnes Chase, 59 Florida Ave., N. W., Washington, D. C.
Maunsell S. Crosby, Rhinebeck, N. Y.
Edward M. Daniel, Lynchburg, Va.
John W. Daniel, Jr., 1794 Lanier Ave., N. W., Washington, D. C.
Mrs. Elizabeth B. Davenport, 45 Green St., Brattleboro, Vt.
Rev. W. L. Dawson, Columbus, Ohio.
Walter Deane, 29 Brewster St., Cambridge, Mass.
Owen Durfee, Box 25, Fall River, Mass.
George C. Embody, 22 Walker Ave., Bradford, Pa.
Lou J. Eppinger, 516 Chene St., Detroit, Mich.
Miss E. W. Fisher, 1502 Pine St., Philadelphia, Pa.
Benj. T. Gault, Glen Ellyn, Ill.
Mrs. Stephen Cobb Goss, 5475 Ridgewood Court, Hyde Park, Ill.
Glen M. Hathorn, 703 B Ave., Cedar Rapids, Iowa.
Hon. Junius Henderson, Boulder, Colo.
Rev. W. F. Henninger, 206 Jefferson St., Tiffin, Ohio.
N. Hollister, Delavan, Wis.
LaRue K. Holmes, Summit, N. J.
Fred C. Hubel, 112 Alexandrine Ave., Detroit, Mich.
Chreswell J. Hunt, 1306 N 53rd St., Philadelphia, Pa.
J. Warren Jacobs, Waynesburg, Pa.
Lynds Jones, 160 North Professor St., Oberlin, Ohio.

J. Eugene Law, 305 Holt St., Pomono, Calif.
W. P. Lyons, 10-11, 41 E. Fourth St., Cincinnati, Ohio.
Miss Rebecca Morehead Leete, North East, Pa.
Herbert H. Marble, Box 14, Somerset, Mass.
Harry B. McConnell, Cadiz, Ohio.
C. H. Morris, McConnelsville, Ohio.
Mrs. Lida B. McMurray, DeKalb, Ill.
Mrs. Olive Thorn Miller, 827 DeKalb Ave., Brooklyn, N. Y.
Walton I. Mitchell, 534 Summit Ave., St. Paul, Minn.
Harry C. Oberholser, 1454 Sheridan Ave , Washington, D. C.
O. H. Pease, Hope, Kansas.
Harold E Porter, 101 West 85th St., New York City.
Charles H. Rogers, 5 West 82d St , New York City.
Dr. W. E. Rotzell, Narberth, Pa.
John H. Sage, Portland, Conn.
W. E. Saunders, Box 58, London, Ont.
Orpheus M. Schantz, Morton Park, Ill.
George H. Selleck, 110 High St., Exeter, N. H.
Dr. A. R. Shearer, Mont Belvieu, Texes.
Prof. P. M. Silloway, Lewiston, Mont.
H. H. Skinner, Oberlin, Ohio.
Robert Windsor Smith, Kirkwood, Ga.
Clarence J. Stone, Branchport, N. Y.
Bradshaw H. Swales, 46 Moffat Building, Detroit, Mich.
John E. Thayer, Box 98 Lancaster, Mass.
Miss Adelaide Utter, Clerk U. S. Circuit Court, Kansas City, Mo.
Warner M. Van Norden, 751 5th Ave., New York City.
Walter F. Webb, Grand Avenue, Rochester N. Y.
Alick Wetmore, North Freedom, Wis.
Burtis H. Wilson, 1102 17th St., Rock Island, Ill.
Sidney S. Wilson, Bank of Commerce Building, St. Joseph, Mo.

Honorary Members.

Prof. H. H. Ballard, Pittsfield, Mass.
Rev. J. H. Langille, Kensington, Md.
Prof. H. Nehrling, Palm Cottage, Gotha, Fla.
Mr. Otto Widmann, Old Orchard, Mo.

Associate Members.

Harold Bowditch, Pond St., Jamaica Plain, Mass.
H. W. Carriger, 2611½ Sutter St., San Francisco, Calif.
Mrs. T. D. Dershimer, Square Top, Wyoming Co., Pa.
Irving A. Field, Cambridge, Mass.
J. L. Floyd, Ada, Ohio.
N. C. Gilbert, DeKalb. Ill.
Harry S. Hathaway, Box 498, Providence, R. I.
Thos. D. Keim, Bristol, Pa.

Miss Caroline Mathews, 90 Main St., Waterville, Me.
William J. Mills, East Point, Ga.
Miss Isabella Sand, Ardsley-on-Hudson, N. Y.
Dr. Chas. E. Slocum, Defiance, O.
Miss Jane L. Spencer, 239 East 1st St., Oswego, N. Y.
Rev. George F. Weld, Lincoln St., Hingham, Mass.
Millard Van Wagner, 47 Leonard St., New York, N. Y.
Edward R. Warren, 20 West Caramillo St., Colorado Springs, Colo.
Norman A. Wood, 1216 S. U. Ave., Ann Arbor, Mich.

ELECTION OF NEW MEMBERS.

The following persons are nominated for Active membership:

Objections to the reception of any of these persons should be sent to the Secretary before April 15.

G. F. Richardson, 116 S. Main St., Mt. Pleasant, Mich.
Norman A. Wood, 1216 S. U. Ave., Ann Arbor, Mich.
C. H. Morris, McConnelsville, Ohio.
W. H. Brownson, Advertiser Office, Portland, Me.
LaRue K. Holmes, Summit, N. J.
Chreswell J. Hunt, 1306 N. 53rd St., Philadelphia, Pa.
Clarence F. Stone, Branchport, N. Y.
Walter F. Webb, Grand Ave., Rochester, N. Y.
Maunsell S. Crosby, Rhinebeck, N. Y.

Nominated for Associate membership:

G. F. Richardson, 116 S. Main St., Mt. Pleasant, Mich.

THE BEST
LAWN
SWING
MADE

AGENTS WANTED

Lawn Swings and Settees, Hammock Chairs, Camp Chairs and Stools, Ironing Tables, Wash Benches, Etc.

Agents easily make

$5 TO $10 PER DAY.

Will furnish samples at reduced prices to those desiring agency. Exclusive territory given. Address,

Clearfield Wooden-Ware Co.,
CLEARFIELD, PA.

DAWSON'S BIRDS OF OHIO.

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones. 720 quarto pages. 322 birds accurately described, their habits interestingly given. Combines ready identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations, to Wilson Bulletin, Oberlin, Ohio.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES.

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds of Ohio are treated. It does not attempt a description of the plumage. Price 50 cents. Address Lynds Jones, Oberlin, Ohio.

Wilson Chapter Publications

CONSIST OF TWO SERIES—OLD AND NEW

The Old Series comprises the following issues:

Ornithologists and Oologists' Semi-annual, Vols. I and II two numbers each; Vol. III, one number. (No. 1 of both Vols. I and II are out of print.) 25 cents a number.

Wilson Quarterly, Vol. IV, two numbers. 25c a number.

The Journal, two numbers. 10 cents a number.

The whole series (available numbers) $1.00.

The New Series comprises

Wilson Bulletins, from No. 1 to 46 inclusive. (Nos. 1, 2, 3, 4, 6, 8, and 24 are out of print.) Several numbers are almost gone.

Of the Wilson Bulletin, No. 5 is a Monograph of the Crow, 41 pages, by Frank L. Burns. Price 25 cents. No. 15 is a study of "The Oberlin Grackle Roost," 18 pages, by Lynds Jones. Price 20 cents. No. 30, "Warbler Songs," 56 pages, by Lynds Jones; in which all North American Warblers are discussed, the songs of nearly all described, and a field key to the adult males given. Price 50 cents. No. 31, "A Monograph of the Flicker," 32 pages, by Frank L. Burns. Price 50 cents. No. 38, "A Summer Reconnoissance in the West," by Lynds Jones and W. L. Dawson; being a study of the birds in fourteen states during a journey of 7000 miles. Price 20 cents. No. 37, "A Sectional Bird Census," by Frank L. Burns. Price 25 cents.

The other numbers consist of "General Notes." Price 15 cents each. The whole available new series for $4.50.

Address all communications to

LYNDS JONES, Oberlin, O.

Stanford University Library
JAN 10 1905

Old Series Vol. XVI. New Series Vol. XI, No. 2

The Wilson Bulletin

No 47

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

June, 1904

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.

CONTENTS

	PAGE
Some Birds of Olympia, Washington. J. M. Keck	33
All Day with the Birds	37
Oberlin all Day with the Birds. Lynds Jones	43
An 'All Day' Bird Study. Burtis H. Wilson	51
The Song of the Dickcissell. P. M. Silloway	52
Spring Migration in Lorain County, O., '04. Lynds Jones	55
A Door-Yard List from Morton Park, Ill. O. M. Schantz	57
Editorial	60
Publications Received	61

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, Ohio, edited by Lynds Jones.

Subscription: Fifty cents a year including postage, *strictly in advance.* Single numbers fifteen cents, unless they are "Special" numbers, when a special price is fixed. The Bulletin, including all "Special" numbers, is free to all paid up member, either Active, Associate or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, 1794 Lanier, Ave., N. W. Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for reviews, should be addressed to Lynds Jones, Oberlin, Ohio.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 20th ot the month preceding publication.

A typical nest of the White-eyed Vireo (*Vireo noveboracensis*).
One egg of the Cowbird appears above the lower edge of the nest.

(Courtesy of Dawson's Birds of Ohio)

THE

WILSON BULLETIN

NO. 47

A QUARTERLY JOURNAL OF ORNITHLOGY

VOL XVI. JUNE, 1904 NO. 2

SOME BIRDS OF OLYMPIA, WASHINGTON.

BY J. M. KECK.

Olympia is a favorite winter resort for many species of birds. The earth is rarely frozen, and both fresh and salt water is open all winter. The lowest temperature last winter was 25 degrees. There was but little snow, and at no time was the ground everywhere covered. Wild fruits abound and remain fresh until spring. In summer the birds are less abundant than in winter. The following list is the result of occasional observations from September 1 to May 5:

1. **Aechmophorus occidentalis.** Western Grebe. Common resident.
2. **Colymbus holbœlli.** Holbœll Grebe. Rare winter visitor.
3. **Colymbus auritus.** Horned Grebe. Migrant, common in spring.
4. **Colymbus nigricollis californicus.** Eared Grebe. Migrant. Seen in spring but not common.
5 **Podilymbus podiceps.** Pied-billed Grebe. Abundant resident. Very tame.
6. **Gavia imber.** Loon. Rare winter visitor.
7. **Gavia pacifica.** Pacific Loon. Rather common in winter.
8. **Gavia lumme.** Red-throated Loon. Common winter resident.
9. **Cerorhinca monocerata.** Rhinoceros Auklet. Rare winter visitor.
10. **Synthliborhamphus antiquus.** Ancient Murrlet, Very common winter resident.
11. **Cepphus columba.** Pigeon Guillemot. Resident, but not common.
12. **Larus glaucescens.** Glaucous-winged Gull. Abundant winter resident.
13. **Larus occidentalis.** Western Gull. Common resident.
14. **Larus argentatus.** Herring Gull. Abundant. Probably resident.
15. **Larus delawarensis.** Ring-billed Gull. Very common. Possibly resident.
16. **Larus brachyrhynchus.** Short-billed Gull. Not common winter resident.

17. **Larus philadelphia.** Bonaparte Gull. Abundant in winter. Possibly resident.

18. **Sterna paradisæa.** Arctic Tern. Common summer resident.

19. **Phalacrocorax pencillatus.** Brandt Cormorant. Common resident.

20. **Phalacrocorax pelagicus.** Pelagic Cormorant. Rare winter visitor.

21. **Merganser americanus.** American Merganser. Abundant in spring.

22. **Merganser serrator.** Red-breasted Merganser. Common in winter.

23. **Lophodytes cucullatus.** Hooded Merganser. Rather common resident.

24. **Anas boschas.** Mallard. Not common resident.

25. **Nettion carolinensis.** Green-winged Teal. Seen in spring. Rare.

26. **Spatula clypeata.** Shoveller. Rare resident.

27. **Aix sponsa.** Wood Duck. Rare resident.

28. **Aythya americana.** Redhead. Not common resident.

29. **Aythya vallisneria.** Canvas-back. Abundant in winter and tame.

30. **Aythya marila.** Scaup Duck. Not very common.

31. **Aythya affinis.** Lesser Scaup Duck. Abundant resident.

32. **Clangula clangula americana.** American Golden-eye. Abundant resident.

33. **Charitonetta albeola.** Buffle-head. Rather common resident.

34. **Harelda hyemalis.** Old-squaw. Rare winter visitor.

35. **Histrionicus histrionicus.** Harlequin Duck. A pair seen April 5.

36. **Oidemia deglandi.** White-winged Scoter. Common winter resident.

37. **Oidemia perspicillata.** Surf Scoter. Common winter resident. Tame.

38. **Chen hyperborea.** Lesser Snow Goose. Rare migrant.

39. **Anser albifrons gambeli.** White-fronted Goose. Not common migrant.

40. **Branta canadensis occidentalis.** White-cheeked Goose. Common migrant. As I have not had the opportunity of closely examining this goose, and as the books give several varieties of Canada geese for this region, I am not absolutely certain of this species. The hunters do not distinguish the different varieties of Canada geese. They report them nesting to some extent in this locality.

41. **Botaurus lentiginosus.** American Bittern. Summer resident.

42. **Ardea herodias.** Great Blue Heron. Resident. Rare in winter.

43. **Ardea virescens.** Green Heron. Not common summer resident.

44. **Nycticorax nycticorax nævius.** Black-crowned Night Heron. Summer resident.

45. **Fulica americana.** American Coot. Summer resident.

46. **Gallinago delicata.** Wilson Snipe. Summer resident. Returns in February.

47. **Ereunetes occidentalis.** Western Sandpiper. Migrant.

48. **Numenius longirostris.** Long-billed Curlew. Probably summer resident.

49. **Colinus virginianus.** Bob-white. Rare resident. Introduced.

50. **Lophortyx californicus.** California Partridge. Rare resident. Introduced.

51. **Bonasa umbellus togata.** Oregon Ruffed Grouse. Common resident.

52. **Phasianus torquatus.** Ring-necked Pheasant. Not common. Introduced.

53. **Columba fasciata.** Band-tailed Pigeon. Not common summer resident.

54. **Accipiter velox.** Sharp-shinned Hawk. Not very common resident.

55. **Aecipiter cooperii.** Cooper Hawk. Rare resident.

56. **Buteo borealis calurus.** Western Red-tail. Not common resident.

57. **Falco sparverius deserticola.** Desert Sparrow Hawk. Rare resident.

58. **Pandion haliaetus carolinensis.** Am. Osprey. Summer resident.

59. **Asio accipitrinus.** Short-eared Owl. Common resident.

60. **Megascops asio kennicottii.** Kennicott Screech Owl. Common resident.

61. **Bubo virginianus saturatus.** Dusky Horned Owl. Not common resident.

62. **Ceryle alcyon.** Belted Kingfisher. Resident. Rare in mid-winter.

63. **Dryobates villosus harrisii.** Harris Woodpecker. Not common resident.

64. **Dryobates pubescens gairdnerii.** Gairdner Woodpecker. Not common resident.

65. **Ceophlœus pileatus abieticola.** Northern Pileated Woodpecker. Rather common resident.

66. **Melanerpes torquatus.** Lewis Woodpecker. Rare. Probably summer resident.

67. **Colaptes cafer saturatior.** Northwestern Flicker. Common resident. A variety of the Red-shafted type.

68. **Chordeiles virginianus henryi.** Western Nighthawk. Summer resident.

69. **Selasphorus rufus.** Rufous Hummingbird. Abundant summer resident.

70. **Selasphorus alleni.** Allen Hummingbird. Rare summer resident.

71. **Cyanocitta stelleri.** Steller Jay. Resident. Very rare in winter.

72. **Perisoreus obscurus griseus.** Gray Jay. Rare resident.

73. **Corvus americanus.** American Crow. Resident. Rare before Feb. 5.

74. **Corvus caurinus.** Northwest Crow. Abundant resident.

75. **Agelaius phœniceus caurinus.** Northwestern Red-wing. Common resident. Rare in mid-winter.

76. **Sturnella magna neglecta.** Western Meadowlark. Common resident.

77. **Scolecophagus cyanocephalus.** Brewer Blackbird. Abundant resident. Very tame.

78. **Coccothraustes vespertinus montanus.** Western Evening Grosbeak. Abundant migrant. A flock of a dozen appeared Feb. 18. In a few days the flock increased to a hundred. About April 1 the flock decreased to about twenty. A few are still here May 5. They are very tame. They feed mainly on maple seeds.

79 **Carpodacus purpureus californicus.** California Purple Finch. Abundant resident.

80. **Acanthis linaria.** Redpoll. Rather common winter resident.

81. **Astragalinus tristis salicamans.** Willow Goldfinch. Common resident. Rare in mid-winter.

82. **Spinus pinus.** Pine Siskin. Common resident.

83. **Passer domesticus.** English Sparrow. Resident. Not so abundant as in eastern cities.

84. **Ammodramus sandwichensis alaudinus.** Western Savanna Sparrow. Summer resident.

85. **Zonotrichia leucophrys gambelli.** Gambel Sparrow. Migrant. Possibly summer resident.

86. **Zamelodia leucophrys nuttalli.** Nuttall Sparrow. Common summer resident.

87. **Zonotrichia coronata.** Golden-crowned Sparrow. Migrant.

88. **Spizella socialis arizonæ.** Western Chipping Sparrow. Common summer resident.

89. **Junco hyemalis oergonus.** Oregon Junco. Common winter resident.

90. **Melospiza melodia morphna.** Rusty Song Sparrow. Common resident.

91. **Passerella itiaca unalaschcensis.** Townsend Sparrow. Winter resident.

92. **Pipilo maculatus oregonus.** Oregon Towhee. Common resident.

93. **Piranga ludoviciana** Louisiana Tanager. Rare summer resident.

94. **Progne subis hesperia.** Western Martin. Common summer resident.

95. **Tachycineta thalassina lepida.** Northern Violet-green Swallow. Common summer resident.

96. **Riparia riparia.** Bank Swallow. Common summer resident.

97. **Ampelis cedrorum.** Cedar Waxwing. Rare resident.

98. **Helminthophila alata lutescens.** Lutescent Warbler. Summer resident.

99. **Dendroica æstiva.** Yellow Warbler. Not common summer resident.

100. **Dendroica coronata.** Myrtle Warbler. Migrant.

101. **Dendroica auduboni.** Audubon Warbler. Common resident.

102. **Dendroica nigrescens.** Black-throated Gray Warbler. Rare summer resident.

103. **Geothlypis tolmiei.** Macgillivray Warbler. Rare summer resident.

104. **Salpinctes obsoletus.** Rock Wren. Very rare resident.

105. Thryomanes bewickii calophonus. Northwest Bewick Wren. Not uncommon resident.

106. Troglodytes aedon parkmani. Pacific House Wren. Not common summer resident.

107. Olbiorchilus hiemalis pacificus. Western Winter Wren. Common resident.

108. Cistothorus palustris paludicola. Tule Wren. Rare summer resident.

109. Sitta canadensis. Red-breasted Nuthatch. Not common resident.

110. Parus atricapillus occidentalis. Oregon Chickadee. Common resident.

111. Psaltriparus minimus. Bush-Tit. Not common resident.

112. Regulus satrapa olivaceus. Western Golden-crowned Kinglet. Abundant winter resident.

113. Regulus calendula. Ruby-crowned Kinglet. Possibly partly resident. Common from Feb. 15 to May 1.

114. Hylocichla ustulata. Russet-backed Thrush. Rare summer resident.

115. Merula migratoria propinqua. Western Robin. Resident. Very rare in mid-winter. Abundant after Feb. 7.

116. Ixoreus nævius. Varied Thrush. Migrant. Common through March.

117. Sialia mexicana occidentalis. Western Bluebird. Resident. Very rare in mid-winter and not very common at any time.

ALL DAY WITH THE BIRDS.

Responses to the call for All Day studies of the birds during May show a gratifying interest in this intensive study of the migrations. It is not an easy task to spend an entire day, always on the keen edge, searching for birds, but no such effort goes unrewarded. One who has given much time to the study of the spring migrations hopes only to be able to spend more time during the next season. Constant study for three weeks or more, during which the birds are passing in the greatest numbers, might prove too wearing to even the most ardent student, but there are those of us who must learn by personal experience.

Reports received range from May 5 to May 23, and represent territory from New York City on the east to Rock Island, Ill., on the west, and latitude from Oberlin to Columbus, Ohio. The reports seem to prove that the height of the migrations for the region covered was between May 6 and 16, a

period of ten days. The Ohio and Illinois reports clearly indicate an exceptional abundance of birds, both in individuals and in species, for practically the whole period. At Oberlin the 21st was marked by a general departure of warblers and sparrows. But a discussion of the Northern Ohio migrations may be found elsewhere.

In the accompanying table, "A" indicates that the species was abundant, "C" that it was common, "T C" tolerably common, "F" few in numbers. "X" indicates that the species was recorded but the individuals not counted, and the numbers give the numbers of individuals seen when a count was made.

Explanations of the work done in the several places represented are given here together, so that the tabular arrangement may not be interfered with, and so that the reader may see at a glance the distribution so far as the reports give it. The Ohio reports are given precedence because they contain the largest number of species,

Oberlin, Ohio, May 9. Temperature 70 at 3 a. m., 54 at 9 a. m. Partly cloudy, clouding rapidly to a cold westerly rainstorm at 6:30 a. m., the rain ceasing in an hour, and afterward a gradual lightening of the clouds, with a brief show of the sun at 5 p. m. Wind westerly, heavy to strong. Time, 3 a. m. to 6:30 p. m. Oak Point and vicinity until 3:30 p. m., then a wheel ride twelve miles to Oberlin and two hours' work in the woods and fields northwest of Oberlin.

LYNDS JONES.

Tiffin, Ohio, May 10. Temperature (at Oberlin) 45 to 62. Wind west-south west, brisk. Showers 9:30 to 10 a. m., 1:30 to 2:30, 3:10 to 4:20 p. m. 4:30 to 10:15 a. m., fields. meadows and two woods one and one-half miles southeast of Tiffin; 11:15 a. m. to 1:30 p. m., due north of Tiffin along the Big Four tracks and a deserted reservoir, three miles; west one mile, 2:30 to 3:10 p. m.; north one mile and back along the Sandusky River, 4:30 to 5:20 p. m.; drive home across country, reaching home at 7:20 p. m.

REV. W. F. HENNINGER AND CARL HEILMANN.

Cadiz, Ohio, May 8. Temperature 60 to 70. Cloudy, with occasional light showers. Wind strong in the afternoon. Beginning at 4 a. m., time in field 11 hours. Walked fifteen miles, rode wheel fifteen miles. In woods and orchards around Cadiz.

HARRY B. MCCONNELL.

Columbus, Ohio, May 23. Temperature between 65 and 80. Clear and still in the forenoon, cloudy with light southwest wind afternoon. 3:00 a. m. to 9:30 p. m. Locality west of Columbus, and on the Ohio State University campus. Z. P. METCALF.

Morton Park, Illinois, May 8. Day warm and cloudy with slight showers. Wind southerly. 5 a. m. to 1 p. m. ORPHEUS M. SCHANTZ

Rock Island, Illinois, May 14. Morning overcoat, cold, warming about noon. No wind. 5 a. m. to 7 p. m. About twelve miles south of Rock Island, Ill., including a drive to Milan. A cultivated region, with woods, thickets, fields and streams. BURTIS H. WILSON.

A TWO-DAY CENSUS.

Northern New Jersey. 6:15 p. m. May 6th to 4:55 p. m. May 8th.

May 6th. Station to house by wagon; 6½ miles; 1½ hours; 6:15 to 7:45. Weather mostly cloudy. 10 species, all seen later.

May 7th. Weather cloudy. 12 species heard from room before leaving it, all seen later.

5.05 a. m. to 8:05 a. m. Weather partly cloudy; 60 degrees at return. Partly in woods, partly along road past fields and a few houses. 41 species, including Yellow-throated Vireo and Chestnut-sided and Black-throated Green Warblers, not seen later

8:55 a. m. to 6:40 p. m. Weather partly cloudy. I walked up a creek till it dwindled to a brook, came back by road, and spent some time in higher woods. The hunt included low and high woods, fields, a creek, and a village. 54 species (59 for the day).

May 8th. 4:50 a. m. to 9:00 a. m. Weather, first half, foggy; second half, mostly fine. Entirely a road hunt: through fields and past farmhouses, a village, and an occasional small wood. 39 species, including Sparrow Hawk (1), Orchard Oriole (2), Scarlet Tanager (2), Red-eyed Vireo (4), House Wren (1), and White-breasted Nuthatch (4) not mentioned before.

9:50 a. m. to 1:30 p. m. Weather fine; 72 degrees at start, 77 degrees at return. Like the long hunt of preceding day, but not so extended. 38 species, none seen this hunt only, except Hairy Woodpecker (2.)

50 minutes spent near house, beginning at 2:30 p. m. yielded six species. 35 of these minutes were in woods, where we saw only several Swifts.

House to station, mostly by wagon; 4 miles; 3:35 to 4:35. 22 species, including Heron, species unknown (1), Mourning Dove, Bob-white (1 pair), and Bank Swallow not seen before, making me 68 species for the two days.

I was alone until Saturday night, (the 7th), when Mr. G. E. Hix joined me, and we were together from then on. He had hunted that afternoon, seeing Marsh Hawk and Cliff Swallow. the only ones seen.

The most striking thing about these hunts is the scarcity of migrants. The Warbler swarms which should have been met with every little way in the woods were almost entirely absent on Saturday and wholly so on Sunday. CHARLES H. ROGERS.

West Chester County, N. Y., May 5. Temperature 75. Clear. 5 a. m. to 7:30 p. m. FREDERICK C. HUBEL.

Cupola, Chester County, Penn., May 8. Weather clear and warm. Peach and cherry trees in full bloom, apple buds just bursting. 5 a. m. till dark. A 200 acre farm, with a creek and woods, carefully worked. CHRESWELL J. HUNT.

	1 Oberlin, Ohio May 9	2 Tiffin, Ohio May 10	3 Cadiz, Ohio May 8	4 Columbus, O. May 23	5 Morton Park, Ill., May 8	6 Rock Island, Ill., May 14	7 West Chester N. Y., May 5	8 Nor. N. J. May 6, 6:15 p.m. to May 8, 8:55 p. m.	9 Cupola, Chester Co. Pa., May 8
Pied-billed Grebe	....	....	....	....	....	....	2	....	....
Herring Gull	2	....	....	....	....	....	50	....	....
Mallard	1	....	....	....	....	....	....	....	....
Great Blue Heron	....	1	....	1	....	x	....	....	....
Green Heron	8	7	....	x	....	....	....	....	x
Virginia Rail	3	....	....	....	....	....	....	....	....
Sora	2	....	....	....	....	....	....	....	....
American Woodcock	3	....	....	....	....	....	....	....	....
Wilson Snipe	7	....	....	....	....	....	....	....	....
Pectoral Sandpiper	....	2	....	....	....	....	....	....	....
Baird Sandpiper (?)	....	2	....	....	....	....	....	....	....
Least Sandpiper	10	....	....	....	....	....	....	....	....
Red-backed Sandpiper	....	2	....	....	....	....	....	....	....
Greater Yellow-legs	2	1	....	....	x	....	....	....	....
Yellow-legs	3	4	....	x	....	....	....	....	....
Solitary Sandpiper	c	4	....	....	....	....	....	2	....
Bartramian Sandpiper	2	2	....	....	....	....	....	....	x
Spotted Sandpiper	c	11	x	x	x	....	....	6	x
Killdeer	c	3	x	x	....	....	....	....	x
Bob-white	2	2	4	x	x	x	1	2	x
Ruffed Grouse	....	....	1	....	....	....	....	1	....
Mourning Dove	c	7	x	x	x	x	2	1	x
Turkey Vulture	....	1	....	x	....	....	....	1	x
Marsh Hawk	....	....	....	....	....	....	2	1	....
Sharp-shinned Hawk	....	....	....	x	....	....	1	....	....
Cooper Hawk	1	....	1	x	....	....	1	....	....
Red-tailed Hawk	2	1	1	x	....	x	2	....	....
Red-shouldered Hawk	2	....	....	x	....	....	8	1	....
Broad-winged Hawk	1	....	....	....	....	....	1	....	....
Bald Eagle	....	....	....	....	....	....	....	1	....
Am. Sparrow Hawk	3	2	....	x	....	....	....	1	x
Am. Osprey	....	....	....	....	....	....	3	....	....
Barred Owl	2	....	1	....	....	....	1	....	....
Screech Owl	1	1	....	1	....	x	2	....	....
Great Horned Owl	1	....	1	1	....	....	....	....	....
Yellow-billed Cuckoo	....	1	....	x	x	....	3	....	....
Black-billed Cuckoo	4	1	....	....	....	....	....	1	....
Belted Kingfisher	2	1	x	x	....	....	7	....	x
Hairy Woodpecker	1	1	x	x	....	x	4	1	....
Downy Woodpecker	4	5	x	x	x	x	7	1	x
Yellow-bellied Sapsucker	1	....	....	....	x	....	7	....	....
Red-headed Woodpecker	c	12	x	x	x	x	1	....	x
Red-bellied Woodpecker	....	1	x	x	....	x	....	....	....
Northern Flicker	c	8	x	x	x	x	12	5	x
Whippoorwill	7	1	....	....	....	x	....	....	....
Nighthawk	2	2	....	x	....	x	1	....	x
Chimney Swift	c	30	x	x	x	x	A	50	x
Ruby-throated Hummingbird	3	1	2	x	....	x	....	....	....
Kingbird	6	2	x	x	....	x	3	4	x
Crested Flycatcher	c	1	x	x	....	x	1	3	x

	1	2	3	4	5	6	7	8	9
Phoebe	c	5	x	x	x	x	14	8	x
Wood Pewee	c	6	x	x	x	x	11		x
Yellow bellied Flycatcher	1								
Green-crested Flycatcher	5	3							
Alder Flycatcher	7	4		x		x			
Least Flycatcher	c	1	x		x		6	4	
Prairie Horned Lark	4	3		x		x			
Blue Jay	c	3	x	x	x	x	12	8	x
Am. Crow	c	11	x	x	x	x	75	27	x
Bobolink	c	45	x	x	x	x	26	3	
Cowbird	c	15	x	x	x	x	4	4	x
Red-winged Blackbird	c	10	x	x	x	x	80	8	x
Meadowlark	c	35	x	x	x	x	23	13	x
Orchard Oriole	6	2	x	x		x		2	
Baltimore Oriole	c	20	x	x	x	x	9	3	x
Purple Grackle								2	x
Bronzed Grackle	c	10	x	x	x	x	8		
Purple Finch	2							16	
Am. Goldfinch	c	8	x	x		x	1	26	x
Vesper Sparrow	c	12	x	x			7	3	x
Savanna Sparrow		1							
Grasshopper Sparrow	4	9	x	x	x			4	x
Henslow Sparrow									x
Lark Sparrow						x			
White crowned Sparrow	c	4	2	1	x		3		
White-throated Sparrow	c	6	x	3	x	x		10	
Tree Sparrow		1					4		
Chipping Sparrow	c	10	x	x		x	40	12	
Field Sparrow	c	2	x	x	x	x	20	7	x
Slate-colored Junco							4	1	
Song Sparrow	c	30	x	x	x	x	70	25	x
Swamp Sparrow	c					x		7	
Towhee	c	4	x	x	x	x	2	2	x
Cardinal	3	6	x	x	x				x
Rose breasted Grosbeak	10	3	x	1	x	x			
Indigo Bunting	c	6	x	x	x	x			
Dickcissel	2	1		x		x			
Scarlet Tanager	c	5	x	x	x	x	1	2	
Purple Martin	5	3		x	x	1	40		
Cliff Swallow	tc			x		x	16	1	
Barn Swallow	c	6	x	x		x	20	22	x
Tree Swallow	2						5		
Bank Swallow	c	3	x	1		x		2	
Rough-winged Swallow	tc	10					1		
Cedar Waxwing	5								
Migrant Shrike	1	1				x			
Red-eyed Vireo	c	3	x	x	x	x		4	x
Warbling Vireo	c	8	x	x					
Yellow-throated Vireo	tc	4	2	x		x			
Blue he ded Vireo	tc	2	1						
White-eyed Vireo					x			4	x
Bell Vireo						x			
Black and White Warbler	c	1	1	x				6	x
Prothonotary Warbler	1				x				
Blue-winged Warbler		6				x		2	x
Golden-winged Warbler		1							
Nashville Warbler	c					x			
Orange-crowned Warbler	1	1				x			

	1	2	3	4	5	6	7	8	9
Tennessee Warbler	c	2	x		....	x	....	....	....
Northern Parula Warbler	....	6	....	....	....	....	....	5	....
Yellow Warbler	c	20	x	x	x	x	4	8	x
Black-throated Blue Warbler	c	1	2	x	x	....	....	....	....
Myrtle Warbler	c	5	3	....	x	....	....	6	x
Magnolia Warbler	c	15	x	x	x	x	....	....	....
Cerulean Warbler	1	1	3	1	....	....	....	....	....
Chestnut-sided Warbler	c	2	x	....	x	x	....	....	....
Bay-breasted Warbler	t c	2	....	x	x	x	....	....	....
Black-poll Warbler	1	6	....	x	x	x	....	....	x
Blackburnian Warbler	c	1	2	x	....	x	1	1	....
Black-throated Green Warbler	c	2	x	....	x	x	....	....	....
Kirtland Warbler	1	....	....	....	....	....	....	....	....
Pine Warbler	....	2	....	.	....	....	....	....	....
Palm Warbler	....	2	....	....	x	....	....	....	....
Prairie Warbler	1	....	....	. .	....	....	....	....	....
Oven-bird	c	3	x	x	x	x	....	6	x
Water-Thrush	7	....	1	....	....	x	....	2	....
Louisiana Water-Thrush	2	1	x	x	....	....	....	....	....
Kentucky Warbler	....	....	x	3	....	....	....	....	....
Connecticut Warbler	2	....	....	....	....	....	....	....	....
Mourning Warbler	t c	1	....	1	....	....	....	....	....
Northern Yellow-throat	c	8	1	x	x	x	....	10	x
Yellow-breasted Chat	t c	5	x	x	....	....	....	2	x
Hooded Warbler	1	....	....	....	....	....	....	....	....
Wilson Warbler	1	1	....	2	....	....	....	....	....
Canadian Warbler	t c	2	2	....	....	....	....	....	x
Am. Redstart	c	11	x	x	x	x	....	7	x
Am Pipit	c	....	....	....	....	....	....	35	....
Catbird	c	50	x	x	x	x	18	18	x
Brown Thrasher	t c	2	x	x	x	x	2	1	x
Carolina Wren	....	....	1	x	....	....	....	....	....
House Wren	c	16	....	x	....	x	10	1	x
Winter Wren	1	....	1	....	....	....	....	....	....
Long-billed Marsh Wren	2	....	....	....	....	....	....	....	....
Brown Creeper	....	....	....	....	....	....	1	....	....
White-breasted Nuthatch	5	2	x	x	....	x	8	4	x
Red-breasted Nuthatch	c	....	....	....	....	....	15	....	....
Tufted Titmouse	5	5	x	x	....	....	....	4	....
Chickadee	7	2	x	x	x	x	20	....	....
Golden-crowned Kinglet	....	4	....	....	....	....	1	....	....
Ruby-crowned Kinglet	t c	1	....	x	x	x	....	....	....
Blue-gray Gnatcatcher	c	20	....	x	....	....	....	....	....
Wood Thrush	c	9	x	x	x	x	16	5	x
Wilson Thrush	c	2	x	....	....	x	2	2	....
Gray-cheeked Thrush	c	....	....	....	x	x	....	....	....
Olive-backed Thrush	c	1	x	....	x	x	....	....	....
Hermit Thrush	2	1	....	....	....	....	1	....	....
Am Robin	c	50	x	x	x	x	A	50	x
Bluebird	c	6	x	x	x	x	18	9	x
Totals	128	112	80	90	56	76	65	66	53

OBERLIN ALL DAY WITH THE BIRDS. 1904.

Two somewhat different styles of "All Day" work were attempted during the past remarkable May migrations. The one followed the general lines of previous work, for the purpose of determining what one man, or two working as one, could do in a single day in recording the birds of this region; the other attempted to cover practically all the different regions and sorts of topography and physical features by working in companies of two persons each, the combined lists so made determining the day's record. This work may be called the "Company All Day" record. It is clear that such cooperative work would result in a much larger list, given the same conditions, than individual work, provided the individual be unable to cover the same territory in a day. Two elements enter into individual work of this sort which seriously hinder the searching of certain haunts of some birds; these are the time available in a single day, and physical endurance. If one could be certain what species were to be found in certain regions it would be possible to clean such region up and pass on to the next, but since this is almost never true, he must determine how much time it will be profitable to spend here and there and yonder, seeking to make each stop count for the most. There must be no actual stop in the lookout for species, for at no time and in no place out of doors is there no possibility of making some new record for the day. One must be on the keen edge constantly.

It has already been hinted that the present May migrations have been remarkable in many ways, in northern Ohio, if not elsewhere. Elsewhere will be found an article which attempts to give some conception of the conditions prevailing in Lorain County during April and May.

For the sake of continuity the "Company All Day" on May 7 is given first. The accompanying table groups the participants, giving to each group credit for the records made. In the same table will be found the writer's personal record, two days later, and two records by Mr. Dawson and the writer, at later dates, when foliage and the progress of the migrations proved unfavorable for more than ordinary lists. The total of 131 species for the "Company All Day," and of

128 for the writer's personal record fully attest the crowded character of this period of the migrations. It will be noticed that no less than 18 species were recorded later which should have been recorded by the company, and that seven other species were certainly in the region covered but were not seen on that day. Some of these would almost certainly have been found had the writer's original plan been carried out, but circumstances made that impossible.

The first group, consisting of Mr. R. L. Baird and Mr. S. D. Morrill, spent Friday night at Oak Point, in order to be ready for work at earliest dawn. Their work included a careful survey of the lake and its shore line in the immediate vicinity of Oak Point, the marshes at the mouth of Beaver Creek in the regions of slack water, and the woods and fields within a radius of a mile or less with the Oak Point grounds as a center. Mr. Baird visited Chance Creek during the evening, but without success.

Messrs. D. E. Nye and H. H. Skinner composed the second group. Mr. Nye spent the early morning in the gorges of Black River at Elyria, while Mr. Skinner spent the early morning hours in the Black Swamp woods, three miles northeast of Oberlin, later meeting Mr. Nye in Elyria. After spending some time with the warblers in the woods in that vicinity, they took trolley to Lorain, and finally to Oak Point, supplementing the work there of the first party, returning home, with Mr. Morrill, by trolley.

The third group, Mr. Harold Vincent and the writer, spent the early morning in the old South Woods, a mile south of Oberlin, and since neither could continue the work afternoon, they practically exhausted the possibilities of that woods before leaving it at ten o'clock, visiting an orchard in the outskirts of town, the water-works reservoir, Arboretum and cemetery, finally returning with a list of exactly 90 species for the morning. During the afternoon two more species were accidentally added. The original plan of this third group contemplated a visit, during the afternoon, to a region where Broad-winged Hawks, Ruffed Grouse, Barred Owls and Carolina Wrens were nesting, none of which were seen during the day.

The accompanying table clearly indicates that the Oak Point region is by far the richest single region of the vicinity. This is more clearly proven when it is known that of the writer's personal list of 128 species 120 were recorded at Oak Point. Here are combined lake shore, marsh, muddy flats, wet woods, open woods, heavy woods, brushy tangles, rolling fields and flat wet fields. Farm houses and extensive orchards entice some birds which usually remain several miles inland.

The day was more nearly ideal than any succeeding day when work of this sort was attempted. The temperature ranged from 57 degrees at 3 o'clock in the morning to 80 degrees during the middle of the day. The air was decidedly chilly during the early morning hours, but the sky was clear until near noon, when threatening clouds appeared, later breaking away to fair. The wind was s. e., light to brisk, dying away in the evening

Species marked A were abundant, C, common, T. C. tolerably common, F, few in numbers Figures indicate the number seen.

May 9th was a very different day. Three o'clock in the morning found the writer at Oak Point, a half-hour too early for the first bird, which proved to be the only Screech Owl of the day. When the Whip-poor-wills began, a few minutes later, the Nighthawks, Cardinals, Catbirds and Field Sparrows were aroused for their first sleepy songs. Within half an hour most birds were singing or calling. At 1:30 a. m. the sky was half cloudy, with a brisk southwest wind to help push the wheel over the dozen miles of road. At 6:30 the wind had shifted to west by south and increased to heavy, driving dark clouds before it, accompanied by a dash of fine rain and a perceptible drop in the temperature. By 10 o'clock the temperature had dropped from 70 degrees at the starting to 54 degrees, the heavy wind and dark sky continuing, but without rain At the approach of the storm all birds dropped from the tree-tops or higher branches to the ground, or near it, and remained low down during the day. They lost their timidity, most warblers permitting an approach within ten feet or less. It was during the last of the fine rain that a fine male of Kirtland Warbler stopped near me and remained within twenty feet, often within eight, for at least ten min-

utes, but he refused to sing. Later the rare Prairie and about noon the still rarer Hooded, were found in the low foliage. In a swampy woods half a mile south of Oak Point, the first Prothonotary Warbler for Lorain County was feeding and singing, permitting a close approach and excellent study. This would have amply atoned for the cold and wet of the day, but a little later a Yellow-bellied Flycatcher, which has eluded me for a dozen years, darted into view from a fringing thicket of the woods. Many other birds which do not ordinarily seek the shelter of the woods were found in the lea of the thickets, where the wind did not reach them. The abundance of all thrushes except the Hermit, exceeded anything which the county has seen during my stay in it. The weather seemed impossible, but since the birds were here the record depended only upon field work under adverse conditions.

The evening of May 13th did not give promise of anything worthy during the following morning, but a determination born of necessity found expression in a trolley ride to Oak Point during that evening, and a search for the soft side of a reasonably clean plank, in a cold, bleak drizzle. The plank proved to be a convenient broken-down bedstead, with dilapidated springs—and mattress, under cover! Being too case-hardened to fear microbes, we slept peacefully until early morning twilight. Clearly, the weather had gone mad, but during its few lucid intervals the birds sang and sported about in spite of rain and cold. The day began at 45 degrees and did not reach 60 during even the noon hour. Rain fell rather more than half the day. The walk overland to Oberlin added but five species to the day's list, thus equalling the largest list previous to this year—113 species.

Not satisfied with the previous record, Monday, May 16, found us again in the field, reversing the course of study by beginning, as in previous years, at the old South Woods at three in the morning. Leaving this usually favored place shortly after 7 o'clock, and home an hour later, we wheeled leisurely to Oak Point, making many stops by the way. The list had reached 94 when the wheels were cached at 11:30, a mile south of the lake. Here the physical strain of two days before began to affect our ability to do effective work. The keen edge had worn off. Shortly after dinner Dawson was

obliged to move homeward. Rather hasty work in the two swamps which had not yet been touched, a brief visit to the lake shore, and a hurried search through a half mile of woods, closed the day at 5:30. Dawson had recorded Cooper Hawk on the way home, making the combined list reach 114. The morning opened with a light frost in low places, but clear until late in the afternoon. The temperature reached 60 degrees during the day, with a brisk westerly wind dropping completely down before night. The foliage seriously interfered with rapid identification, but was little more troublesome than on the 14th. Up to the 11th neither blossoms nor leaves were far enough advanced to give trouble.

The work of the 16th closed a series of record breaking all day studies. While it was serious work, sapping the vitality, it yet gave full value in return in showing the exceptional character of the season of migration. Three times within ten days the previous best record was broken, and the fourth day equalled. A new high-water mark was made, which nothing but another exceptional season can hope to equal, while a new record for co-operative work has been established as the beginning of a long series of such studies, we may well hope. A series of such co-operative studies in any given region can hardly fail to throw light upon some of the vexing questions of local migration, and furnish material for the broader study when made in connection with similar studies in contiguous regions.

	May 7. Company.			May 9	May 14	May 16
	Baird Morrill	Nye Skinner	Jones Vincent	Lynds Jones	Dawson Jones	Dawson Jones
Pied-billed Grebe	1	1	.....			.....
Herring Gull	1	1	.....	2	7	4
Bonaparte Gull	.....	.....	.....	.....	1	.....
Common Tern	2	10	.....	.....	.....	12
Black Tern	.....	.....	.....	.....	2	.....
Red-breasted Merganser	1	.....	.....	.....	.....	.....
Mallard	.....	.....	.....	1	.....	1
Lesser Scaup Duck	1	1	.....	.....	.....	.....
American Bittern	4	1	.....	.....	.....	.....
Least Bittern	.....	.....	.....	.....	.....	1
Green Heron	2	1	.....	3	.....	3
King Rail	1	.....	.....	.....	.....	1
Virginia Rail	3	1	.....	3	4	2
Sora	1	.....	.....	2	4	5
American Woodcock	3	.....	2	3	5	3
Wilson Snipe	1	.....	.....	7	.....	1
Least Sandpiper	.....	.....	.....	10	.....	.....
Greater Yellow-legs	1	.....	.....	2	.....	1
Yellow-legs	3	.....	.....	3	1	.....
Solitary Sandpiper	f e	2	1	c	c	c
Batramian Sandpiper	.....	.....	1	2	1	2
Spotted Sandpiper	t c	c	2	c	c	c
Killdeer	t c	3	.....	c	c	c
Bob-white	f e	2		2	1	1
Mourning Dove	t c	t c	c	c	c	c
Turkey Vulture	3	1	.....	.....	.....	1
Marsh Hawk	1	.....	.....	.....	.....	.....
Sharp-shinned Hawk	1	1	.....	.....	1	.....
Cooper Hawk	.....	.....	.....	1	1	1
Red-tailed Hawk	1	.....	.....	2	1	.....
Red shouldered Hawk	.....	1	2	2	2	2
Broad-winged Hawk	.....	.....	.....	1	.....	.....
Am. Rough-legged Hawk	1	.....	.....	.....	.....	.....
Bald Eagle	2	.....	.....	.....	I	.....
Am Sparrow Hawk	f e	1	3	3	3	1
Am Osprey	1	f	.....	.....	.....	.....
Barred Owl	.....	.....	.....	2	1	1
Screech Owl	.....	t	.....	1	.....	
Great Horned Owl	1		.....	1	1	.....
Yellow-billed Cuckoo	1	1	.....	.....	.....	.....
Black-billed Cuckoo	1	.....	2	4	.....	.....
Belted Kingfisher	f e	2	1	2	1	.....
Hairy Woodpecker	1	2	2	1	3	5
Downy Woodpecker	f e	2	3	4	t c	t c
Yellow-bellied Sapsucker	.....	t c	.....	1	.....	.....
Red-headed Woodpecker	c	t c	c	c	c	c
Red-bellied Woodpecker	.....	1	.....	.....	1	.....
Northern Flicker	c	5	c	c	c	c
Whippoorwill	c	1	1	7	10	1
Nighthawk	.....	.....	.....	2	2	.....

	May 7	May 7	May 7	May 9	May 14	May 16
Chimney Swift	A	c	c	c	c	c
Ruby-throated Hummingbird	2	2	.....	3	3	3
Kingbird	f e	t c	c	6	t c	c
Crested Flycatcher	c	t c	2	c	c	c
Phoebe	f c	t c	2	c	c	c
Wood Pewee	f e	t c	x	c	c	c
Yellow-bellied Flycatcher	.....	.....	.....	1	.....	.....
Green-crested Flycatcher	f e	2	c	5	F	t c
Alder Flycatcher	f e	.....	1	7	F	F
Least Flycatcher	t c	t c	c	c	c	c
Prairie Horned Lark	t c	t c	4	4	F	F
Blue Jay	c	3	c	c	c	c
American Crow	c	6	c	c	c	c
Bobolink	A	c	c	c	c	c
Cowbird	c	t c	c	c	c	c
Red-winged Blackbird	A	c	c	c	c	c
Meadowlark	c	c	c	c	c	c
Orchard Oriole	1	1	4	6	F	F
Baltimore Oriole	f c	c	c	c	c	c
Bronzed Grackle	c	c	c	c	c	c
Purple Finch	.....	.....	x	2	.....	.....
American Goldfinch	c	t c	c	c	c	c
Vesper Sparrow	c	c	c	c	c	c
Grasshopper Sparrow	c	3	10	4	4	F
White-crowned Sparrow	15	7	.....	c	c	c
White-throated Sparrow	t c	3	c	c	c	c
Chipping Sparrow	f e	c	c	c	c	c
Field Sparrow	t c	c	c	c	c	c
Song Sparrow	c	c	c	c	c	c
Lincoln Sparrow	.....	.....	.....	.....	.....	1
Swamp Sparrow	t c	2	8	c	c	c
Towhee	t c	c	c	c	c	c
Cardinal	f e	3	2	3	t c	t c
Rose-breasted Grosbeak	f e	t c	x	10	t c	t c
Indigo Bunting	f e	2	c	c	c	c
Dickcissel	.....	.....	.....	2	.....	.....
Scarlet Tanager	f e	t c	c	c	c	c
Purple Martin	1	1	2	5	F	F
Cliff Swallow	f e	1	.....	t c	F	F
Barn Swallow	A	c	c	c	c	c
Tree Swallow	.....	.....	.....	2	F	F
Bank Swallow	A	2	.....	c	c	c
Rough-winged Swallow	A	3	.....	t c	t c	t c
Cedar Waxwing	.....	7	.....	5	7	20
Migrant Shrike	1	.....	.....	1	1	1
Red-eyed Vireo	.....	.....	c	c	c	c
Warbling Vireo	c	2	c	c	c	c
Yellow-throated Vireo	f e	3	x	t c	t c	t c
Blue-headed Vireo	f e	5	x	t c	t c	F
Black and White Warbler	f e	t c	c	c	F	F
Prothonotary Warbler	.....	.....	.....	1	1	.....
Blue-winged Warbler	2	c	x	.....		F
Golden-winged Warbler	.....	.....	1	.....	.....	.....
Nashville Warbler	t c	t c	c	c	c	c
Orange-crowned Warbler	.....	.....	.....	1	1	.....
Tennessee Warbler	.....	t c	c	c	c	c

	May 7	May 7	May 7	May 9	May 14	May 16
Northern Parula Warbler	.....	.....	.....	.....	1	.....
Cape May Warbler	.....	.....	.....	.....	.....	2
Yellow Warbler	c	c	c	c	c	c
Black-throated Blue Warbler	tc	c	c	c	c	c
Myrtle Warbler	tc	c	c	c	tc	F
Magnolia Warbler	c	c	c	c	c	c
Cerulean Warbler	.....	c	c	1	.....	c
Chestnut-sided Warbler	c	c	c	c	c	c
Bay-breasted Warbler	1	5	tc	tc	f	4
Black-poll Warbler	2	.....	.....	1	.....	.....
Blackburnian Warbler	c	c	c	c	c	c
Black-throated Green Warbler	c	c	c	c	c	tc
Kirtland Warbler	.....	.....	.....	1	.....	.....
Palm Warbler	c	2	1	.....	.....	.....
Prairie Warbler	.....	.....	.....	1	.....	.....
Oven-bird	c	c	c	c	c	c
Water-Thrush	6†	.....	2	7	4	6
Louisiana Water-Thrush	.....	1	.....	2	4	2
Connecticut Warbler	.....	2	1	2	.....	.....
Mourning Warbler	.....	.....	1	tc	.....	1
Northern Yellow-throat	c	c	c	c	c	c
Yellow-breasted Chat	2	1	2	7	tc	tc
Hooded Warbler	.....	.....	.....	1	.....	.....
Wilson Warbler	fe	3	1	1	1	1
Canadian Warbler	tc	3	2	tc	tc	F
American Redstart	c	c	c	c	c	c
American Pipit	5†	.....	c	c	c	tc
Catbird	tc	c	c	c	c	c
Brown Thrasher	fe	3	tc	tc	tc	tc
House Wren	2	c	c	c	c	c
Winter Wren	5	.....	.....	1	.....	.....
Short-billed Marsh Wren	.....	.....	.....	.....	2	2
Long-billed Marsh Wren	5†	.....	.....	2	4	4
White-breasted Nuthatch	fe	3	f	5	f	F
Red-breasted Nuthatch	tc	tc	2	c	c	tc
Tufted Titmouse	tc	3	4	5	f	F
Chickadee	.....	3	4	7	f	F
Ruby-crowned Kinglet	fe	.....	c	c	.....	2
Blue-gray Gnatcatcher	fe	1	c	c	tc	tc
Wood Thrush	tc	c	c	c	c	c
Wilson Thrush	c	c	c	c	c	c
Gray-cheeked Thrush	1	3	c	c	c	c
Olive-backed Thrush	A	c	c	c	c	c
Hermit Thrush	.....	1	2	2	.....	.....
American Robin	c	A	c	c	c	c
Bluebird	fe	c	c	c	c	c
Totals	115	105	92	128	113	114
Grand Total		131				

Total for May 7 to 16, inclusive, 152.

AN "ALL DAY" BIRD STUDY.

BY BURTIS H. WILSON.

I left the house on the morning of the 15th, at 5 a. m., returning for breakfast at 6, and then returning to the fields to remain until 12 o'clock. The night had been very cold and a heavy dew had fallen, making the air so chilly that it was nearly noon before my overcoat could be discarded. It was an ideal day for bird observation, as there was no wind, and the cold of the morning hours made the birds sluggish, so that it was nearly noon before the Warblers began to seek the higher tree-tops. I noticed, however, that the cold did not prevent the Warblers from singing almost incessantly, even as early as I started out. During the day eggs were found in the nests of Crows, Blue Jays, Mourning Doves and Brown Thrashers, also a Robin's nest with one egg and two newly hatched young.

On account of the extremely backward spring, not only the migrations, but the nest-building, is at least two weeks later than usual with our early arrivals. After losing about forty-five minutes at the dinner table, I again sought the fields, spending the time until 5:15, when we sat down to supper. At 5:45 Mr. Montgomery's son and I started to drive back to town. We reached Milan, on the Rock River, at 7 o'clock, where I entered an electric car and ended my day's observations. Now comes what is to me the interesting part of my story.

During the day I did not see Red-winged Blackbirds, House Wrens, Yellow Warblers, Meadowlarks, Bank Swallows, or Wood Thrushes, until on returning to town we approached within three miles of Rock River, when one by one these birds were seen. This is also true of these birds on my way to the farm on the previous day, and also of several trips made in former seasons over the same ground, the only exception being the House Wren, which usually nests around the farm buildings, but has been seen there only a few times so far this year. Only one Purple Martin was seen during the day, but if we had reached Milan a little earlier more would, without doubt, have been seen, as they nest in the village. But if I had been early enough to see the Martins, I would

have missed the three Great Blue Herons which flew east, high over our heads, when we were still about a mile and a half from the river. Another peculiar fact is that I saw no Grasshopper Sparrows or Vesper Sparrows during the day. However, the latter bird does not seem to be as abundant this year as usual. I noted only one Cliff Swallow, but as they seldom go far from the barn where they nest, and their colonies are not very common, it is not to be wondered at. The most abundant migratory birds seen during the day were the Warblers (mostly Chestnut-sided, Nashville, Tennessee, and Redstarts), and the Gray-cheeked and Olive-backed Thrushes, the Thrushes being very abundant.

THE SONG OF THE DICKCISSELL. *Spiza americana.*

BY P. M. SILLOWAY

Whiling away a recent afternoon over some old piles of ornithological literature, I found an interesting article entitled "Dickcissell in His Illinois Haunts." One paragraph particularly attracted my attention: "The song of this interesting bird invariably begins with three notes, very much resembling the syllables *Dick! Dick! Dick!* These are followed by an indescribable warble with the notes running rapidly together." In his excellent characterization of the song, the writer perhaps unconsciously made use of a term which effectually limits Dickcissell to only one song, the one beginning with the three notes mentioned. Now it happened that during my last summer in Illinois, 1898, I made some special observations regarding the musical ability of Dickcissell, and it may not be amiss to record a part of the notes then entered in my journal. I trust that my friend Mr. Hess will not take this contribution as a criticism of his excellent article in that old Oologist, but merely as a further addition to our fund of recorded observations.

During July, 1898, my home at Virden, Illinois, was near a forty-acre pasture, which was cut into two nearly equal

parts by a railroad. A walk of one hundred yards along the railroad would take me to the pasture, which was tenanted by many families of Dickcissell. The volubility of this songster is well known. Sitting on the porch of my home, I could hear long after sundown the ringing phrases of Dickcissell, wafted to my ears on the clear summer air. It was my custom to walk over to the pasture about four o'clock every afternoon, and follow the tall, tree-like hedge which enclosed the pasture. During these walks I noted down many songs of Dickcissell, with a view to make a complete summary at the proper time. Circumstances, however, called me from Illinois before I accumulated sufficient data, but such as I have to offer may prove of interest. The words suggested by the songs are not vital to the descriptions, but are given as the nearest rendering into English occurring to my mind when the notes were made.

On July 13 I first fixed one particular Dickcissell as a part of a certain detail of the meadow. He soon became a landmark in my walks. His first song was like this: *Dick, dick, ciss ciss sell*, and this rendition proved to be his favorite production. Frequently I could hear in it: *Quick, quick, sell sell sell*, both songs being strongly emphasized at the last syllable Only rarely would this songster offer: *Dick, dick, dick, sell sell sell.*

In another angle of the pasture I was sure to hear a performer who was seemingly addicted to bad habits, for as I drew near his station, he would generally take a stand on a tall weed and shrill out lustily: *Quick, quick, gi'me a chew.* Frequently he would become so importunate as to repeat his plea: *Quick, quick, gi'me a chew, gi'me a chew.* The characteristic production of this songster began with only two notes, but sometimes he would vary the performance by uttering three *dick* or *quick* notes at the beginning. Once I was in another frame of mind, perhaps, though I am free to say that I do not *chew*, for I heard this songster utter, *Dick, dick, miniature*, twelve times in a minute, shortly after he had shouted *Gi'me a chew.*

In the same portion of the pasture where I met the first Dickcissell, another nearby performer would regularly chant: *Dick, dick, quee quee quee, ciss ciss sell.* After a generous

measure of this song, he would change it to: *Dick, dick, dick, ciss ciss sell.* One tenant of the pasture offered a most original production as his characteristic song, which seemed to run like this: *Quick, quick, looky here, skeezics.* After trifling with me in the foregoing manner several times, the last songster would boldly omit the introductory notes, and call out: *Looky here, skeezics, see see see.* It is interesting to note that this songster had four or five songs easily distinguishable. One of his common offerings was like this: *Dick, dick, queerily;* again he would say: *Quick, queerily, quee vee.*

On July 20 I have the following production of another performer: *Dick, dick, dick, queerily, chew chew chew.* This song was frequently varied by changing the number of the *dick* notes, as sometimes only two were given, and occasionally four *dicks* formed the beginning measures.

On July 21 I found a performer who gave the following variation of the characteristic song: *Dick, dick, dick, ciss sell.* Sometimes he would utter only two *dicks*, and frequently, by way of surprise, he would call: *Dick, dick, dick, dick,* omitting the customary phrase in closing.

July 23 brought a new rendition of my July 20th songster's production. This is what I heard: *Dick, dick, dick, what d'ye see see sell.* This peculiar song was varied by changing the number of introductory notes to only two, and sometimes by giving four. Another different song heard on that day went like this; *Quee, quee, quee, quee,—Dick ciss sell,* uttered sometimes with only two *dicks* and three *quees*, like this: *Quee, quee, quee,—Dick, dick, ciss sell.*

July 26 furnished the following records: *Dick, dick,—sell sell sell, see,* which was varied to: *Dick, sell sell see.* The latter, offering only one *dick*, is the most unusual performance of Dickcissell. The same songster furnished this record: *Dick, dick, sell sell sell, quee quee quee;* and also this: *Dick, sell sell sell, quee,* as well as: *Dick ciss sell, quee quee quee.* The following is the most unique record taken, but I think it will stand comparison with the original, *Ciss sell, ciss sell, que, ciss sell.*

SPRING MIGRATION IN LORAIN COUNTY, O., 1904.

BY LYNDS JONES.

There are migrations and migrations, but the migrations of 1904 have proved to be in a class by themselves. For eleven years I have studied both spring and fall migrations at Oberlin with more or less care, and for six years at Grinnell, Iowa. Only once during this period of seventeen years has there been anything like what we have witnessed this year. That was in the spring of 1888, at Grinnell, Iowa, when the weather conditions were somewhat similar during the last days of April and the first two weeks of May.

In 1888 the weather had been about normal up to the middle of April, when a decided warm wave of considerable extent called the early warblers, and other birds which travel with them, north. This wave subsided on the 28th under the pressure of a decided change to cold, cloudy weather, particularly cloudy nights. This unfavorable condition prevailed, with frost on the 13th, 14th and 16th of May, with continued cloudiness and northerly winds, until the 17th. The warm wave which followed brought the birds in swarms, bright males, young, and females, all in the same company, some of them apparently already mated. There were scattering arrivals during the interval, but these were largely confined to hardy individuals and did not represent the forward movement of the host. Some of the migrants tarried until June 10.

The season of 1904, while clearly below the average in temperature for nearly the whole of March, seemed to be opening on the first of April, but speedily made good the traditions of the day, for snow fell on the 3rd. A feeble warm wave on the 8th, 9th and 10th made very little impression on the birds. Another warm wave, covering the 21st, 22nd, 23rd and 24th, caused a sprinkling of arrivals, but the suc-

ceeding eight days were cold and wet, with northerly winds. Then followed a period of warmth and hope for eight days, with maximum temperatures ranging from 62 degrees on the 3rd to 84 degrees on the 7th, culminating on the 9th with a morning storm of rain and cold. More cold on the 10th, 11th and 12th, followed by two warm, and these by six cold days, closed the season so far as arrivals were concerned. At the present writing (May 27) a few of the transient warblers remain.

There had been no distinct movement up to May 4th, but scattering records of hardy individuals proved that the host was not far to the south. Clearly, the weather a hundred miles south was more favorable for the movement of the birds than at Oberlin, during the first five days of May. On the 6th matters began to turn to decidedly favorable, and on the 7th the belated birds rushed in by hundreds. The three succeeding nights were impossible for northern movement. In fact, only the night of the 12th gave much opportunity for migration in the darkness, and evidently not many birds were ready then, and they were obliged to wait until the 21st and 22nd, when the hosts moved north, leaving the 23rd with only the laggards.

As in 1888, brilliant males, females and young, were found together, and apparently in nearly equal numbers. Mating was clearly in progress. Very few birds, except the Raptores and Robins, had succeeded in making much progress in raising broods. Red-winged Blackbirds, which often have young before this time, had scarcely even begun their nests.

One of the most noticeable features of this exceptional migration was the superabundance of Grey-cheeked and Olive-backed Thrushes. For days they were more numerous than Robins, overflowing from the woods into the bordering fields. Wilson and Hermit Thrushes were also unusually numerous, but their numbers were insignificant compared with the others. Many of the warblers were also more than usually numerous, notably the Magnolia, Blackburnian, Chestnut-sided, and Black-throated Blue. On the other hand, the Cuckoos and Wood Pewee were hard to find, and the Chickadee almost refused to be seen or heard. Red-breasted Nuthatches were fairly common for more than a week. One of

the disappointing features was the almost entire absence of any but the regular shore-birds. Lake Erie and the mud flats seemed to have no attraction for them.

The birds of greatest interest, the occurrence of which is worthy of special record are, first, Prothonotary Warbler, twice recorded, being the first county record; Kirtland Warbler, the second county record; Hooded Warbler, the third county record, at least three individuals seen; Yellow-bellied Flycatcher, the first seen in Ohio by the writer; Short-billed Marsh Wren, the third county record; Brewster Warbler, singing almost like the Blue-winged; and a Yellow Warbler with a dusky forehead, suggesting a cross between the Yellow and Northern Yellow-throat, but probably a freak, since the dark forehead, running from eye to eye, was the only Yellow-throat character. It fraternized with the Yellow Warblers, fed like them, acted like them.

Taken all together, this migration season has been a capital illustration of the effect of weather upon the later spring migrations. The weather had continued unfavorable, in the main, for northward movements in the more northern regions, while becoming gradually favorable south. The birds promptly pushed north as far as conditions warranted, then halted to await a change. For the bunching of species which do not regularly migrate at the same time, in a region like Northern Ohio, it has never been surpassed in my experience, and does not seem likely to be repeated in the near future. One who could fail to improve the opportunities for bird study which these conditions presented, must be either luke-warm or else sinfully busy!

A DOOR-YARD LIST FROM MORTON PARK, ILL.

BY ORPHEUS M. SCHANTZ.

The "Door-yard List" from Denmark, Iowa, in the March BULLETIN, was so interesting to me that the thought came that possibly the observations of my wife and I, made in our 50 by 125 foot yard during the last three years, might be of interest to others.

We live about seven miles southwest of Chicago, in a little suburb which, fifteen years ago, contained not a house,

tree nor shrub of any kind. To-day there is a comparatively small area containing trees or shrubs that amount to anything. Four blocks square will cover the bird field, except the prairie birds, for there is open prairie nearly all around us. The nearest timber of any importance lies along the Des Plaines River, about four miles west of Morton Park.

About half a mile north is a section of land, largely vacant, partially surrounded by a belt of trees, some of which are of good size. On the north and east of this tract the trees are planted so as to make a wide belt; on the south and west they are planted in a double row. The trees are principally box elders, soft maples, cottonwood, silver poplar, and balm of gilead.

In our little yard, and in front of it, are, altogether, ten Carolina poplars and cottonwoods, all about ten years old. These trees are very prolific in insect life, and with the profusion of vines—Virginia creeper, bitter sweet, night shade—it may be readily understood why we have been favored with so many "callers" during the spring and fall migration. We have always had the prairie birds, Meadowlarks, Horned Larks, Bobolinks, Dickcissels, quite a variety of Sparrows, and occasionally Bob-white, in the vacant fields around us.

Five or six years ago we first began to notice the migrants, and of course had to have a bird book. Mabel Osgood Wright's Bird Craft proved a great help, and was in constant use for two years, when we discovered Chapman's Hand-Book, since then we have never failed to identify any newcomer. We scarcely ever have to refer to the book now, because we have become quite intimately acquainted with most of the residents and transient visitors. With the exception of the prairie birds, very few nest in Morton Park. Those that do are Robin, Bluebird, Chimney Swift, Chipping Sparrow and Yellow Warbler.

The first pair of Robins nested under a porch roof, within a few feet of the front door of a large residence. When the old gentleman who owned the house discovered the nest, he locked the door and allowed none to use it until the young birds were nearly full-fledged. On the second of March, this year, a flock of fourteen Robins made their appearance, evidently the descendants of the pair mentioned above.

Our first year's record was 25, the second 40, the third 75, and each year since then between 80 and 100. Each year brings either some new record, or brings back some visitors that have missed coming for a year or two.

By sitting on a broad window seat in our bedroom, we could look down into a wide spreading cottonwood, and in that tree we have identified a great many warblers and vireos that ordinarily are so hard to find and see satisfactorily. Last year we caught a fine male Florida Gallinule, which I think was tired and stopped to rest, as there was nothing the matter with it. We took it to Lincoln Park, where it may still be found. The park authorities were glad to get it.

The following birds were found in our door-yard:

Florida Gallinule, Mourning Dove, Screech Owl, Yellow-billed Cuckoo, Black-billed Cuckoo, Hairy Woodpecker, Downy Woodpecker, Red-headed Woodpecker, Northern Flicker, Whip-poor-will, Nighthawk, Chimney Swift, Ruby-throated Hummingbird, Kingbird, Wood Pewee, Least Flycatcher, Yellow-bellied Flycatcher, Blue Jay, Crow, Bobolink, Prairie Horned Lark, Meadowlark, Orchard Oriole, Baltimore Oriole, Bronzed Grackle, Yellow-bellied Sapsucker, Purple Finch, Goldfinch, Redpoll, Snowflake, Vesper Sparrow, White-crowned Sparrow, White-throated Sparrow, Tree Sparrow, Chipping Sparrow, Field Sparrow, Junco, Song Sparrow, Fox Sparrow, English Sparrow, Towhee, Rose-breasted Grosbeak, Indigo Bunting, Dickcissel, Scarlet Tanager, Summer Tanager, Purple Martin, Loggerhead Shrike, Red-eyed Vireo, Philadelphia Vireo, Warbling Vireo, Yellow-throated Vireo, White-eyed Vireo, Bell Vireo, Black and White Warbler, Cape May Warbler, Yellow Warbler, Black-throated Blue Warbler, Myrtle Warbler, Magnolia Warbler, Chestnut-sided Warbler, Bay-breasted Warbler, Black-poll Warbler, Blackburnian Warbler, Black-throated Green Warbler, Palm Warbler, Oven-bird, Water-Thrush, Northern Yellow-throat, Yellow-breasted Chat, Wilson Warbler, Redstart, Mockingbird, Catbird, Brown Thrasher, Bewick Wren, House Wren, Winter Wren, Brown Creeper, White-breasted Nuthatch, Red-breasted Nuthatch, Chickadee, Golden-crowned Kinglet, Ruby-crowned Kinglet, Wood Thrush, Wilson Thrush, Gray-cheeked Thrush, Olive-backed Thrush, Robin, Bluebird.

THE WILSON BULLETIN

A Quarterly Magazine Devoted to the Study of Living Birds
Official Organ of the Wilson Ornithological Club

Edited by LYNDS JONES
PUBLISHED BY THE CLUB, AT OBERLIN, OHIO

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number, postpaid.

Price in all countries in the International Postal Union, 65 cents a year, 20 cents a number.

Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr. Frank L. Burns, Berwyn, Penn., or to Mr. John W. Daniel, jr., Washington, D. C.

EDITORIAL.

The editor will be at Woods Hole, Mass., after June 25, where letters should be addressed, until September 1st. After that date, and until the September BULLETIN is mailed, address Oberlin, Ohio.

Bird-Lore has established the Christmas Census, we have established the New Year Census and the May "All Day" record, and now are ready to add another to this intensive field study of the birds. Shall we not at least try to fittingly celebrate the birth of our nation by making a list of the birds to be found on that day? Would not such a use of the day be more in accord with the real spirit of the anniversary than the usual noisy and dangerous celebration? To those who record 75 or more species on July 4, 1904, the editor will present volumes 9 and 10 of the Wilson Bulletin, or a copy of Bulletin 31, "A Monograph of the Flicker." This summer study will include only the breeding birds and will therefore be of value as showing what birds nest in your vicinity. We trust that this beginning of the study of the breeding birds will prove so interesting that careful studies may follow. It is only by the careful study of the breeding birds of many localities that we shall be able to know for certain the actual distribution of many of our birds. Present day knowledge of summer distribution is largely general; we should make it exact by thoroughly working our own localities.

In asking for New Year records and May records, and now in asking for July 4th records, the editor has in mind a general campaign of field study such as few have ever undertaken systematically. He hopes to prove the intrinsic interest of this sort of work by inducing many to undertake it in these small ways so that they will be glad to broaden out to more and more studies, first looking toward monthly lists of the birds of each locality represented by an observer, and finally making these monthly records into strictly seasonal records for the purpose of determining accurately the bird population during each season. We

should then have Winter Birds, Birds of the Spring Migrations, Breeding Birds, Birds of the Fall Migrations. At first these would run into each other, but as the work grew and experience increased the different groups would separate themselves out, and the records would be of inestimable value in determining distribution. Monthly lists are not only possible but entirely feasible and within the reach of all who work in the field a little. Of course these monthly lists will not be complete, not even for the best of us, and likely not even if we could spend all our time in the field studying, for birds are not stationary structures and man's ability is limited, but they will be not less interesting for that reason. The check-book method of recording briefly field studies ensures annual, seasonal and monthly lists, for the record is a daily record. It is not complete, to be sure, but what is recorded is definite. It is also the easiest sort of record to keep. It can be made to give records of special localities if the observer wishes Thus the writer's check-book records the birds found at the water-works reservoir, on the college campus, in his yard, at Oak Point, while giving the whole general record, all in one column for each species for one day. Simply transcribing gives each list, or only one, at one's pleasure. At best book-keeping is irksome, but here the labor is small for the results obtained. Furthermore, the records, when there are any, are definite even if meagre. They may be made as full as one pleases. Our plea is for more field study looking toward the exact determination of the birds of your region. Work easily, but carefully.

PUBLICATIONS RECEIVED.

Bird Life Stories, Book One, by Clarence Moores Weed.

The book contains twenty-four chapters, each chapter treating of one familiar species. The author has taken the text from the writings of one of four of our most famous writers upon bird life, and therefore, the book is designed as a classical reader wholly devoted to birds. The four writers—Alexander Wilson, John James Audubon, Thomas Nuttall, and Charles E. Bendire -are sufficient guaranty of the matter selected for these twenty-four familiar birds. The book is clearly designed on the lines of The Nature Calendar series which the author of this book has begun for the purpose of combining the usually uninteresting part of learning to read with a training of the senses to recognize the pupil's natural surroundings. It is unfortunate that the excellent quality of the text and general makeup of this admirable book should be marred by the quality of the colored pictures. The three color photograph process is responsible for not a little of the trouble, the mechanical process of printing being evidently carelessly done, but the taxidermy is not always satisfactory. We shall await the appearance of the other two numbers of this series with interest. The series is published by the Rand, McNally Company, Chicago. L. J.

Wild Birds in City Parks. By Herbert Eugene Walter and Alice Hall Walter, Third Edition.

'The new features of this edition are the addition of forty-five bird —the majority of which are shore and water birds—and a simple field

key.' Following 'General Hints' designed to aid in placing the birds under observation in the proper group, a brief description of each of the 145 species treated is given, with the names of other species with which it might be confused. There is a 'Table of Occurrence' giving the number of mornings out of the 454, during the years from 1898 to 1903 inclusive, when each was recorded. A 'Table of Arrival,' arranged systematically by families shows at a glance the year, month, and day of the first appearance of each species. A chapter of 'General Hints' upon the water and shore birds, where the larger groups are given, is followed by 'Particular Hints' upon each of the water birds given. A 'Migration Chart,' 'Key,' 'Glossary' and 'Supplemental List,' with the index, complete this closely packed little book of 66 pages. Inside the back cover there is a large checked blank for recording the migrations of the species given in the book, with additional spaces for other birds. This little book cannot fail to be of great service to those who, living in large cities, would know the birds which may be found in the parks.

L. J.

Gleanings No. III. The Haunts of the Golden-winged Warbler. By J. Warren Jacobs. Published by the author.

This thirty page brochure contains an admirable account of the summer haunts of the Golden-winged Warbler in the region of Waynesburg, Pa., from studies carried on every summer since 1891. The whole subject of nesting is interspersed with interesting descriptions of the region and with unusually good half-tones of both the region and the nesting places and nests. A color chart prepared by hand, and half-tone of a series of eggs close the work, with descriptions of sets of eggs. Mr. Jacobs is to be congratulated upon this close study of a little known warbler, and its clear presentation in interesting form.

L. J.

Birds of the Huachuca Mountains, Arizona. By Harry S. Swarth. Pacific Coast Avifauna No. 4. Cooper Ornithological Club of California. April 15, 1904.

In two pages of introduction the author acquaints us with the mountain range, its position and physical features. Sixty-three pages are devoted to 195 species and subspecies, most of them being copiously annotated. The paper is a model of faunal literature, treating of a very interesting region.

L. J.

The Metallic Colors of Feathers from the Sides of the Neck of the Domestic Pigeon. By R. M. Strong, Reprinted from the Mark Anniversity Volume, Article XIII, pp. 263-277, plate XX, 1903.

This important contribution is the result of long continued careful study of the phenomena of metallic color reflections, chiefly from the neck of the domestic pigeon. Dr. Strong seems to have well nigh exhausted the lines of investigation and concludes that 'The metallic colors of these feathers are probably thin-plate interference colors or

Newton's rings effects which are produced where spherical pigment granules come in contact with the outer transparent layer. The pigment also has the very important function of absorbing light not reflected to the eye as metallic color. The colors seen without a microscope are mixtures of colors from innumerable small points.' L. J.

Birds from Benguet Province, Luzon, and from the Islands of Lubang, Mindoro, Cuyo, and Cagayancillo. By Richard McGregor. Bulletin of the Philippine Museum, No. 3, January 30, 1904.

This number of the Philippine Museum Bulletin is devoted to Zoographical Notes for Lubang, Mindoro, Verde, Cuyo, Agutaya, Cagayancillo, and Benguet Province, Luzon, to Undescribed Plumages and Notes on the Rarer Species, to New Localities for Known Species, and to a list of Species from Irisan, Benguet Province, Luzon. L. J.

Birds of Alleghany and Garrett Counties, Western Maryland. By G. Eifrig. From The Auk, Vol. XXI, No. 2, April, 1904.

The author prefaces the list proper with a brief but lucid discussion of the physical features of the region and the intricate life zones as illustrated by birds, mammals, and plants. The list contains mention of 180 species, with brief annotations. We welcome it as a thoroughly reliable local faunal list. L. J.

Wisconsin Arbor and Bird Day Annual, 1904. Compiled by Maud Barnett. Issued by C. P. Cary, State Superintendent.

This hundred page magazine is prepared with special reference to the school children, with pictures, sketches of birds and trees, selections of poetry, and articles written expressly for this Annual, or selected for their fitness. A number of full page photographs of trees from nature by A. W. Mumford, of Chicago, add to the attractiveness of the pages. The compiler has done a good work which will certainly stimulate nature study among the children for whom it was intended. L. J.

Discovery of the Breeding Area of Kirtland's Warbler. By Norman A. Wood. From Bull. Mich. Ornith. Club, Vol. V, pp. 3-13, March, 1904.

This interesting account of the uncovering of the last warbler mystery is accompanied by four half-tones of nests and typical conditions of breeding, and a half-tone of the only egg found. Mr. Wood has here given us really more information about this rare bird, whose nesting was unknown until he discovered it, than we have of several other warblers. He has also given us careful and numerous illustrations of the song as heard in the breeding grounds. It is interesting to notice that none of the songs which he has transcribed for us resemble that which the writer listened to for a full half hour in Oberlin (see Wilson Bulletin, No. 32, page 2, July, 1900.) We can only hope that collectors will respect the rights of these birds to life, liberty and the pursuit of

happiness instead of rushing in and causing their extermination.

L. J.

The Migration Route of Kirtland's Warbler. By Chas. C. Adams. Bull Mich. Ornith. Club, Vol. V, pp. 14-21, March, 1904.

This second contribution to the life history of *kirtlandi*, an attempt to determine its route of travel from the Bermudas, its winter home, to its known breeding range in Michigan, is well timed. It is accompanied by a like study of the migration route of the Prothonotary Warbler, both being illustrated by maps. This discussion is necessarily limited to the known occurrences of the bird, and from the data at hand its route of migration seems to be from its winter home westward to the Mississippi river, thence north to the mouth of the Ohio, one branch continuing up the Mississippi to St. Paul, on its course giving off branches at the Illinois river, the birds reaching southern Michigan by skirting the southern end of Lake Michigan, and another branch entering northern Illinois. The Ohio offshoot again divides, one course being up the Ohio to Cincinnati, the other up the Wabash to northern Indiana and northern Ohio and eastern Michigan, there following the course of Lake Huron to Mackinac. The course thus pursued almost exactly corresponds to the post-glacial drainage of the regions covered. No doubt the birds pass further up the Ohio and finally cross to Lake Erie by one of the main tributaries, possibly the Muskingum or Scioto. Now that this bird has been brought so prominently before us there should be additional records of migrating birds.

L. J.

Unpublished Letters of John James Audubon and Spencer F. Baird. By Ruthven Deane. From "The Auk," Vol. XXI, No. 2, April, 1904.

This paper consists of a letter from Baird to Audubon, and Audubon's reply, concerning Audubon's proposed western trip, and a recommendation of Baird to a position in the "National Institute," by Audubon. These letters give us a hint of the esteem in which Audubon, then sixty-two years old, held Baird, then but nineteen. L. J.

Warbler Songs and Notes. By G. Eifrig. Reprinted from the Ottawa Naturalist, Vol. XVIII, 1904.

This paper is given to brief descriptions of the songs of 22 of the warblers which visit the region of Ottawa. L. J.

Amateur Sportsman, The Vol. XXX, Nos. 5, 6, Vol. XXXI, Nos. 1, 2
American Ornithology, Vol. IV, Nos. 4, 5.
Atlantic Slope Naturalist, The, Vol. I, No. 6.
Bird-Lore, Vol. VI, No. 3.
Condor, The, Vol. VI, Nos 2, 3.
Journal of the Maine Ornithological Society, The, Vol. VI, No. 2.
Maine Sportsman, The, Vol. 11, Nos. 127, 128, 129.
Monthly Bulletin, The, of the Division of Zoology, Pa. State Dept. Agri., Vol. I, Nos. 4. 11-12, Vol II, No. 1.
Naturaliste Canadien, Le, Vol. XXXI, Nos. 2, 3.
Ohio Naturalist, The, Vol. IV, Nos. 5, 6, 7.

THE BEST LAWN SWING MADE

AGENTS WANTED

Lawn Swings and Settees, Hammock Chairs, Camp Chairs and Stools, Ironing Tables, Wash Benches, Etc.

Agents easily make

$5 TO $10 PER DAY.

Will furnish samples at reduced prices to those desiring agency. Exclusive territory given. Address,

Clearfield Wooden-Ware Co.,
CLEARFIELD, PA.

DAWSON'S BIRDS OF OHIO

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones. 720 quarto pages. 322 birds accurately described, their habits interestingly given. Combines ready identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations to WILSON BULLETIN, Oberlin, Ohio.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds of Ohio are treated. It does not attempt a description of the plumage. Price is 50 cents. Address Lynds Jones, Oberlin, Ohio.

FOR SUMMER READING

The Nature Library

has exceptional charm and value

IS it a pleasant hour with Nature you wish? Or would you confirm a recollection of some point brought up during a recent outing, identify a species, or put your finger on some fugitive fact in Nature? These are but a few of the many uses the books have aside from their principal value as a complete guide to the study of American Natural History, and their attractiveness as entertaining reading.

ARE there little ones growing up about you? Think what it means to a child to have at hand books which attract the interest to subjects sure to result in health and satisfaction all through life.

THE long out-door season will mean much more to you and to the youngsters if these beautiful volumes are at hand for counsel and entertainment. All the year they keep the country in the home, and all that is best and most attractive in Nature is ever before you.

"As necessary as a dictionary, but far more interesting"

TEN SUPERB LARGE VOLUMES

4,000 pages, 10¼ x 7¾ inches; 300 plates in full colors; 450 half tone photographs; 1,500 other illustrations, and *a General Introduction by John Burroughs*

You will incur no obligation and and you will become informed about a most notable work by sending to us the coupon below.

CUT THIS OUT

Doubleday, Page & Co., New York City

Gentlemen:

You may send me at your expense the new booklet containing sample color plates, black and white half-tones, specimen text pages, etc., of the Nature Library. Include also particulars of price and terms.

Name.....

Address.....

City.....

Wil. Bul. 7. '04

We wish to submit our elaborate booklet, which will show better than anything except the books themselves the beauty, authoritativeness, and usefulness of the new edition of

The Nature Library

Doubleday, Page & Co.

34 Union Square

NEW YORK CITY

University
JAN 10 1907
S

Old Series Vol. XVI. New Series Vol. XI, No. 3

The Wilson Bulletin

No 48

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

September, 1904

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.

CONTENTS

	PAGE
Notes on the Holboell Grebe (illustrated). ROBT. J. SIM	67
An Ornithological Reconnoissance of the Grand Reservoir, Ohio, in 1904. W. F. HENNINGER	74
July Fourth Censo-Horizons. LYNDS JONES	76
Spring Migration Along Lake Erie's Shore. ROBT. J. SIM	80
Additions to Winter Birds of Wayne County, Mich. B. H. SWALES, JR.	82
Some Barn Swallow's Nests. C. J. HUNT	83
Lawrence Warbler Breeding in New York City. G. E. HIX.	84
An Addition to the Birds of Ohio. LYNDS JONES.	85
All Day With the Birds	85
Some Needed Work	88
Editorial	89
Notes	92
Publications Received	94

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, Ohio, edited by Lynds Jones.

Subscription. Fifty cents a year including postage, *strictly in advance.* Single numbers 15 cents, unless they are "Special" numbers when a special price is fixed. The BULLETIN, including all "Special" numbers, is free to all paid up members, either Active, Associate or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, 1794 Lanier Ave. N. W. Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for reviews, should be addressed to Lynds Jones, University of Chicago, Chicago, Ill.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 15th of the month preceding publication.

HOLBŒLL GREBE (*Colymbus holbœllii*) IN CHARACTERISTIC ATTITUDES

THE

WILSON BULLETIN

NO. 48

A QUARTERLY JOURNAL OF ORNITHLOGY

VOL. XVI. SEPTEMBER, 1904 NO. 3

NOTES ON THE HOLBOELL GREBE

(*Colymbus holbœllii*)

BY ROBERT J. SIM.

Learning of the capture of a Holboell Grebe near Ashtabula, O., I set out on the first opportunity—Feb. 21, 1904—to see if I could procure the bird. I found it in the possession of Mr. J. J. Topper of Plymouth. This man said that the bird had been seen in the neighborhood for about a week when he took it in thinking to prevent it from starving. It seemed to have come down to a pond near by which was covered with a sheet of "glary" ice, and was unable to rise. Judging from the tracks in the snow as Mr. Topper said, the bird had been trying to reach some wild rose hips that still clung to the bushes. Before the grebe came into my possession strips of fresh fish and raw oysters had been offered it to no purpose, but a little piece of bread had been forced down its throat. This must have been about all the nourishment the bird had had for nearly two weeks. I found it exceedingly thin but able to run about. Bringing it home I kept it in a room where we saw much of each other every day. On the third day of May it was liberated on a small clear pond near Mills Creek, west of Jefferson. Thus it will be seen that this bird was kept in confinement for over two months, during which time opportunity for considerable study was afforded. Below its various occupations and actions are described as closely as my notes allow.

FOOD, FEEDING AND DRINKING.

On the first day strips of raw meat and fresh lettuce were placed in a dish of water before the grebe. These were poked tentatively, then fished out onto the floor, but not eaten. The bird seemed inclined to flop himself into the dish. Towards night two small pellets of raw beef steak were forced down his throat. On the second day I placed a four inch wild fish (shiner?) in a dish filled with water. This was set on the floor in front of the bird. He gave the fish a slight poke whereupon it swam around violently. Making a quick thrust he caught it, grasping it crosswise with the bill—not impaling it. The fish then went through a course of pinching from head to tail, being hitched along from side to side in the bill. It was then turned about and gulped down head first. Later in the day three out of four strips of raw white-fish were eaten, each about the size of a man's finger. These the grebe bruised and shook until small fragments flew several feet around. At this time of the year live food was scarce, but we succeeded in finding a few small aquatic animals. By the twenty-seventh of February the grebe had eaten—all volununtarily—the following:

10 live gold fish—2 to 5 inches long.
2 pieces raw steak (taken from water).
1 four-inch wild fish.
2 large tad-poles.
7 medium sized dragon-fly larvæ.

In swallowing the large gold-fishes the birds jaws seemed to be distended laterally and he gulped so violently that the back of his head struck his back with a hollow "tunking" sound. This operation apparently jarred the fish past the sticking point. When very hungry the grebe swallowed the fishes alive. Of the cray fishes offered him only the small or soft ones were eaten, and no great relish was shown. Earthworms, when their season came, were eaten with avidity, but raw beef-steak (lean) was the principal article of diet with the bird during his stay with us. This came to be taken from the hand, the floor, or water indifferently. In swallowing food the grebe always threw his head outside the normal in violent gulpings, in this respect, as in most, differing from a contemporary coot captive which drew the food into his

throat simply by movements of the tongue and jaws. Unless the diver had already been filled up he invariably ate all the earth-worms and beetles which were placed in the coot's dish.

Only once did the bird really drink, I believe. This was on the first day of our companionship when I put him into a small tank of water. On this occasion he drank eagerly, immersing the bill for an instant, then tipping the head back after the manner of most birds, and repeating this many times.

BATHING AND PREENING

It seems strange that a water-bird should be so indifferent to his natural element. When placed in a tank of clear water, warm or cold, our bird became very wet and soon tried to jump out. After his bath he looked snakey, so closely did the wet feathers adhere to his slender body. I do not think, however, that the under-lying down became much wet; for the bird usually managed to dry himself in an hour or two. A regular bath, such as he took at once after being released, consisted of much water treading, plunging and flapping. In fact the feathers could scarcely be raised one from another they were so completely soaked.

In drying and replacing the feathers the bill was thrust, open into the plumage, then closed and flicked outward, sending drops of water in all directions. The bill was worked over every part of the plumage except, of course, that of the head, and this was rubbed vigorously on the back and wings; or was scratched with the middle-toe nail of the rapidly vibrating foot. When I gently scratched or rubbed the back or sides of the grebe's head he immediately rubbed his head on his back as though the sensation that my scratching caused suggested his own method of producing it. When the bird's back was scratched no such movements were observed. In preening the feathers of the breast and belly he would stand nearly erect on his toes, retaining this position for several seconds, then would step along so as to rest on a dry spot. Often the bird stood up vigorously and frankly, flopping his wings for several seconds at a time. I say frankly because in contrast the coot had a timid, furtive way of doing it.

SLEEPING.

The grebe slept from dark until dawn, and generally for about an hour after a hearty meal. In roosting he ordinarily



HOLBŒLL GREBE (*Colymbus holbœllii*) GOING TO SLEEP

lay on his keel, his feet projecting behind. As the bird became sleepy the feathers gradually fluffed out and the head settled down upon the back. Soon one foot would be seen to twitch slightly. This twitching and curling up of the toes would increase in violence until, with a preliminary rapid waving movement the foot was tucked under the wing which in turn was quickly covered with the flank feathers. Then the other foot was similarly hidden. Thus the feet were entirely concealed beneath the wings and feathers, not even the heel-joints showing.

While in this position the bird allowed me to lay back the feathers and then gently raise the wing. The flat, flipper-like foot was found to lie at full length against the warm, down-covered side of the body, the toes reaching nearly to the "arm-pit." After tucking his feet away one by one, he would draw his head far back and with a peculiar wagging motion, settle it amongst the feathers of the back. With this movement the bill was thrust quite out of sight among the feathers on either side of the neck, more commonly on the right side.

NOTES.

Early in the morning what seemed to be the song was heard. It consisted of a series of cough-like notes "*Cah..... Cah.....Cah*" the bird would say, shaking his head in a most painful manner. This song, if it could be called such, was given four or five times at dawn, and each consisted of four or five of the coughs. When frightened into a desire to inspire fear the grebe opened his mouth very wide (by moving both halves) and gave a rather loud scream which became grating, like the crow's song, towards the end. It somewhat resembled a whinner though, of course, was not so loud. A much more pleasing note than either of the above may be suggested by the syllables, "*Wit-'tah*,"—not loud but rather high, the first part being higher than the second. It was usually given when the bird had just taken wing exercise, or had waddled across the room. It was often given, too, when the bird's back was stroked.

SWIMMING AND DIVING.

In swimming he sat rather deep, the tail and heels usually being submerged. At all times when folded the wings were

quite concealed under the side and flank feathers and humerals. In ordinary swimming the feet struck out alternately. The tarsi extended downward and outward. In diving the bird was not observed to spring forward in the common grebe manner, but rather let himself down very quickly as though drawing his head back through a hole. When it was below the surface I could scarcely realize that the creature before me was a bird, so slender was he and so swiftly did he dart about and shoot through the tangle of aquatic vegetation. It was amazing. The wings were entirely covered by the feathers and the feet struck out simultaneously at the sides, far astern. Their movements could scarcely be followed.

STANDING AND WALKING.

Contrary to the common belief in regard to grebes, this individual was never seen to rest upon his heels, though observed untiringly. The ordinary standing position was with the body somewhat inclined forward, the shoulders humped up, and the tarsi descending forward and outward on a slant—and so entirely free from the ground. The bird could stand for nearly a quarter of a minute or run ten or twelve feet before dropping to his breast. While vigorously preening or jabbing at his feathers he would sway around and sometimes had to take a step or two to regain balance. He seemed to become fatigued after standing for a moment, the strain being felt most, I think, in the toe-joints and in the muscles and tendons which flex the digits; for even with the tarsus slopping back as it did (see sketch) the toes were necessarily pressed upward to an unnatural degree. This conclusion was strengthened by the fact that the toe-joints became much swollen and developed corn-like protuberances. In pattering rapidly along the bird held his body in a semi-erect attitude and it swayed but little, on account of the quickness of the steps; but when walking slowly along he swung noticeably to the side whose leg for the moment supported his weight. He sometimes took food from the floor without first lying down, though he usually assumed the ventropodal position when feeding. When going under chairs and other low articles the bird progressed in short leaps, giving a spring with both legs together and coming down on his breast.

CURIOSITY, FEAR, ETC.

When curiosity and suspicion were aroused the bird's neck was upstretched very slim and the feathers of the body, too, were pressed close. The feet were drawn up under the body at the sides ready for a leap. The suspicious object was regarded intently with one eye at a time. When frightened and disturbed the bird either ran and tumbled to the darkest corner, or defended himself by delivering powerful blows with the nearly closed bill. During the first few days the bird bit me so often and so viciously that my hands were covered with scratches; but thereafter learned that it was useless to try to intimidate me thus. When the coot was let loose in the grebe's room a fight soon took place in which the gray fellow silently bit and scratched with bill and nail, while the grebe sat raining blows with his spear like bill. Their wings were raised threatingly but, I think, not used. Fearing that one might hack the other "into pieces small" I parted them and no more trouble followed during their week of companionship.

COLOR, SENSE, MEMORY, ETC.

During the first few days of the grebe's stay with us we fed him gold-fish which were taken from a dish by the use of a small net made of white mosquito bar. Soon, however, when the bird saw me pick up the net, he hurried over expecting, apparently, to see a gleaming fish drop out. And the moment I even drew a white handkerchief from my pocket he came toddling up with outstreched neck, eagerly watching every movement. Thus I could amuse my friends by making our pet come to calls or signals which, by themselves, would have no effect whatever.

A gray overcoat or a long yellowish gray smock had no terrors for the grebe, but let me enter the room with a broad brimmed gray hat or a scarlet or black skull cap on my head and he would make for the darkest corner. A dark red gown worn by my sister had a similar effect, though most dresses did not frighten him.

He would rest on the hand or lie contentedly on my knee as long as allowed to do so, or would climb onto my foot and let me raise him up on it. But all this was changed the moment that the bird was liberated. He swam at once to the farther side of the pond, and do what I would he would not

return nor could I approach within many feet of him. He was just as shy as any wild grebe and paid no attention to pieces of meat tossed out to him. He saw me then as he has seen men before. I was no longer the great mass standing over him. One is reminded of the little girl at the menagerie who couldn't see the elephant. There he was towering up before her; but he was so big that she could make nothing of him, so she still asked, "Where is the elephant?"

AN ORNITHOLOGICAL RECONNOISSANCE OF THE GRAND RESERVOIR, OHIO, IN 1904.

BY W. F. HENNINGER

The fact that in former years the Grand Reservoir, in Mercer and Auglaize counties, Ohio, was an interesting place for birds, as also that in Dawson's recent investigations it received but scant attention, induced Mr. Karl Heilmann, of Tiffin, and myself to take a summer trip to the Reservoir, to find out what the conditions of bird life would be at the present time. Along the northern side of the Reservoir is the pike from St. Marys to Celina, the tracks of the Lake Erie & Western R. R. and the Western Ohio Traction Co. It is obvious that this part of the Reservoir showed nothing of interest concerning birds. The western part from Celina to the southeast showed us one interesting species, the Yellow-bellied Sapsucker, for the finding of this bird at this time of the year would tend to strengthen Mr. Oberholser's only breeding record for this species in the state. The basis of our work and supplies was Montezuma, on the southwest end of the Reservoir, a quiet little village, the monotony of which is changed only by the advent of a stranger, or an occasional dog fight in which most of the citizens participate with great glee. There we heard that the oil wells in the Reservoir had driven the nesting birds out quicker than anything else.

The Cormorants had ceased to nest since 1886, the Gadwall even before that. One of the natives told me he had not seen a young duck for the last four years, though Mallards and Blue-winged Teal were supposed to nest there still. This did

not seem very encouraging and the first day on the water, which was higher than for the last 13 years, only confirmed this. We saw great numbers of the Great Blue Heron, flocks of 18 and 25, and one Coot, the only one seen on the trip, and one bird which by elimination we thought to be the Sandhill Crane, but identification was not as satisfactory as it should have been. As soon as we reached the oil derricks bird life was extinct.

We rowed back in a fierce gale and concluded to start out bright and early the next morning on the pike to St. Marys. So 6 a. m., July 2, found us out on the road with hip boots, gun and camera. After walking nine miles to the east we turned off to the Reservoir, and three miles east of the oil wells struck a place which was promising. The trees fringing the Reservoir were standing in about three feet of water, mostly willows, water ash and a few oaks. With their green arches they were keeping almost every ray of sunlight away from the oozy recesses beneath. Spotted Sandpipers and Redstarts were common and in a few moments I saw a female of the Prothonotary Warbler, which we observed for about five minutes. Later on in a similar place we saw another female of the same species. This would indicate that it is still a very rare summer resident at the Reservoir. The call note, to my ear, faintly resembled the chirp of the Yellow Warbler, while the colors in general, setting aside the size, would remind one of the Blue-winged Warbler minus the wing bars.

In this shady retreat we also found the Chestnut-sided Warbler in several specimens. Most likely they had bred in the neighborhood. Rowing across a space of open water we soon entered a dense swampy margin and here was the only place where the Long-billed Marsh Wren could be found, and then sparingly only, not to be compared with the vast numbers at the Sandusky Bay marshes. Bitterns were booming plentifully, but the Least Bittern was not there. Gallinules and Rails were also absent, while a Wapakoneta sportsman, who is also a practical taxidermist, had found a nest of the King Rail on the previous week off Russell's Point at the Lewiston Reservoir. After poking around in the cat-tails for a while, we suddenly started a female Blue-winged Teal, but in spite of a very diligent search, we did not find the nest. The

bird was in an excellent condition and flew rapidly away, proving that she was in no way a crippled bird. Evidently the Blue-winged Teal is a rare summer resident at the Grand Reservoir. No other water birds were seen. Tree Swallows and Martins were very common and I am surprised to see how commonly, in northern Ohio, the Purple Martin takes to the woods and swamps to nest, and how rare comparatively the species is, while in southern Ohio, it is hardly ever seen away from the bird houses and is a common bird. The ordinary Ohio land birds were all common at the Reservoir with the exception of the House Wren, which was rather rare. All told 67 species of birds were heard or seen.

It certainly does not pay an ornithologist to make a visit to the Grand Reservoir, except in the migration seasons, the Licking Reservoir no doubt being the most profitable inland body of water in Ohio. The interesting things which Mr. Dury found at the Grand Reservoir in former years are gone for good, and after coming to this conclusion we wearily tramped the nine miles back to Montezuma, packed our grips and went home.

JULY FOURTH CENSO-HORIZONS, 1904.

Judging from the writer's experience, the Independence Day bird work was far less interesting and inspiring than the May migration work, and it was entered into with that lack of enthusiasm which summer heat is pretty certain to bring about. However, the work actually done proves that there is, after all work to be done in summer which will count quite as much as the May work in the final result. Heat, foliage, and last but not least, insects, make bird study in summer irksome. Take away the annoyance of the insects and I venture to assert that the present hesitancy about going into the woods in July would give place to commendable enthusiasm. Let us hope that so much of the millennial time will soon come.

The reports which have been received cover a pretty wide range of country, but are not as well scattered as the May reports. They are interesting in disclosing what are in the breeding birds of the several regions. One would suppose that a

larger number of breeding birds might easily be found in each of the regions represented.

The regions from which reports have been received, begining in the extreme east, follow:

WOODS HOLE, MASS. R. L. Baird, I. A. Field, and Lynds Jones. 5 to 7 and 8 to 12 a. m., 1 to 5 p. m. Mainland in the morning, Nonamesset Island and the surrounding water the latter part of the morning and afternoon. Little work done after dinner hour. Weather fair.

CROSSWICK, N. J. Charles H. Rogers. First hunt 5:50 to 8 then 9:25 to 5:05. Weather fine. From Crosswick through Yardville to and around Laurie's Pond and return.

ENGLEWOOD, N. J. George E. Hix. Mostly damp woods and extensive marsh; no fields or orchards and only about a mile of road. 7:15 a. m. to 6:55 p. m. Weather fine.

CANANDAIGUA, N. Y., AND VICINITY. Frank T. Antes. 4:30 a. m to 5 p. m. on July 4th; 8 to 10 a. m. on July 5th. Village of Canandaigua, five miles of lake shore, a marsh at the foot of the lake, and a small tract of woodland about seven miles up the lake. Weather fine.

GRASMERE RHINEBECK, N. Y. Clinton G. Abbott and Maunsell S. Crosby. 6 to 8 and 9 to 10 a. m. Area covered about 150 acres. Fine day.

CUPOLA, WELSH MOUNTAIN, CHESTER COUNTY, PA. Thos. D. Keim and Chreswell J. Hunt. 5 a. m. to 7 p. m. Distance covered over 15 miles. Weather Fine.

GENEVA, O. A. W. Galpin and Robert J. Sim. Cowles Creek marsh and lake shore. 4:30 to 12 a. m. Hot day, partly clear, becoming rainy p. m. 20 acres in the marsh, and woods and thickets near.

WATERFORD, OAKLAND COUNTY, MICH. Alexander W. Blain, Jr. 9 a. m. to 7:30 p. m. Warm day, showers in p. m. In Woods, fields, meadows, marshes, streets of town, along railroad tract, and along the lakes.

HOPE, KANS. O. H. Pease. 1 to 2:40 p. m. Heavy rain the night before, day warm, partly cloudy and sultry. A small grove and hedge fence. Distance about 1½ miles.

	Woods Hole, Mass. Baird, Field, Jones	Crosswick N J C. H Rogers	Englewood, N. J. Geo. E. Hix	Canandaigua, N. Y. F. T. Antes	Grassmere, Rhinebeck, N. Y. Abbot, Crosby	Cupola, Pa. Keim, Hunt	Geneva, O., Galpin, Sim	Waterford, Mich. A. W. Blain, Jr.	Hope, Kans. O. H. Pease
Loon	8							2	
Herring Gull	2								
Laughing Gull	2								
Common Tern	c								
Roseate Tern	c								
Mallard								5	
Wood Duck							1		
American Bittern				x					
Broad-winged Hawk	1								
Least Bittern							2	2	
Great Blue Heron						2		1	
Green Heron	2	3			x	3		1	
Blk-crowned Night Heron					x	1			
American Woodcock								1	
Spotted Sandpiper	4			x	x		4	6	
Bartramian Sandpiper						2			
Killdeer		1		x		8	3	4	
Bob-white		1			x			1	1
Ruffed Grouse								1	
Mourning Dove		2		x	x	1		2	15
Turkey Vulture		5				4			
Marsh Hawk								1	
Sharp-shinned Hawk				x					
Red-tailed Hawk								1	
Red shouldered Hawk					x				
White-winged Scoter	1								
Am. Sparrow Hawk		6					2		
Screech Owl									2
Yellow-billed Cuckoo		1	1			3		1	
Belted Kingfisher	3	4		x	x	1	2	1	
Hairy Woodpecker		2			x	1		1	
Downy Woodpecker		6	1		x	3		3	
Red-headed Woodpecker						4	3	2	
Northern Flicker	4	17	1	x	x	8	5	6	8
Whip-poor-will								1	
Nighthawk									
Chimney Swift	10	40	4	x	x	c	10	15	
Ruby-throat Hum'gbird	3				x	1		1	

Kingbird	5	7	1	x	x	5	6	10	19
Crested Flycatcher		4	2	x	x		1	3	
Phoebe		1	1	x	x	5	2	6	
Wood Pewee	10	5		x	x	3	3	5	
Green-crested Flycatcher		2	1			1			
Least Flycatcher				x					
Prairie Horned Lark							2	10	
Blue Jay		2	1		x	7		1	2
American Crow	c	8	6	x	x	c	9	20	
Starling			1						
Bobolink			1	x	x			30	
Cowbird	4	4	4	x	x			8	2
Red-winged Blackbird	c	1	11	x	x	c	20	c	
Meadowlark	2	19		x	x	c		a	
Western Meadowlark									8
Orchard Oriole		4			x	4		1	7
Baltimore Oriole	3		1	x	x	1		2	11
Purple Grackle		x	1		x	c			
Bronzed Grackle	c			x				10	c
Purple Finch				x					
American Goldfinch	5	2		x	x	8	3	c	1
Vesper Sparrow	10	1	1	x	x	a	3	4	
Savanna Sparrow				x	x				
Grasshopper Sparrow					x	9		1	
Lark Sparrow									
Chipping Sparrow	10	22	4	x	x	3	1	c	
Field Sparrow	4	5	3	x	x	c	4		
Song Sparrow	c	25	35	x	x	c	11	a	
Swamp Sparrow		2	9						
Towhee	c	3	1		x	13		2	
Cardinal		1				1			
Rose-breasted Grosbeek		2	2		x			1	
Indigo Bunting	2	2	4	x	x	5	3	5	
Dickcissel									15
Scarlet Tanager		2	1	x	x	4	1	1	
Purple Martin		1		x		4		c	2
Cliff Swallow	1							10	
Barn Swallow	c	12	2	x	x	11	5	a	5
Tree Swallow	2		3					4	
Rough-winged Swallow		4		x		6		2	
Cedar Waxwing	c		6	x	x		1	8	
Migrant Shrike								1	
Red-eyed Vireo	c	17	14	x	x	c	5	2	
Warbling Vireo		1		x	x			5	
Yellow-throated Viero		4		x	x				
White-eyed Vireo		3				3			
Bell's Vireo									2
Black and White Warbler	2	2		x	x				

Worm-eating Warbler			1		x	2				
Golden-winged Warbler					x					
Yellow Warbler	c	5	3	x	x	1	1			5
Chestnut-sided Warbler	c		1		x	2				
Prairie Warbler	c									
Blk-throat green warbler	c									
Northern Parula Warbler	1									
Kentucky Warbler			2							
Oven-bird	c	3	9	x	x	6			1	
Louisiana Water-Thrush			2	x	x					
Northern Yellow-throat	c	5	9		x	c	3			
Yellow-breasted Chat	1		1			6	1			
American Redstart	2		4	x	x				4	
Mockingbird										1
Catbird	c	12	11	x	x	c	3		2	12
Brown Thrasher	c	4			x	4	1		1	15
House Wren		4			x	7			1	3
Long-billed Marsh Wren			8	x			2		1	
White-breasted Nuthatch		1	8	x	x		1			
Tufted Titmouse		5				1				
Chickadee	6			x	x				3	
Carolina Chickadee		2								
Blue-gray Gnatcatcher									c	
Wood Thrush		3	2		x	4			5	
Wilson Thrush	3		1	x	x					
American Robin	c	35	20	x	x	c	10		5	3
Bluebird	4	3	3	x	x	2	4		5	1
Hooded Warbler			9			3				

SPRING MIGRATION ALONG LAKE ERIE'S SHORE.

BY ROBERT J. SIM.

While I was staying along the shore of Lake Erie near Geneva, Ohio, last May (1904) I was much interested in noting the diurnal flights of birds, which, the longshoremen assured me, occur every year. These flights took place on several fine days before the middle of the month. I made observation only on days when the movement was not conspicuous, that is, on May 7th, 12th and 13th. The weather was warm, the sky cloudless but hazy, and a rather strong south breeze was felt.

The birds were all passing from west to east in a belt about one-fourth of a mile in width, the main body passing along over the beach. For the most part these flights were

over before noon, and the earlier part of the morning seemed to be the favorite time. My notes are not complete because the flights were noticed a day or two before any records were kept, and because I had other bird work in hand and could not attend strictly to the migrators.

The *Raptores* were well represented. Those which I shall call *Buteo* were not identified with absolute certainty though I should feel little hesitancy in calling them American Rough-legs. They were all of the same species. The size was that of the Red-shoulder or a trifle greater. The plumage above seemed to be rather dark, and below the birds were whitish with a dusky band varying in width and distinctness across the belly. The tails all appeared like that of a young Red-tail—that is, rather finely barred with dark and light. None of these birds stopped, but the Sharp-shins often did so, alighting among the trees and causing great disturbance among the small birds. Some of these small hawks had their crops vastly distended. Most of them were carefully looked at through an 8-power prism glass and were found with one exception to be in the brown, immature plumage. The Ospreys flew low, sometimes alighting on one of the trees along the bank. They varied much in the completeness of the necklace.

The Blue Jays flew just above the top of the trees, sometimes stopping for a moment in them. One flock stopped in an oak tree to feed. Nineteen seemed to be a favorite number for a flock, but they ranged from single birds to a flock of thirty. The flocks trailed out sometimes being in all a hundred yards long. These jays were absolutely silent except in the cases of stray ones.

Below I give the species of birds, with the number seen, and the dates.

	MAY 7	MAY 12	MAY 13
Marsh Hawk	3		
Sharp-shinned Hawk	72	24	5
Cooper Hawk	2		
Buteo	18		
Am. Sparrow Hawk	2	1	1
Am. Osprey	5	1	
Kingbird	13		
Bluejay	120	few	148

Am. Goldfinch..... 10.....3 small flocks.....
Am. Pipit.....200.....3 flocks..... .....3 flocks

On the 12th there was a conspicuous movement of swifts and swallows. They flew mostly in the fore part of the morning and towards night, going east. In the morning every one or two minutes a small company went by. In proportion they were as below:—

Bank Swallow by far the most; Barn Swallow 2nd; Chimney Swift 3rd; Cliff Swallow 4th; Tree Swallow 5th; Rough-winged seldom.

In the afternoon the flight continued from 3 to 6:30 p. m. Every two or three minutes a scattered, feeding flock of 20 to 50 passed along. Species as follows:

Cliff Swallow predominating; Barn Swallow 2nd; Swift 3rd; Bank Swallow 4th and an occasional Tree Swallow.

ADDITIONS TO LIST OF THE WINTER BIRDS OF WAYNE COUNTY, MICH.

BY BRADSHAW H. SWALES.

Since the publication of my list of the Winter Birds of Wayne county, Michigan, in THE WILSON BULLETIN, March, 1903, I have been able to add the following to the list of 61 species above recorded.

62. LARUS ARGENTATUS. Herring Gull. Accidentally omitted. A common winter resident going north in late March.

63. LARUS MARINUS. Black-backed Gull. Reported to be occasionally taken at the Flats in winter. One shot on the Detroit river during March, 1904, and sent in to a local taxidermist.

64. LOPHODYTES CUCULLATUS. Hooded Merganser. I observed a male December 3, 1903, at Belle Isle. Another was sent in later in the month to L. J. Eppinger.

65. CHARITONETTA ALBEOLA. Bufflehead. I noticed one bird at L. J. Eppinger's which was shot on the Detroit river in December, 1903.

66. ANAS BOSCHAS. Mallard. A number were taken during December, 1903, on the Detroit river.

67. ANAS OBSCURA. Black Duck. Several observed February 3, 1903, on the lower part of the Detroit river.

68. NETTION CAROLINENSIS. Green-winged Teal. A late bird was shot in early December, 1903, off Fighting Island, by F. Bryant.

69. ACCIPITER ATRICAPILLUS. American Goshawk. J. B. Purdy records a bird shot December 24, near Plymouth. (Bull. Mich. Ornith. Club, II, 38.)

70. CERLYLE ALYCON. Belted Kingfisher. Chas. Freiburger records one January 14, 1903, at the Detroit Water Works. (Bull. Mich. Ornith. Club, IV, 28.)

71. PINICOLA ENUCLEATOR LEUCURA. Canadian Pine Grosbeak. The first birds of this species were recorded in Wayne county, on November 9, 1903, when two birds were shot near Detroit and sent in to L. J. Eppinger. Jas. B. Purdy writes me that he shot one December 9th near Plymouth. On March 6, 1904, I met with two near Palmer Park, north of Detroit.

SOME BARN SWALLOW NESTS.

BY CHRESWELL J. HUNT.

The Barn Swallows shape and vary their clay nests according to the site selected for them. The most common form seems to be that placed on the side of a beam, in which case the nest is usually in the form of a reversed half cone—the top being a half circle while the base is pointed. I also have a nest which was built where a lath projected two inches from the base of the beam. In this case the birds used this lath end for a foundation and built the nest above it. This nest is in the usual shape, the top being a half circle, but the bottom, where it rested upon the lath, is flat instead of pointed. Another nest I have seen was placed in a corner where two boards came together at right angles. It had the usual pointed base but was exceptionally long.

I have had a pair of Barn Swallows under observation for the past four summers. Their nests were built under the barn-yard roof. They first built the nest, above referred to, placed upon the lath. They used this nest for two seasons when I removed it and sawed off the protruding lath. The next year they built a nest upon the beam right beside the

spot where the old one had been. This nest was of the usual pointed shape. Both of these nests were lined with straw—no feathers being used.

This summer (1904) what I believe to be the same pair of birds built a nest on top of a strip of wood about five inches wide. This nest is shaped almost like a Phœbe's, being a high circular nest. The centre of this nest was made almost entirely of feathers into which the eggs sank and were almost hidden from view. While the female sat upon the nest laying an egg the male would always be perched upon a near-by stick, apparently on guard.

It is a wonder these birds selected this place for a nesting site as it is just above a door where people are passing every little while all day long. In constructing this nest the birds made themselves just about twice as much work as they would have had building one of the half cone style. As I believe they were successful in rearing all their former broods this new style of nest was hardly built for more protection. Whether it is less arduous for them to build a nest on a solid foundation than to fasten it to the side of a beam I am unable to say. At any rate this circular nest took just about double the material for its construction.

LAWRENCE WARBLER (*Helminthophila lawrencei*) BREEDING IN BRONX PARK, NEW YORK CITY.

BY GEORGE E. HIX.

On May 18, Dr. Wm. Wiegman found a typical male Lawrence Warbler (*Helminthophila lawrencei*) mating with a female Blue-winged Warbler (*H. pinus*). At that time they were collecting material but the nest was not found until after the young, six in number, were hatched. I first saw Lawrence on the 11th of June. At that time the young had apparently been hatched a couple of days. On the 17th the nest was empty, but contained fresh excrement. When last seen there was nothing to show whether the young birds would be typi-

cal *pinus* or tend toward *lawrencei*. This of course could not be determined until the juvenile plumage had been moulted and the first winter plumage assumed. The two songs heard were both different from that of the Blue-winged Warbler, being somewhat between that and the Golden-winged.

AN ADDITION TO THE BIRDS OF OHIO

BY LYNDS JONES.

Mr. W. E. Clyde Todd calls my attention to a record which entirely escaped my notice when compiling my 'Revised Catalogue of the Birds of Ohio'. The published note follows.

'Smith's Longspur in Ohio.

By Clark P. Streator.

Smith's Longspur (Calcarious pictus) Collected at Garretsville, Ohio, on Jan. 29,'88. I observed a large flock of strange birds busily engaged in feeding upon the seeds of ragweed. They would only stay a moment in a place and were very shy, but I was lucky enough to secure two very fine specimens. I belive this to be the first time this species has been taken in Ohio'. Ornithologist and Oologist, 13, page 95.''

This species should be added to the list of Accidendal species in the Revised Catalogue.

ALL DAY WITH THE BIRDS.

This may seem an inopportune time for discussing a subject which has special reference to an all day study of the birds during the height of the spring migrations. It is certainly true that this all day habit grew out of a special effort put forth to determine as nearly as possible what and how many species of birds might be found in a limited region in one day, but it has long outgrown merely that. These spring migration all day studies proved so valuable in many ways that entire days were given in other seasons to the

same sort of study, with no less interesting and valuable results. Some of my friends are making monthly records of the birds, planning at least four all day studies during each month. Some, with less opportunity for study, are making seasonal lists, based primarily upon three all day studies, one at the begining, one at the middle and one at the close of the season, supplemented by such records as may be made during the intervals. It is clear that the person who can spend a considerable time with the birds during each month will have at command monthly, seasonal and annual lists. In making any seasonal divisions one must largely disregard the calendar seasons and group the birds according to whether they are resident during the year, whether they are seen only during the winter months, or winter weather, or more exactly, whether they come from the north and spend the winter, whether they are transient visitors, passing the region twice each year, or whether they come from the south, breed and pass south again. Careful study will always disclose the group to which each species belongs during a series of years.

If we would understand geographical distribution, local and general, fully, a considerable time must be spent with the birds at all seasons. While the mapping of areas of distribution of birds must be based upon breeding birds, because it is during the breeding season that they are really settled for a time, while at other times they are wandering about to a greater or lesser extent, their distribution during the winter season is no less interesting, and capable of throwing light upon some of the problems of summer distribution. The routes of migration must also be regarded in the light of geographical distribution, and the mapping of these routes is of more importance in the discussion of the distribution and origin of present day species than many persons think. These remarks apply principally to the question of continent distribution, it is true, but how are we to exactly know about such general distribution unless we begin with local distribution? Hence, the local list becomes of great importance as furnishing the material for such exact knowledge of general distribution. It may not be clear how the all day studies at any particular time bear upon this question of geographical distribution. My answer is that they furnish the means of knowing what

birds are in the region at the time of the study. Ten consecutive hours spent in a study of the birds only once a week will prove of more value in determining what birds inhabit the region than triple that number if scattered over the week in two hour lots, for the simple reason that the ground can be so much more thoroughly covered. The greatest value of the short studies lies in coming in touch with the birds often so they can be learned more readily. But it is one thing to learn the birds and quite another thing to learn what ones inhabit a given region. To you who must spend a considerable time learning the birds the shorter studies will be the more attractive, but once the task of learning is completeed, let the longer studies have a place, for the sake of the local list.

The reader will at once infer that 'All Day' bird studies for times not hitherto indicated will be suggested. At the risk of being considered an 'All Day' crank (might as well be killed for an old sheep as a lamb!) I want to propose to each reader who has become inoculated with this 'All Day' germ some one full day's study in the succeeding months. The earlier in October such a study can be arranged the more birds will be recorded, but any day will be better than none. I would certainly be considered demented if I suggested Thanksgiving day for such activity, but for students and educators, at least, the Saturday following the feast might prove both interesting and healthful. For December your services are already solicited. I would urge that Mr. Chapman be given hearty support in extending his Christmas Bird Census over the whole country. You cannot properly enjoy the day without getting near the heart of Nature.

Another line of study is suggested elsewhere—that of laying more stress upon the fall migrations. It is now too late to note the beginnings of the southward movement by many of the breeding birds, but it is not too late to record the final departure of many species. Plan now to give next year's southward movement proper attention.

SOME NEEDED WORK.

Those who imagine that we are nearing the end of bird study—that there is not much left worth spending time upon —should search through all available bird books for such common things as most phases of breeding habits, the problems of song, most questions relating to food and the manner and times of feeding, the effect of light and wear and age upon color of the feathers. In short, there is nothing yet fully known even in the field of the more evident matters relating to the birds. In the less evident, having to do with mental processes, physiological processes and the like almost nothing has been done. Anyone should be able to throw light upon the questions of nest building, deposition of the eggs, period of incubation, young in the nest and out of it, and anything else relating to the life as revealed in the nesting season. There are doubtless more than a dozen species of birds nesting within your reach every year, and yet it is not a hazardous venture that you are ignorant about all of these suggested questions. Ought you to be? Such work will prove intensely interesting and of great value. It must certainly be within the reach of all who live where trees grow. For such work the birds need not be disturbed to the point of leaving the nest in order to carry on the necessary observations. If the nest is too high to be looked into without climbing too near it and where a step-ladder cannot be used, a small mirror on the end of a stick will often suffice. But if the study taxes your ingenuity, so much the better!

THE WILSON BULLETIN

A Quarterly Magazine Devoted to the Study of Living Birds
Official Organ of the Wilson Ornithological Club

Edited by LYNDS JONES

PUBLISHED BY THE CLUB AT OBERLIN, OHIO

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number postpaid.

Price in all countries in the International Postal Union 65 cents a year, 20 cents a number.

Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr. Frank L. Burns Berwyn, Penn., or to Mr. John W. Daniel, Jr., Washington, D. C.

EDITORIAL.

The editor will be in Chicago, Ill., from October 1st, until June 25. All communications and publications should be addressed to him at the University of Chicago, Chicago, Ill., between those dutes, or until further notice.

The editor spent another delightful summer at Woods Hole, Mass., and the islands in the vicinity, in company with Mr. R. L. Baird and Mr. I. A. Field. Nine days were spent on Muskeget island with Mr. Baird, and the other places of interest to an ornithologist were visited. Many interesting things were learned about the gulls and terns which will be shared with our readers later. A short trip among the lake Erie islands furnished an interesting comparison between the salt water and fresh water Common Terns. Much more consecutive time is needed before the whole history of the terns nesting in either region can be learned.

We hope that many bird students have undertaken a careful study of the autumn migrations, which began more than a month ago. We are now just in the midst of the southward movement, when fall work becomes the most interesting. The most important part of this fall work are the exact records of southward departure, or at least of that part of the work which remains of this season. Record should be kept of both last seen and the departure of the bulk of the species. Very few migration note books contain much information about any phase of the southward movement. Some of us have to be stay-at-homes all summer, and to such must we look for complete records of the southward movement.

We are pleased to note that Mr. Frank M. Chapman, the best known writer on birds at the present time, has undertaken the task of collecting a complete life history of our warblers by the co-operative methods upon which our Club was founded. Let every member of this Club and every reader of this BULLETIN aid him in his good work to the full extent of your ability. There are possibilities in this field almost beyond conception if all who can will lend their aid. One great part of our work as a Club has been in raising up a generation of bird lovers who would know how to study the birds for furthering our knowledge of them. Now that an opportunity offers let us show Mr. Chapman that our efforts have been productive.

We are also pleased to note that SCIENCE, for August 26, announces the establishment of 'A Station for the Study of Bird Life' by Mr. Charles C. Worthington, at Shawnee, Monroe county, Pennsylvania, with Mr. William E. D. Scott in charge. Mr. Scott is well known as the Curator of the Department of Ornithology at Princeton University. He has written of his experiment with birds there, particularly in regard to inheritance of song. This station is under the charge of the Worthington Society. All phases of bird life will be studied, under normal as well as artificial conditions. We may expect some valuable results from this establishment.

Members and readers will be interested to know that Mr. J. Warren Jacobs' collection of the 'Eggs of Native Pennsylvania Birds' is now on exhibition at the St. Louis Worlds Fair. It is installed with the Pennsylvania economic zoological exhibit in the Palace of Agriculture. It contains the eggs of 159 species which have been known to nest within the state, several having been added to the collection since it was exhibited at the Chicago Worlds Fair, eleven years ago. All will be interested to see the collection about which Mr. Jacobs has written so entertainingly.

Ill health has hastened the removal of our Vice-president, Rev. W. L. Dawson, to the Pacific coast. For the present his address will be Blaine, Washington. We trust that the bracing air and salubrious climate of that region may speedily restore him to complete health and vigor.

N. B.—The editor has concluded to issue a sheet of errata with the index, both to appear with the December BULLETIN. He will be grateful for notice of any errors appearing in contributed articles. He will also be grateful for articles and notes intended for the December BULLETIN. Everything intended for the December BULLETIN should reach the editor at his Chicago address not later than November 15.

NOMINATIONS FOR 1905.

Nominations for officers for 1905 are now in order. The list of present officers and members may be found in the March, 1904, BULLETIN. Every member should manifest interest enough in the Club to make full nominations. A postal card is sufficient to contain the required number of names. Address it to Lynds Jones, University of Chicago, Chicago, Ill. If you will give this matter your immediate attention there will be no dearth of nominations.

IMPORTANT NOTICE: There are ready for distribution title pages for the following volumes of THE WILSON BULLETIN: New Series Vols. 3, 4, 5, 7, 8, 10, 11. All who wish these title pages may secure them by giving notice to the editor before the December number is mailed. Unless otherwise directed, all such requests will be met by enclosing the title pages wanted with the December BULLETIN. They will be mailed earlier if a one cent stamp accompanies the request.

NOTES.

BOBOLINKS INCREASING AT BOULDER, COLORADO.

Bobolinks, once supposed to be rare in Colorado, are common in at least three localities in Boulder county this year, and their songs have been much enjoyed by the writer and others. Last year I saw one east of Boulder and Dr. J. R. Brockett reported one on the University campus at Boulder, and the year before Mr.L.C. Bragg took one near town, but this year I have seen about a dozen every day that I have visited one locality. Popularly our Lark Bunting has been oftencalled "Bobolink," though there is no good reason for the confusion, as they are readily distinguished. JUNIUS HENDERSON.

A JANUARY KINGFISHER.

On January 19th of this year I observed a Belted Kingfisher about two miles below this place on the bank of the Hudson river. The mercury stood at 2 degrees below zero and the river was frozen solid from shore to shore.

I also recorded a Sparrow Hawk the same day, but this is not unusual as I have seen several during the past winter.

Ossining, N. Y., May 6, '04. FREDERICK C. HUBEL.

NOTES ON THE NESTING OF THE BLACK TERN.

On June 27, of this year, Mr. Karl Heilmann and the writer found a nest of the Black Tern with three fresh eggs in the Sandusky Bay marshes. The eggs were placed on a few broken fragments of old flags on what seemed to be a pancake of muck. After I had photographed the nest and eggs and put the dripping camera into the boat, I examined the foundation. The pancake of muck rested on some floating vegetation. This in turn was resting on an old sunken muskrat house, probably three or four years old, which had formed a kind of a submarine decayed island. This would indicate that both Langdon as well as Jones and Dawson were right in their respective statements which seem to be at variance; it is the combination of both which gives us the correct fact at least in this one particular instance. New muskrat houses or those a year or so old were not used, though in abundance.

We also saw a pair of Belted Piping Plovers on this date, the nest had probably been destroyed by a recent storm, which had driven the waves of Lake Erie into the bay clear over the sand-dunes of the Cedar Point peninsula. W. F. HENNINGER.

NORTHERN OHIO NOTES.

Roseate Tern, (*Sterna dougalli*,) was found in a company of Common Terns which were flying above Big Chicken island, lake Erie, August 26, 1904. While this island lies in Canada it is so near the border of Ohio that the occurrence of this rare Ohio tern here should not

be allowed to pass unnoticed. The bird was in full breeding dress, and gave evidence by its actions, of belonging to the island. Many of the other birds were in molt, but this specimen gave no evidence of molt.

Black Tern.(*Hydrochelidon nigra surinemansis.*) No birds were to be seen at their regular breeding places in the Sandusky marshes, August 24 to 27, but they were found in numbers at the Chicken islands with the Common Terns, most of them in transition molting plumage. Some had almost completed the molt, while others had scarcely more than begun. A few Black Terns were also found with small companies of Common Terns roosting on the sand spits among the Bass islands.

It may be interesting to note that North Harbor island, which was a favorite breeding place for the Common Terns in 1901, gave little evidence of having been inhabited by more than a handfull of birds the past season. Likewise, Chick island, which was a low reef supporting a considerable growth of Smart-weed (*Polyganum*) and numerous nesting terns in 1901, was reduced to a wave washed reef entirely devoid of either vegetation or tern nests on the day of our visit, August 26. However, roosting terns of both species, Herring and Bonaparte Gulls, were numerous enough to almost completely cover the rocks. The gulls were also in molting plumage.

The lagoon of Middle Bass island, which yielded such rich bird life in 1901, was full to the brim, and surrounded by three feet of water on all sides. Consequently there were no mud flats, and no shore birds. A flock of Shoveller—decoys gave me palpitation of the heart until their true structure was revealed, when three small boys in a boat rowed through their midst. A single Least Bittern made an additional record for the lagoon. Black Terns were also in evidence here. Numerous Pied billed Grebes seemed entirely at home, and probably bred here during the summer, with the King Rails and Florida Gallinules. both of which were feeding everywhere.

The only shore birds recorded among these islands were Sanderlings, Turnstones, Spotted Sandpipers and Killdeers. This was disappointing, in view of the records of others at Oberlin earlier in the month. I have noticed, however, that many of the shore birds seem to prefer inland ponds and mud flats during their southward journey.

LYNDS JONES.

THE BEST LAWN SWING MADE

AGENTS WANTED

Lawn Swings and Settees, Hammock Chairs, Camp Chairs and Stools, Ironing Tables, Wash Benches, Etc.

Agents easily make

$5 TO $10 PER DAY.

Will furnish samples at reduced prices to those desiring agency. Exclusive territory given. Address,

Clearfield Wooden-Ware Co.,
CLEARFIELD, PA.

DAWSON'S BIRDS OF OHIO

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones. 720 quarto pages. 322 birds accurately described, their habits interestingly given. Combines ready identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations to WILSON BULLETIN. Oberlin, Ohio.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES.

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds of Ohio are treated. It does not attempt a description of the plumage. Price is 50 cents. Address, LYNDS JONES, Oberlin, Onio.

Wilson Chapter Publications

CONSIST OF TWO SERIES—OLD AND NEW

The Old Series Comprises the Following Issues:

Ornithologists' and Oologists' Semi-annual, Vols. I and II, two numbers each; Vol. III, one number. (No. 1 of both Vols. I and II are out of print.) 25 cents a number.

Wilson Quarterly, Vol. IV, two numbers. 25 cents a number.

The Journal, two numbers. 10 cents a number.

The whole series (available numbers) $1.

The Whole Series Comprises

Wilson Bulletins, from No. 1 to 48 inclusive. (Nos. 1, 2, 3, 4, 6, 8, and 24 are out of print.) Several numbers are almost gone.

Of the Wilson Bulletin, No. 5 is a Monograph of the Crow, 41 pages, by Frank L. Burns. Price 25 cents. No. 15 is a study of "The Oberlin Grackle Roost," 18 pages, by Lynds Jones. Price 20 cents. No. 30, "Warbler Songs," 56 pages, by Lynds Jones; in which all North American Warblers are discussed, the songs of nearly all described, and a field key to the adult males given. Price 50 cents. No. 31, "A Monograph of the Flicker," 32 pages, by Frank L. Burns. Price 50 cents. No. 38, "A Summer Reconnoissance in the West," by Lynds Jones and W. L. Dawson; being a study of the birds in fourteen states during a journey of 7000 miles. Price 20 cents. No. 37, "A Sectional Bird Census," by Frank L. Burns. Price 25 cents.

The other numbers consist of "General Notes." Price 15 cents each. The whole available new series for $5.00.

Address all communications to

LYNDS JONES, Oberlin, Ohio

University Library
JAN

Old Series Vol. XVI. New Series Vol. XI, No. 4

The Wilson Bulletin

No 49

Published Quarterly by the
WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

December, 1904

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.

CONTENTS

	PAGE
"Kearsarge" Birds, E. H. and H. E. PORTER - -	97
An October All-Day at Blaine, Washington, W. L. DAWSON -	100
A Summer Porch List at Hinsdale, Ill., ESTHER CRAIGMILE -	105
Some Observations on a Captive Red-Tailed Hawk, (*Buteo borealis*), W. F. HENNINGER - - -	107
Brewster Warbler, (*Helmnithophila leucobronchialis*) in Chester County, Pennsylvania, FRANK L. BURNS - -	109
Brewster Warbler Again in Ohio W. F. HENNINGER -	109
Editorial - - - - - - -	111
Election of New Members - - - -	112
Election of Officers for 1905 - - - -	112
Notes from Sandusky, Ohio, E. L. MOSELEY - -	112
Publications Received - - - - -	113

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, Ohio, edited by Lynds Jones.

Subscription: Fifty cents a year including postage, *strictly in advance.* Single numbers 15 cents, unless they are "Special" numbers when a special price is fixed. The BULLETIN, including all "Special" numbers, is free to all paid up members, either Active, Associate or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, 1794 Lanier Ave., N. W. Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for reviews, should be addressed to Lynds Jones, 5623 Drexel Ave., Chicago, Ill.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 15th of the month preceding publication.

THE

WILSON BULLETIN

NO. 49

A QUARTERLY JOURNAL OF ORNITHOLOGY

VOL. XVI DECEMBER, 1904 NO. 4

"KEARSARGE" BIRDS.

BY E. H. AND H. E. PORTER.

Kearsarge, N. H., is situated in a narrow valley of the White Mountains, having Moat Mountain and the noted White Horse ledges on the west, Mt. Kearsarge on the north, and Rattlesnake Range on the east. Through a "notch" to the northwest, the railroad leads to Fabyans and the higher peaks. Three villages are situated in this valley: Kearsarge, North Conway and Intervale. These villages are all interesting from an ornithological standpoint owing to the very varied country, but at Kearsarge village the birds are mainly those of the woods and fields, almost the only water being the diminutive Kearsarge brook.

Our summer home is some two hundred feet from the road, the intervening space being lawn, on which the Goldfinches and Robins are especially numerous. With the exception of a few acres of hay-fields, the remaining twenty-four acres of land are woods, through which Kearsarge brook runs.

This list was compiled during July, August and a part of September. Barely six *long* hunts were made during that period, the birds seen merely chanced to meet our sight, but we always carried glasses. In the list those species marked * were seen on our own place, those marked † breeding there.

In addition to those species named, the Towhee, Black-throated Blue Warbler and Pileated Woodpecker were heard.

1. **Botaurus lentiginosus.** American Bittern. This species was seen once in July, but several times afterwards. Breeds in the Intervale marshes.

2. **Ardea herodias.** Great Blue Heron. Only once seen, the only

record of this species, as far as can be ascertained, in the valley.

3. **Accipiter velox.** *Sharp-shinned Hawk. Commonest of the hawks.

4. **Accipiter cooperi.** *Cooper Hawk. Semi-occasional.

5. **Buteo borealis.** *Red-tailed Hawk. Fairly common during September.

6. **Buteo lineatus.** *Red-shouldered Hawk. One instance noted.

7. **Falco sparverius.** Am. Sparrow Hawk. Breeds on Peaked Mountain of the Rattlesnake Range.

8. **Ceryle alcyon.** †Belted Kingfisher. One within sight or hearing almost constantly.

9. **Dryobates villosus.** *Hairy Woodpecker. Occasional.

10. **Dryobates pubescens medianus.** †Downy Woodpecker. Common in the early summer.

11. **Colaptes auratus luteus.** *Northern Flicker. *Rather rare.*

12. **Antrostomus vociferus.** *Whip-poor-will. One seen.

13. **Chordeiles virginianus.** *Nighthawk. A small flock appeared nearly every evening.

14. **Chætura pelagica.** *Chimney Swift. Common until September 5.

15. **Trochilus colubris.** *Ruby-throated Hummingbird. Uncommon.

16. **Tyrannus tyrannus.** *Kingbird. Common. Breeds at Intervale.

17. **Sayornis phœbe.** *Phœbe. Occasional.

18. **Contopus virens.** Wood Pewee. One instance.

19. **Empidonax flaviventris.** †Yellow-bellied Flycatcher. One family near Kearsarge Brook.

20. **Empidonax minimus.** Least Flycatcher. One seen in August.

21. **Cyanocitta cristata.** *Blue Jay. Infrequent at all times.

22. **Corvus brachyrhynchos.** †American Crow. Comparatively few seen until the first part of September.

23. **Molothrus ater.** Cowbird. One recorded from the Enchanted Woods.

24. **Agelaius phoniceus.** Red-winged Blackbird. Breeds between Kearsarge Village and Intervale.

25. **Sturnella magna.** Meadowlark. One individual remained from July 13 to 17, part of the time at Intervale.

26. **Carpodacus purpureus.** †Purple Finch. A frequent visitor to our lawn, but difficult to find when not there.

27. **Astragalinus tristis.** †American Goldfinch. Common, especially on the lawns.

28. **Pooecetes gramineus.** †Vesper Sparrow. Infrequently seen.

29. **Zonotrichia leucophrys.** White-crowned Sparrow. One instance.

30. **Zonotrichia albicollis.** *White-throated Sparrow. Fairly common on the slopes of the mountains.

31. **Spizella socialis.** †Chipping Sparrow. Abundant.

32. **Spizella pusilla.** †Field Sparrow. Rather rare.

33. **Junco hyemalis.** †Slate-colored Junco. Abundant.

34. **Melospiza cinerea melodia.** †Song Sparrow. Very common occasionally, at other times surprisingly rare.

35. **Cyanospiza cyanea.** *Indigo Bunting. Common along the roads.

36. **Hirundo erythrogaster.** *Barn Swallow.

37. **Petrochelidon lunifrons.** *Cliff Swallow.

Two more surprisingly infrequently recorded species.

38. **Ampelis cedrorum.** †Cedar Waxwing. Very common.

39. **Lanius borealis.** *Northern Shrike. Once recorded.

40. **Vireo olivaceus.** †Red-eyed Vireo. Commonly seen near the house.

41. **Vireo flavifrons.** *Yellow-throated Vireo. By far the commonest of the Vireos.

42. **Vireo solitarius.** *Blue-headed Vireo. Often seen, especially in September.

43. **Vireo noveboracensis.** *White-eyed Vireo. Once recorded, the only time in the county, as far as can be ascertained.

44. **Mniotilta varia.** *Black-and-white Warbler. Infrequent.

45. **Helminthophila rubricapilla.** *Nashville Warbler. Commoner than the previous species.

46. **Helminthophila peregrina.** *Tennessee Warbler. Twice seen.

47. **Dendroica coronata.** *Myrtle Warbler. Rather common.

48. **Dendroica maculosa.** †Magnolia Warbler. Somewhat more common.

49. **Dendroica blackburniæ.** Blackburnian Warbler. We saw one in very thick woods—a beautiful male. His song was almost precisely that of the Black and White Warbler, and the song was almost all on the same note, contrary to the usual one.

50. **Dendroica virens.** †Black-throated Green Warbler. By far the commonest of the Warblers, and commonest of all Kearsarge birds with the exception of the Chickadees, Robins, Bluebirds, Song Sparrows, Chippies, Crows and Juncos.

51. **Dendroica vigorsii.** *Pine Warbler. Rather rare.

52. **Seiurus aurocapillus.** *Oven-bird. Fairly common.

53. **Geothlypis trichas brachidactyla.** *Northern Yellow-throat. Rare.

54. **Wilsonia canadensis.** Canadian Warbler. Twice seen.

55. **Setophaga ruticilla.** † American Redstart. Common until August 15.

56. **Galeoscoptes carolinensis.** *Catbird. Casual.

57. **Toxostoma rufum.** *Brown Thrasher. Once recorded.

58. **Certhia familiaris americana.** *Brown Creeper. Rare at all times.

59. **Sitta carolinensis.** *White-breasted Nuthatch. Fairly common.

60, **Sitta canadensis.** *Red-breasted Nuthatch. Rarer than the previous species.

61. **Parus atricapillus.** †Chickadee. Common.

62. **Parus hudsonicus.** *Hudsonian Chicadee. Very rare.

63. **Regulus satrapa.** *Golden-crowned Kinglet. Common in September.

64. **Hylocichla mustelina.** †Wood Thrush. Rare.

65. **Hylocichla fuscescens.** *Wilson Thrush. Commoner.

66. **Hylocichla swainsonii.** *Olive-backed Thrush. Once seen.

67. **Hylocichla guttata pallasii.** *Hermit Thrush. Commonest of the thrushes.

68. **Sialia sialis.** *Bluebird. Common.

69. **Passer domesticus.** English Sparrow. Only once seen in Kearsarge village.

In addition to these, nineteen species were recorded from the village by four other bird-lovers, including Warbling Vireo.

The most remarkable facts are the extreme abundance of the Black-throated Green Warbler, the infrequency of all warblers and the absence of the Northern Parula Warbler, Scarlet Tanager, all Grouse and the Yellow-bellied Sapsucker.

AN OCTOBER ALL-DAY AT BLAINE, WASHINGTON.

BY WILLIAM LEON DAWSON.

Of course it was cloudy. That it was so instead of *rainy* was a mark of special favor, received by the Bird-Man with becoming gratitude, for every loyal Puget-Sounder knows that rain has the right of way from October first *on*. The clouds formed a great leaden canopy centering over Blaine, with no visible support by way of tent poles upon either horizon, but with certain airy hangings and draperies at the edges which not even the rising sun could thrust aside. But in his attempt to do so the draperies caught fire, warming from saffron to Saturn red and rolling up from the east in billows of flame which threatened the eternal hills. Mt. Baker, our patron saint and watchful sentinel, situated some fifty miles to the east and south, stood apart from the conflict, but reflected something of the heavenly ardor from its new-fallen snows until the sun found a rift in the curtain of cloud and shot a full glance at the mountain, whereupon it cast a huge shadow athwart the sky, like the umbration of the needle upon the dial. The northeast in glory and the southeast in the shadow of our glorious mountain—it was worth the sacrifice of a few early birds to have seen it !

The first half hour of indecision, from six o'clock on, was spent awheel, ranging the sidewalks of the still silent town, gleaning the bird-notes from orchard and garden and unreclaimed ravine. The writer was just congratulating himself upon the abundance of native birds hereabouts, undisturbed

as yet by the presence of the all but universal scourge, when "*Yark, yark, scrape, chirp, chirp*" came from a little fir-tree a block away, in the center of the central business block of town. His obscene majesty, the English Sparrow, has arrived! His half century of conquest is appropriately consummated upon this last square rood of Uncle Sam's undivided possession. Blaine stands at the exact northwestern corner of the United States, and the goal of the Gamin is reached.

The aliens can be very crafty if they choose, and they do choose while they are in the minority. I have searched the townsite over, during the two months past, fearing the dread presence, yet resolved to know the worst, without having heard a single chirp from the *domesticus* until to-day. Yet I am assured that the creatures have been here in small numbers for two or three years past. Nine Sparrows were frightened, upon this occasion, from the little fir-tree, and all became instantly silent upon their escape.

The cataloguer's attention was next directed toward the water birds. Drayton Harbor, which is the inner sea sanctum of Blaine, is some seven miles in circumference at high-tide, and save for a narrow channel, is shut off from the wide waters of Semiahmoo Bay by a sand-spit a mile in length. As flood-tide approaches, the number of sea birds upon the harbor is augmented to several thousand by the arrival of ducks, mostly Scaups, "Bluebills," and Scoters, or "Black Ducks." These birds, upon entering or leaving the harbor, usually fly low over the sand-spit and are here assaulted by a battery of ever-ready guns. As a result of the steady maintainance of the firing line, the ducks upon the harbor are nervous and unapproachable—save by eight-power binoculars.

The constant residents of the harbor are Grebes. Western Grebes (*occidentalis*), to the number of a hundred or more, move about singly or in small groups, occasionally calling to each other in shrill notes, like the squeaking of rusty windlasses. They are graceful creatures—cruelly graceful with their dagger beaks and eyes of fiery red—and yet there is something swan-like in the carriage of the head upon the mobile neck.

Holbœll Grebes (*Colymbus holbœllii*), appear in lesser

numbers and are less conspicuous by reason of their compact build and blended colors.

Horned Grebes (*Colymbus auritus*), abound. As a rule they frequent the shallows, where escape by diving is not always convenient; but, unlike their congeners, they take to wing with great readiness. At times they will fly back and forth upon the merest whim, or for exercise.

The presence of a school of smelt or herring will attract all the Grebes at once. On this all-day trip the writer drew near a busy company of several hundred birds, and when they took flight the pattering of tiny feet and the dipping wing-tips was like the fall of a small cloud-burst upon the water.

Gulls, at high-tide, are like street car conductors off duty. There is nothing for them to do but to haunt unavailingly the scenes of their former activity, or to gather in languid companies and discuss the prospects of the next shift. At such times they are glad to find log-booms or floats to rest upon; but in the absence of these (as at present in Drayton Harbor) they sit upon the water or drift about on pieces of mill waste, or else desert the harbor altogether.

Bonaparte Gulls (*Larus philadelphia*), which have swarmed to the number of thousands for six weeks past, are now represented by a few stragglers, wounded birds and their faithful mates; while Ring-billed Gulls (*Larus delawarensis*) have become common, and California Gulls (*Larus californicus*) are slowly increasing in numbers.

The Terns, Arctic (?) and Common (*Sterna paradisæa* and *hirundo*) have disappeared, and after them has apparently followed their arch-persecutor, the Parasitic Jaeger (*Stercorarius parasiticus*), a specimen of which I shot from "the spit" on September 30th.

In contrast with all this was the trip into the interior, made in the afternoon. The gray cloud-cap still hung over everything, but it included within its dome distant Baker, whose uplifting presence made one feel that his world was good enough, however circumscribed.

Mile after mile was done off over gravel pikes, past stump-scarred clearings, tiny orchards and deserted townsites, through somber forests of fir and hemlock, and through wildernesses of second growth, alders, willows and evergreens.

But the bird-world was oppressed by the cloud-cap. It had had its breakfast, and, since there was not light enough to encourage thoughts of supper, it betook itself early to bed. There was little to be heard except the *tick, tick, tick* of the Western Winter Wren (*Olbiorchilus hiemalis pacificus*) and the drowsy *tss, tss; tss tsee* of the Western Golden-crowned Kinglet (*Regulus satrapa olivaceus*).

At 3 p. m., as I stood upon a little bridge over a ravine choked full with a jungle of vine maple, goooseberry bushes, devil's club, and ferns, an incautious chickadee, grumbling over its hard piliow, led the Bird-Man to attempt the chickadee call. (The notes of these Oregon Chickadees, *Parus atricapillus occidentalis*, have a slightly different pitch from those of *Parus atricapillus*, and I have not quite mastered them yet). As a result, the bushes began to yield up sleepy Chickadees. Western Golden-crowned Kinglets emerged unexpectedly from snug sleeping-bags hidden in the vegetation under my feet. The Northwest Bewick Wren (*Thryomanes bewickii calophonus*) scolded, W. W. W. (which is note-book shorthand for Western Winter Wren) ticked apprehensively, and the fruitful bush began to yield a tribute of Ruby-crowned Kinglets, aroused from deeper slumbers, but cheerful and forgiving still. It remained only for the *Maior domo*, which in these parts is the Oregon Towhee (*Pipilo maculatus oregonus*), to peep out and through the curtains to see that the children were not taking harm, and then the disturber of midnight peace, at 3:00 p. m., withdrew "in good order."

But even after this disheartening adventure, Fortune perched upon the Bird-Man's banner in the shape of a California Pygmy Owl (*Glaucidium gnoma californicum*). This pocket edition of the powers that prey stood out boldly upon the topmost splinter of a wayside stub and challenged scrutiny. The gnoma gave his back to the road, but every now and then turned a careful eye upon the stranger. Then all at once the bird whirled backward and launched himself like a bolt across the road, at a mouse some sixty feet away. Seizing the "wee, timerous, cowerin' beastie" at the very entrance of its hole, the bird maintained its grasp with both feet and supported itself by wings outstretched upon the ground. Not until the squeakings of the victim had quite ceased, did the captor rise

and disappear by rapid flight into the wood.

There are always a few staple species which try the patience of the ardent horizonist by deserting on the "All-Day." Formerly they have been the most inevitable of birds afield, but now, forsooth, they must lurk in hiding with the most cunning intent to defeat the ends of science. Among such that I would hold up on this occasion to well deserved scorn, are Kingfisher (*Ceryle alcyon*), (think of it!), Audubon Warbler (*Dendroica auduboni*), and Herring Gull (*Larus argentatus*). Others which might have been gracious are Mongolian Pheasant (*Phasianus torquatus*), Northern Pileated Woodpecker (*Ceophlœus pileatus abieticola*), Harris Woodpecker (*Dryobates villosus harrisii*), and California Creeper (*Certhia familias occidentalis*). With these additions I feel that the "All-Day" would have been fairly representative of the vicinity of Blaine at this season.

Horizons, Blaine and vicinity, October 13, 1904, 5. a. m. to 5 p. m.:

BLAINE, 5:00 TO 8:00 A. M.

Rusty Song Sparrow,—*Melospiza cinerea morphna.*
Northwest Crow,—*Corvus caurinus.*
Northwest Bewick Wren,—*Thryomanes bewickii calophonus.*
Western Meadowlark,—*Sturnella magna neglecta.*
Western Robin,—*Merula migratoria propinqua.*
California (?) Finch,—*Carpodacus purpureus californicus.*
Oregon Junco,—*Junco hyemalis oregonus.*
Western Golden-crowned Kinglet,—*Regulus satrapa olivaceus.*
Western Winter Wren,—*Olbiorchilus hiemalis pacificus.*
English Sparrow,—*Passer domesticus.*
American Pipit,—*Anthus pensilvanicus.*
Oregon Towhee,—*Pipilo maculatus oregonus.*

DRAYTON HARBOR, 8:00 TO 10:00 A. M.

Horned Grebe,—*Colymbus auritus.*
Western Grebe,—*Echmophorus occidentalis.*
White-winged Scoter,—*Oidemia deglandi.*
Greater Scaup Duck,—*Aythya marila.*
Lesser Scaup Duck,—*Aythya affinis.*
Bonaparte Gull,—*Larus philadelphia.*
Holbœll Grebe,—*Colymbus holbœllii.*
Western Gull,—*Larus occidentalis.*
Surf Scoter,—*Oidemia perspicillata.*
Marbled Murrelet,—*Brachyramphus marmoratus.*
Glaucus-winged Gull,—*Larus glaucescens.*

Ruddy Duck,—*Erismatura jamaicensis.*
Red-breasted Merganser,—*Merganser serrator.*
Loon,—*Gavia imber.*

SEMIAHMOO POINT (The Sand Spit), 10:00 TO 12:00 A. M.

Western Bluebird,—*Sialia mexicana occidentalis.*
Nuttall Sparrow,—*Zonotrichia leucophrys nuttalli.*
Northwest Flicker, —*Colaptes cafer saturatior.*
Sandwich Sparrow,—*Ammodramus sandwichensis alaudinus.*
Ring-billed Gull, —*Larus delawarensis.*

BLAINE BEACH, 12 M.

California Gull, —*Larus californicus.*
Northwest Coast Heron,—*Ardea herodias fannini.*

BLAINE-LINDEN ROAD, 1:30 TO 4:00 P. M.

Steller Jay, —*Cyanocitta stelleri.*
Oregon Ruffed Grouse,—*Bonasa umbellus sabini.*
Oregon Chickadee,—*Parus atricapillus occidentalis.*
Pine Siskin,—*Spinus pinus.*
Ruby-crowned Kinglet,—*Regulus calendula.*
California Pygmy Owl, —*Glaucidium gnoma californicum.*
Gairdner Woodpecker,—*Dryobates pubescens gairdneri.*

A SUMMER PORCH LIST, AT HINSDALE, ILL.

BY ESTHER CRAIGMILE.

It was almost discouraging to look forward to a summer without bird tramps, but that was the outlook from the end of June to the middle of August, 1904. But there is something to be seen and heard, even from one's own door, if eyes and ears have been trained. The location was a few miles west of Chicago, half a mile distant from Flag creek. To the east, west and north, rolled the fertile prairies, while a wood of small trees stretched to the south, bordering the creek until it reached the Des Plaines river, four miles distant.

A dense thicket, several rods in length, grew along the roadside, furnishing food and shelter for not a few birds. Song Sparrows, Chippies, Catbirds, Thrashers, Chewinks, Cowbirds, and Indigo Buntings were always in evidence there. Numbers of water birds were to be seen mornings and evenings, going to and from their feeding grounds.

Previous to this year the fields have been alive with hundreds of Dickcissels. No nests were found this summer,

and their notes were only heard twice on two successive days late in June. "What has become of Dick?" was a common question among bird friends.

The Yellow Warbler has always been abundant until this year. Not one was seen or heard, so the Song Sparrows were alone responsible for the young Cowbirds. It was not an uncommon sight to see a huge young Cowbird pursuing a Song Sparrow along the wire fence, refusing to be hushed until it had been gorged with food.

It seemed a little strange to miss both these birds this summer. Heretofore their presence has been more marked than any other variety on this list.

The most unique experience of the summer was a morning serenade. All the Bartramian Sandpipers of the creek bottom must have been present. It was barely three o'clock, and their weird, wind-like whistles sounded like music from another planet.

The Nighthawks did not appear until the middle of August, and then in migration for the most part. One bright midday a flock of one hundred were seen flying south at considerable height. Near this same spot two years ago, late in August, I counted six hundred Nighthawks in a space of fifteen minutes. It was about six o'clock in the evening, and the air was just alive with them. The area in which I counted was not more than a quarter of a mile in width, and it was impossible to estimate the actual size of the wave. It was a continuous passage—there was no looking back on their part—and standing in an open tract in the woods, I was able to count those in my range with some degree of accuracy.

Here is the summer list in the order in which they were seen or heard:

1. Bartramian Sandpiper.
2. Indigo Bunting.
3. Dickcissel.
4. Song Sparrow.
5. Vesper Sparrow.
6. Grasshopper Sparrow.
7. Goldfinch.
8. Wood Pewee.
9. American Bittern.
10. Red-winged Blackbird.
11. Crow.
12. Blue Jay.
13. Barn Swallow.
14. Chimney Swift.
15. Kingbird.
16. Bob-white.
17. Flicker.
18. Catbird.
19. Brown Thrasher.
20. Red-headed Woodpecker.

21. Cowbird.
22. Bobolink.
23. Mourning Dove.
24. Meadowlark.
25. Yellow-billed Cuckoo.
26. Robin.
27. Bluebird.
28. Field Sparrow.
29. Marsh Hawk.
30. Prairie Horned Lark.
31. Purple Martin.
32. Baltmore Oriole.
33. Red-eyed Vireo.
34. Rough-winged Swallow.
35. Northern Yellow-throat.
36. Screech Owl.
37. Great Blue Heron.
38. Black-crowned Night Heron.
39. Migrant Shrike.
40. Hairy Woodpecker.
41. Downy Woodpecker.
42. Chickadee.
43. Bronzed Grackle.
44. Chewink; Towhee,
45. Scarlet Tanager.
46. American Sparrow Hawk.
47. Pigeon Hawk.
48. Nighthawk.
49. Semipalmated Plover.

SOME OBSERVATIONS ON A CAPTIVE RED-TAILED HAWK (*Buteo borealis*).

BY W. F. HENNINGER.

On August 15th of this year a Red-tailed Hawk was brought to me, that had been shot in the wing while pouncing down on a chicken in a barn-yard. Just a few days before that another had been shot at at the same place, while with a chicken in his talons he was trying to get to an orchard. On August 31st I saw one pursuing a Red Squirrel. This goes to confirm my statement made elsewhere (Wilson Bulletin, December, 1902, p. 138) that the Red-tailed Hawk feeds principally on birds and squirrels. The wounded Hawk brought me is not an adult and I concluded to experiment with my captive to find out just what he would eat. After refusing food for three days he began to eat. So far he has eaten beef, raw, boiled and roasted; also raw and boiled pork, kidney and liver. He would not touch cured meats; nor will he eat veal, except forced by hunger. A Barn Owl, which I had in captivity for a week in 1898, would also refuse veal. He always liked squirrel, whenever offered him. On September 1st I gave him two Yellow-legs and two Field Sparrows, and although he had just been fed with some raw beef, immediately devoured them with great greed; in fact, he would always jump down from his perch and with great ferocity go for birds

at once. He always began with the head, picking it clean, and then tearing it off with his bill would crush it at once and devour it, after which the rest of the bird would follow. He always preferred birds to anything else. The second week in September a Sora was brought to me. This I let lie for three days, till, in the heat, it had begun to decay and smelled very badly. After he had literally stuffed himself with raw meat, about one pound, I threw this decayed Sora in his cage. He went at it at once and ate it in about five minutes. This, in my opinion, does not merely show that the Red-tailed Hawk will eat carrion, and most any bird of prey, when hungry, will do that, but, as he certainly was not hungry at the time, that Red-tails prefer birds to other food. Again, I do not like to make a statement to this effect, as there is so much evidence to the contrary (compare: Hawks and Owls, by A. K. Fisher,———), but it certainly agrees with my observations in Pike and Scioto Counties, as well as my field observations in Seneca County. On September 29th I put a large dead rat in his cage. He looked at it for about ten minutes before touching it, and I hadn't fed him for two days at that. Then he ate the head and the inside, leaving the skin, hind legs and tail intact, and would not eat this till the next day, though I did not feed him anything else. This certainly does not show a great fondness for rats and mice. He never went at them with the greed he showed for squirrels and birds. On October 21 and 22 I fed him raw and fried fish, which he ate immediately. So far I have never been able to get him a snake or frogs, but if I can keep him over the winter will try him on these next spring. Thus I cannot consider the Red-tail as harmless as other ornithologists do, yet would not agree with the majority of sportsmen in calling him a harmful species, as killing of squirrels is not a detriment, but a beneficial service. From a utilitarian standpoint, I can only hope for the extermination of the squirrels, especially the Red Squirrel, though not from an aesthetic or sentimental standpoint.

THE NEW YEAR CENSUS.

Not long after this number reaches its readers a new year will be at hand. If you are keeping a yearly record of the birds, your old note-book will be laid aside and a fresh one placed in readiness. It means something æsthetically, if not practically, how the new note-book is begun, whether the first page is well filled or not. To me it means more than I care to admit. We have begun these New Year Censuses well, and I hope and trust that 1905 will not fall behind 1904 in the records that shall be made. 1905 begins on Sunday, and it is therefore proposed that for those who do not study birds on that day, to make the record on the 2nd. How much each one can do will depend upon the region, the time, and the weather. The editor will be unable to participate in this contest, so the prize offered last year cannot be repeated. To every one who secures a bona fide list of twenty-five or more species, not including English Sparrow, a year's subscription to THE WILSON BULLETIN will be given. Make as large a list as possible for Mr. Frank M. Chapman's Christmas Census, and then beat it for the New Year Census! Send the lists to Lynds Jones, 5623 Drexel Avenue, Chicago, Ill.

BREWSTER WARBLER (*Helminthophila leucobronchialis*) IN CHESTER COUNTY, PENNSYLVANIA.

Near the close of an "all-day with the birds" I was fortunate enough to meet with a single individual of this type, phase, or whatever it may be, near Devon, on May 13th, 1904. It was found at the border of a large grove, where I watched it sometimes as near as twenty feet, from an old cart-road. It was altogether like the Blue-winged Warbler above,—with the white wing bars and black bar through the eye—but the under parts were white instead of the rich yellow of that bird. I could detect no trace of yellow on its breast. In action, dropping from branch to branch of the smaller trees so near, it was very like the Blue-wing—with which I am familiar. Although I cannot but regret that I had no means of securing it at the time, I have no hesitation in recording it as a Brewster's Warbler, after I had observed it through a good pair of field glasses for over fifteen minutes. FRANK L. BURNS, Berwyn, Pa.

BREWSTER WARBLER AGAIN IN OHIO.

(*Helminthophila leucobronchialis*)

BY W. F. HENNINGER.

On September 17th of this year while out to study the annual fall

migration in a very favorable woods 1½ miles southeast of Tiffin we were able to add several species to the county list. Of warblers 18 species were seen, among them a Brewster Warbler. While kneeling down watching a Black and White Warbler catching a moth, I suddenly spied in a low sapling a warbler that at the first glance I recognized as a new species.

The upper parts were bluish gray with a black streak through the eye, lower parts yellowish shading into white on forepart of breast, throat and chin, two small yellowish wing bands, yellowish crown patch hardly discernible, the whole plumage overlaid with a kind of greenish cast as if some one had breathed over it. The bird was observed for about 8 or 9 minutes, was very agile, but uttered no sound, not even the usual migration "chip" of other warblers. The bird was perhaps not quite typical but it was a Brewster nevertheless and to my knowledge the first fall record for this species in the state, and I was only sorry I had left my gun at home.

THE WILSON BULLETIN

A Quarterly Magazine Devoted to the Study of Living Birds
Official Organ of the Wilson Ornithological Club

Edited by LYNDS JONES

PUBLISHED BY THE CLUB, AT OBERLIN, OHIO

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number postpaid.
Price in all countries in the International Postal Union, 65 cents a year, 20 cents a number.
Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr. Frank L. Burns, Berwyn, Penn., or to Mr. John W. Daniel, Jr., Washington, D. C.

EDITORIAL.

Readers will be pleased to know that the Collection of Mr. J. Warren Jacobs which was displayed at the St. Louis Fair and mentioned in the last Bulletin, received from the committee on awards a gold medal as a just recognition of the expense of time and money as well as the discrimination which Mr. Jacobs has put upon his unusually complete collection.

The local papers of Chardon, Ohio, report a flight of some 500 Passenger Pigeons at that place on October 22. It is unfortunate that this report was not confirmed in a manner above question. The presence of so many of these once abundant birds is difficult to account for. No breeding place of any such extent as this would indicate are known, if, indeed, any at all are known. Numerous reports of considerable numbers in the Northwest, in South America, and even in foreign countries where they never lived, have proved to be myths.

Mr. Frank L. Burns, of Berwyn, Pa., has kindly consented to assist the editor in collecting copy for the March Bulletin. Matter intended for publication in that number should reach Mr. Burns before February 15.

The editor will be at 5623 Drexel Ave., Chicago, Ill., until July, 1905. Requests for back numbers of the Bulletin, or notices that the current number has not been received should be addressed to Mrs. Lynds Jones, 160 North Professor St., Oberlin, Ohio, to insure prompt attention.

We want to inaugurate a campaign for increasing our membership the coming year, to twice its present size. This can be done if every member will either induce some friend to become a member, or send to any one of the officers a list of names of persons who are interested in birds. Will not every member make this a personal matter? Increased membership means a better grade of work, and a better Bulletin.

ELECTION OF NEW MEMBERS.

The following persons are presented for membership:

FOR ACTIVE MEMBERSHIP.

P. A. Taverner, Detroit, Mich.: Carl Fritz Henning, Boone, Iowa; Frederick B. McKechnie, Ponkapog, Mass.

FOR ASSOCIATE MEMBERSHIP:

Miss Clara S. Wheatley, Medina, Ohio.

ELECTION OF OFFICERS FOR 1905.

The election of officers has resulted as follows:

President—Lynds Jones, Oberlin, Ohio.

Vice-President—W. L. Dawson, Blaine, Wash.

Secretary—John W. Daniel, Jr., Washington, D. C.

Treasurer—Frank L. Burns, Berwyn, Penn.

Executive Council. H. C. Oberholser, John H. Sage, A. W. Blain, Jr.

It is in order to send your annual membership dues to the treasurer to the address given above.

Matter intended for the March Bulletin may be sent to Mr. Frank L. Burns, Berwyn, Pa., or to the editor at 5623 Drexel Ave., Chicago, Ill.

NOTES.

NOTES FROM SANDUSKY, OHIO. Mr. Webster Ransom reports a Sycamore Warble, (*Dendroica dominica albilora*) on April 24.

I have secured two specimens of Red-legged Black Duck (*Anas obscura rubripes*) this spring.

Two specimens of Widgeon (*Mareca penelope*) were mounted by John Herb this spring.

James Galloway informs me that about twenty years ago James Dildyne shot six King Eiders (*Somateria spectabilis*) at one shot.

Miss Edith Dixon, of Milan, reports a Blue Grosbeak (*Guiraca cærulea*) at that place, well identified, but the specimen was not secured.

William Harting caught an uninjured Horned Grebe (*Colymbus auritus*) asleep, on Sandusky Bay near Cedar Point, on April 13. The bird was brought home and fed, and on April 16, was taken to the bay and released. It was caught twice during the day. It seemed to have no fear of him.

The specimen of Jaeger shot at Sandusky October 10, 1889, and reported in Cook's Birds of Michigan, page 27 and requoted in Jones' Birds of Ohio, page 26 proves to be a Parasitic Jaeger (*Stercorarius parasiticus*).

Webster Ransom reports a parrot that frequented his orchard, six miles south of Sandusky, in the summer of 1903. It resembled a Carolina Paroquet (*Conurus carolinensis*).

E. L. MOSELEY, Sandusky, Ohio.

PUBLICATIONS RECEIVED.

A Preliminary Review of the Birds of Nebraska, with synopses. By Lawrence Bruner, Robert H. Wolcott, Myron H. Swenk.

While this long promised "Birds of Nebraska" comes to us as a "Preliminary Review," and therefore much different in character from what one might be led to expect, seems to be well adapted to its avowed purpose, "A Working List." While it is preliminary in character, the claim of the authors that it "is believed to represent exactly the state of our knowledge at the present day." seems to be borne out by an examination of the list of species and annotations. The book is really a series of artificial keys, each key followed by an annotated list of the species covered by that key. The keys are intended for work with the specimen in hand, and to the writer's mind, will prove somewhat difficult to those unfamiliar with this method of identification. They belong to the older, rather than the newer system. However, properly used, they will identify. There is a complete index, but we miss a summary of any sort. Such a summary as we find in "The Birds of Colorado," would be a valuable addition. The book is bound in cloth and makes a very welcome and valuable addition to faunal literature.

L. J.

North American Fanna. No. 23. November 24, 1904. A Biological Reconnaisance of the Base of the Alaska Peninsula. By Wilfred H. Osgood.

In this paper of 86 pages, including index, we are given a glimpse of the physical features of the region treated in two maps and eleven half-tones, and in the word pictures which occupy a prominent part of the introduction. Some 43 mammals are given as occurring in the region, and 137 birds are listed. In compiling the lists, previous work is drawn upon It is of interest to note that the bird list contains no new species or sub species. One born to such work cannot avoid a feeling of slight envy of those who are fortunate enough to get out into such remote and interesting fields while he congratulates the author upon his good work.

L. J.

A Discussion of the Origin of Bird Migration. By P. A. Travener. From The Auk, Vol. XXI, No. 3, July, 1904, pp. 322-333.

The author discusses the whole question of migration, but strongly emphasizes one point in previous discussions which has remained obscure. It is that the necessity for the northward movement was at first brought about (and the same necessity continues to the present time) by the fact that during the winter the tropics contain all the bird life they can support without the enormous increased demand for food which would result from the appearance of nestfulls of young. The southward movement began because returning cold destroyed the food north. Of course the laws of adaptation and the survival of the fittest must be brought in to complete the argument. The paper is a valuable one in bringing into prominence a factor which must have exerted a determining influence at the beginnings of this vast movement.

L. J.

Amateur Sportsman, Vol. XXX1, Nos. 6, 7, 8.
American Ornithology, Vol. IV, Nos. 9, 10, 11, 12.
Bird-Lore, Vol. VI, Nos. 5, 6.
Boys and Girls, Vol. II, No. 4; Vol. III, Nos. 1, 2, 3.
Bulletin of the Michigan Ornithological Club, Vol. V, No. 3.
Bulletin 153, Ohio Agricultural Experiment Station.
Bulletin 3, Ohio Department of Agriculture.
By The Wayside, Vol VII, No. 4.
Condor, The, Vol. VI, Nos. 5, and 6.
Maine Sportsman, Vol. XII, Nos. 134, 135
Monthly Bulletin, The, Vol. II, Nos. 6, 7.
Naturaliste Canadien, Le, Vol. XXXI, Nos. 9, 10, 11.
Ohio Naturalist, The, Vol. V, No. 1.
Oologist, The, Vol. XXI, No. 10.
Ornithologische Monatschrift, Vol. XXIX, Nos. 9, 10.
Plant World, The, Vol. VII, Nos. 9, 10, 11.
On a Collection of Birds and Mammals from Mount Sanhedrin, California. By Witmer Stone. From Proceedings of The Academy of Nat. Sci. of Philadelphia, July, 1904.

ERRATA.

Page 13, line 8, for "autumn" read "summer."
Page 19, line 32, " "1904" read "1903."
Page 35, line 10, " "*holbœlli*" read "*holbœllii.*"
Page 33, line 20, " "*Synthliborhamphus*" read "*Synthliboramphus.*"
Page 36, line 23, " "*ocrgonus*" read "*oregonus.*"
Page 36, line 27, " "*itaca*" read "*iliaca.*"
Page 94, line 31, " "permiting" read "permitting."
Page 95, lines 16, 17, 18, for "occurence" read "occurrence.;'

INDEX.

Acanthis linaria, 36.
Accipiter atricapillus, 83.
cooperi, 6, 35, 98.
velox, 35, 98.
Actitis macularia, 6.
Æchmophorus occidentalis, 33, 104.
Aix sponsa, 6, 34.
Agelaius phoniceus, 7, 98.
caurinus, 35.
Ammodramus sandwichensis alaudinus, 36, 105.
Ampelis cedrorum, 7, 36, 99.
Anas boschas, 34, 82.
obscura, 82.
rubripes, 112.

obscura + boschas, 22.
Anser albifrons gambeli, 34.
Anthus pensilvanicus, 104
Antrostomus vociferus, 6, 98.
Ardea herodias, 6, 34, 97.
fannini, 105.
virescens, 34.
Asio accipitrinus, 35.
Astragalinus tristis, 7, 98.
salacamans, 36.
Auklet, Rhinoceros, 33.
Aythya affinis, 34, 104.
americana, 34.
marila, 34, 104.
vallisneria, 34.
Bittern American, 6, 19, 34, 48, 75, 78, 97, 106.
Least, 21, 48, 75, 78' 93.
Blackbird, Brewer, 35.
Northwest Red-winged, 35.
Red-winged, 7, 17, 21, 41, 49, 51, 56, 79, 98, 106.
Bluebird, 9, 18, 20, 24, 42, 50, 58, 59, 80, 99, 100, 107.
Western, 37, 105.
Bob-white, 16,17, 18, 19, 20, 34, 39, 40, 48, 58, 59, 78, 94, 106.
Bobolink, 41,49, 58, 59, 79, 92, 107.
Bonasa umbellus, 6.
sabini, 105.
togata, 35.
Botaurus lentiginosus, 6, 34, 97.
Brachyramphus marmoratus, 104.
Branta canadensis occidentalis, 34, 97.
Bubo virginianus saturatus, 35.
Bufflehead, 17, 34, 82.
Bunting, Indigo, 7, 41, 49, 59, 79, 99, 105, 106.
Lark, 92.
Bush-Tit, 37.
Buteo borealis, 6, 98, 107.
calurus, 35.
lineatus, 6, 98.
Butorides virescens, 6.
Calcarius pictus, 85.
Canvas-back, 34.
Cardinal, 17, 18, 19, 20, 41, 45, 49, 79.
Carpodacus purpureus, 7.
californicus, 35, 104.
mexicanus frontalis, 24.
Catbird, 8. 24, 43, 45, 50, 59, 80, 99, 105, 106.

Ceophlœus pileatus abieticola, 6, 35, 104.
Cepphus columba, 33.
Certhia familiaris americana, 8, 99.
occidentalis, 104.
Ceryle alcyon, 6, 35, 83, 98, 104.
Chætura pelagica, 6, 98.
Charitonetta albeola, 34, 82.
Chat, Yellow-breasted, 42, 50, 59, 80.
Chen hyperborea nivalis, 21, 34.
Chewink, 105, 107.
Chickadee, 8, 16, 17, 18, 19, 20, 24, 42, 50, 56, 59, 80, 99, 107.
Hudsonian, 99.
Oregon, 37, 103, 105.
Chordeiles virginianus, 6, 98.
henryi, 35.
Cistothorus palustris paludicola, 37.
Clangula clangula americana, 34.
Coccothraustes vespertinus montanus, 36.
Coccyzus americanus, 6.
erythrophthalmus, 6.
Colaptes auratus luteus, 6, 98.
cafer saturatior, 35.
Colinus virginianus, 34.
Columba fasciata, 35.
Colymbus auritus, 33, 102, 104, 112.
holbœllii, 33, 66, 57, 101, 104.
nigricollis californicus, 33.
occidentalis, 101.
Compsothlypis americana usneæ, 8,
Contopus virens, 7, 98.
Conurus carolinensis, 112.
Coot, American, 21, 22, 34, 69,75.
Cormorant, Brandt, 34.
Double-crested, 74.
Pelagic, 34.
Corvus americanus, 35.
brachyrhinchos, 98.
caurinus, 35, 104.
Cowbird, 7, 32, 41, 49, 79, 98, 105, 106, 107.
Crane, Sandhill, 75.
Creeper, Brown, 8, 17, 18, 19, 20, 42, 59, 99.
California, 104.
Crow, American, 16, 17, 18, 19, 20, 24, 41, 49, 51, 59, 79, 98, 99, 106.
Fish, 20.
Northwest, 35, 104.
Cuckoo, Black-billed, 6, 40, 48, 56, 59.
Yellow-billed, 6, 40, 48, 56, 59, 78, 107.
Curlew, Long-billed, 34.
Cyanocitta cristata, 7, 98.
stelleri, 35, 105.

Cyanospiza cyanea, 7, 99.
Dendroica æstiva. 8. 36.
auduboni. 36. 104.
blackburniæ, 8, 99.
cærulescens. 8.
dominica albilora. 112.
coronata. 8. 36. 99.
maculosa. 8. 99.
nigrescens. 36.
pensylvanica, 8.
rara, 3.
striata, 8.
vigorsii. 8. 99.
virens, 8. 99.
Dickcissel. 41. 49. 52. 53, 54, 58. 59. 79. 105. 106.
Dove. Mourning. 18, 19, 24, 39, 40, 48, 51, 59, 78, 107.
Dryobates pubescens medianus, 6. 98.
gairdneri. 35. 105.
villosus, 6. 98.
harrisii. 35. 104.
Duck, Black, 82. 101.
Lesser Scaup. 34. 48. 101. 104.
Harlequin, 34.
Red-legged Black, 17. 112.
Ruddy, 105.
Scaup, 17, 34. 101. 104.
Wood, 6, 34. 78.
Eagle. Bald. 6, 40. 48.
Golden, 19.
Eider, King. 112.
Empidonax flaviventris, 98.
minimus, 7, 98.
Ereunetes occidentalis, 34.
Erismatura jamaicensis. 105.
Falco sparverius, 6. 98.
deserticola, 35.
Falcon, Peregrine, 22.
Finch, California Purple, 35. 104.
House, 27.
Purple, 7. 20. 41, 49, 59, 79.
Flicker, Northern. 6, 17. 18, 19, 24, 40, 48, 59, 78. 98, 106.
Northwestern, 35.
Flycatcher, Alder, 41, 49.
Crested, 6, 24, 41, 49, 79.
Green-crested. 41, 49, 79.
Least, 7, 41, 49. 59. 79. 98.
Olive-sided. 7.

Yellow-bellied, 41, 46, 57, 59, 98.
Fulica americana, 34.
Gadwall, 22, 74.
Galeoscoptes carolinensis, 8, 99.
Gallinago delicata, 34.
Gallinule, Florida, 21, 59, 75, 93.
Gavia imber, 6, 33, 105.
lumme, 33.
pacifica, 33.
Geothlypis tolmiei, 36.
trichas brachydactila, 8, 99.
Glaucidium gnoma californicum, 103, 105.
Gnatcatcher, Blue-gray, 42, 50, 80.
Goatsucker, 11.
Golden-eye, American, 17, 34.
Goldfinch, American, 7, 16, 17, 18, 19, 20, 24, 41, 49, 59, 79, 82, 98, 106.
European, 17.
Willow, 36,
Goose, Canada, 20,34.
Lesser Snow, 34.
White-cheeked, 34.
White-fronted, 22, 34.
Goshawk, 83.
Grackle, Bronzed, 7, 16, 24, 41, 49, 59, 79, 107.
Purple, 41.
Grebe, Eared, 33.
Holbœll, 33, 66, 67, 101, 104.
Horned, 102, 104, 112.
Pied-billed, 33, 40, 48, 93.
Western, 33, 101, 104.
Grosbeak, Blue, 112.
Canadian Pine, 17, 83.
Evening, 20.
Rose-breasted, 7, 41, 49, 59, 79.
Western Evening, 36.
Grouse, Oregon Ruffed, 35, 105.
Ruffed, 6, 16, 18, 40, 44, 78.
Guillemot, Pigeon, 33.
Guiraca cærulea, 112.
Gull, Black-backed, 17, 82.
Bonaparte, 21, 34, 48, 93, 102, 104.
California, 102, 105.
Glaucus-winged, 104.
Herring, 17, 18, 19, 33, 40, 48. 78, 82, 93, 104.
Laughing, 78.
Ring-billed, 33' 102, 105.
Short-billed, 33.
Western, 33, 104.

Haliæetus leucocephalus, 6.
Harelda hyemalis, 34.
Hawk, American Rough-legged, 48, 81.
American Sparrow, 6, 17, 18, 19, 20, 39, 40, 48, 78, 81, 92. 98, 107.
Broad-winged, 40, 44, 48, 78.
Cooper, 6, 20, 35, 40, 47, 48, 81, 98.
Desert Sparrow, 35.
Florida Red-shouldered, 17.
Marsh, 39, 40, 48, 78, 107.
Pigeon, 107.
Red-shouldered, 6, 40, 48, 78, 81, 98.
Red-tailed, 6, 16, 17, 18, 19, 20, 40, 48, 78, 98, 107, 108.
Sharp-shinned, 19, 35, 40, 48, 78, 81, 98.
Sparrow, 10.
Western Red-tailed, 35.
Helminthophila alata lutescens, 36.
lawrencei, 84, 85.
leucobronchialis, 109
peregrina, 99.
pinus, 84, 85.
rubricapilla, 8, 99.
Helodramas solitarius, 6.
Heron, Black-crowned Night, 6, 13, 75, 34, 78, 107.
Great Blue, 6, 34, 40, 52, 75, 78, 97, 107.
Green, 6, 34, 40, 48, 78.
Northwest Coast, 105.
Hirundo erythrogaster, 7, 99.
Histrionicus histrionicus, 34.
Hummingbird, Allen, 35.
Ruby-throated, 6, 40, 49, 59, 78, 98.
Rufous, 35.
Hydrochelidon nigra surinamensis, 93.
Hylocichla guttata pallasii, 8, 100.
fuscescens, 8, 99.
mustelina, 99,
swainsonii, 100.
ustulata, 37.
Icterus galbula, 7.
Iridoprocne bicolor, 7.
Ixoreus nævius, 37.
Jaeger, Parasitic, 102, 112.
Jay, Blue, 7, 16, 17, 18, 19, 20, 41, 49, 51, 59, 79, 81, 98, 106.
Gray, 35.
Steller, 35, 105.
Junco hyemalis, 7, 46, 98.
oregonus, 36, 104.
Junco, Oregon, 36, 104.

Slate-colored, 7, 16, 17, 18, 19, 20, 24, 41, 49, 98, 99.
Killdeer, 40, 48, 78, 93.
Kingbird, 6, 22, 40, 49,59, 79, 81, 98, 106.
Kingfisher, Belted, 6, 18, 35, 40, 48, 78, 83, 92, 98, 104.
Kinglet, Golden-crowned, 6, 16, 17, 18, 19, 20, 24, 42, 59, 99.
Ruby-crowned, 8, 24, 42, 50, 59, 103, 105.
Western Golden-crowned, 37, 103, 104.
Lanius borealis, 99.
Lark, Horned, 10, 17,18, 19, 20, 58.
Prairie Horned, 18, 20, 24, 41' 49, 59, 107.
Crested, 9.
Tufted, 9.
Wood, 9, 12.
Larus argentatus, 33, 82, 104.
brachyrhinchus, 33.
californicus, 102, 105.
delawarensis, 33, 102, 105.
glaucescens, 33, 104.
marinus, 82.
occidentalis, 33, 104.
philadelphia, 34, 102, 104.
Loon, 6, 33, 78, 105.
Pacific, 33.
Red-throated, 33.
Lophodytes cucullatus, 82.
Longspur, Smith, 85.
Lophortyx californicus, 35.
Mallard, 34, 40, 48, 74, 78, 82.
Martin, Purple, 7, 41, 49, 51, 59, 76, 79, 107.
Western, 36.
Mavis, 11.
Meadowlark, 17, 19, 20, 24, 41, 49, 51, 58, 59, 79, 98, 107.
Western, 35, 79, 104.
Megascops asio kennicottii, 35.
Melanerpes torquatus, 35.
Melospiza cinerea melodia, 7, 98.
morphna, 36, 104.
georgiana, 7.
Marcea penelope, 112.
Merganser serrator, 34, 105.
Merganser,
American Hooded, 34, 82.
Red-breasted, 34, 48, 105.
Merula migratoria, 8.
propinqua, 37, 104.
Mniotilta varia, 8, 99.
Mockingbird, 59, 80.

Molothrus ater, 7, 98.
Murrelet, Ancient, 33.
Marbled, 104.
Myiarchus crinitus, 6.
Nighthawk, 6, 24, 40, 44, 48, 59, 78, 98, 106, 107.
Western, 35.
Nightingale, 11, 12.
Nettion carolinensis, 34, 83.
Numenius longirostris, 34.
Nuthatch, Red-breasted, 8, 16, 18, 20, 37, 42, 50, 56, 59, 99.
White-breasted, 8, 16, 17, 18, 19, 20, 24, 39, 42, 50, 59, 80, 99.
Nuttalornis borealis, 7.
Nycticorax nycticorax nævius, 6, 15, 34.
Oidemia deglandi, 34. 104.
perspicillata, 34. 104.
Olbiorchilus hiemalis pacificus, 37. 103, 104.
Old-squaw, 34.
Oriole, Baltimore, 7, 24, 41, 49, 79, 107.
Orchard, 39, 41, 49, 59, 79.
Osprey, American, 35, 40, 48, 81.
Oven-bird, 8, 24, 42, 50, 59, 80. 99.
Owl, American Long-eared, 19.
Barn, 107.
California Pygmy, 103, 105.
Barred, 17, 18, 19, 40, 44, 48.
Dusky Horned, 35.
Great Horned, 18, 27, 28, 40, 48.
Horned, 11.
Kennicott Screech, 35.
Screech, 13, 19, 24, 40, 45, 48, 59, 78, 107.
Short-eared, 35.
Snowy, 18, 22.
Pandion haliaetus carolinensis, 35.
Paroquet, Carolina. 112.
Parus atricapillus, 8, 99, 103.
occidentalis, 37, 103, 105.
hudsonius, 99.
Partridge, California, 35.
Passer domesticus, 9, 36, 100, 101, 104.
Passerculus sandwichensis, savanna, 7.
Passerella iliaca unalaschensis, 35.
Perisoreus obscurus griseus, 35.
Petrochelidon lunifrons, 7, 99.
Pewee, Wood, 7, 24, 41, 49, 56, 59, 79, 98, 106.
Phalacrocorax pelagicus, 34.
pencillatus. 34.

Phasianus torquatus, 35, 104.
Pheasant, Ring-necked, 35, 104.
Mongolian, 104.
Phœbe, 6, 27, 41, 49, 79, 98.
Say, 27.
Pigeon, Band-tailed, 35.
Passenger, 111.
Pinicola enucleator leucura, 83.
Pipilo erythrophthalmus, 7.
maculatus oregonus, 36, 103, 104.
Pipit, American, 42, 50, 82, 104.
Piranga erythromelas, 7.
ludovicianus, 36.
Plover, Semi-palmated, 107.
Podilymbus podiceps, 33.
Pooecetes gramineus 7, 98.
Progne subis, 7.
hesperia, 36.
Psaltriparus minimus, 37.
Quiscalus quiscula æneus, 7.
Rail, King, 48, 93.
Virginia, 40, 48.
Red-head, 22. 34.
Redpoll, 16, 36, 59.
Red-wing, Northwest, 35.
Redstart, American, 8, 42, 50, 52, 59. 75, 80, 99.
Regulus calendula, 8, 105.
satrapa, 8, 99.
olivaceus, 37, 103, 104.
Riparia riparia, 7, 36.
Robin, American, 9, 18, 24, 42, 50, 51, 56, 58, 59, 80, 99, 107.
English, 11.
Western, 37, 104.
Salpinctes obsoletus, 36.
Sanderling, 93.
Sandpiper, Baird, 40.
Bartramian, 48, 78, 106.
Least, 40, 48.
Pectoral, 40.
Semipalmated, 21.
Solitary, 6, 40, 48.
Red-backed, 40.
Spotted, 6, 40, 48, 75, 78, 93.
Western, 34.
Sapsucker, Yellow-bellied, 6, 20, 48, 59, 74, 100.
Sayornis phœbe, 6, 98.
saya, 27.
Scolecophagus cyanocephalus, 35.
Selasphorus alleni, 35.

rufus, 85.
Scoter, Surf, 34, 104.
White-winged, 22, 34, 78, 104.
Seiurus aurocapillus, 8, 99.
noveboracensis, 8.
Setophaga ruticilla, 8, 99.
Shoveller, 34, 93.
Shrike, Loggerhead, 20, 58.
Migrant, 41, 49, 79, 107.
Northern, 17, 19, 99.
Sialia sialis, 9, 100.
Mexicana occidentalis, 37, 105.
Siskin, Pine, 86, 105.
Sitta canadensis, 8, 37, 99.
carolinensis, 8, 99.
Skylark, 9, 10, 11, 12.
Snipe, Wilson, 22, 34, 40, 48.
Snowflake, 17, 59.
Somateria spectabilis, 112.
Sora, 40, 48, 108.
Sparrow, Chipping, 7, 41, 49, 58, 59, 79, 98, 99, 105.
English, 9, 24, 36, 59, 100, 101, 104, 109.
Field, 7, 24, 41, 45, 49, 59, 79, 98, 107.
Fox, 59.
Gambel, 36.
Grasshopper, 41, 49, 52, 79, 106.
Golden-crowned, 36.
Henslow, 41.
Lark, 41, 79.
Leconte, 24.
Lincoln, 49.
Nuttall, 36, 105.
Rusty Song, 36, 104.
Sandwich, 105.
Song, 7, 16, 17, 18, 19, 20, 41, 49, 59, 79, 98, 99, 105, 106.
Swamp, 7, 17, 41, 49, 79.
Townsend, 36.
Tree, 16, 17, 18, 19, 20, 41, 59,
Vesper, 7, 19, 41, 49, 52, 59, 79, 98, 106.
Western Savanna, 36.
White-crowned, 41, 49, 59, 98.
White-throated, 7, 17, 18, 20, 24, 41, 49, 59.
Spatula clypeata, 34.
Sphyrapicus varius, 6.
Spinus pinus, 36, 105.
Spiza americana, 52.

Worm-eating, 80.
Yellow, 4, 8, 24, 36, 42, 50, 51, 57, 58, 59, 75, 80, 106.
Water-Thrush, 8, 42, 50, 59.
Louisiana, 42, 50, 80.
Waxwing, Cedar, 7, 20, 36, 41, 49, 79, 99.
Whippoorwill, 6, 23, 40, 45, 48, 59, 78, 98.
Widgeon, 112.
Wilsonia canadensis, 8, 99.
Woodcock, American, 40, 48. 78.
Woodpecker, Downy, 6, 16, 17, 18, 19, 20, 24, 40, 48, 59, 78, 107.
Gairdner, 35, 105.
Hairy, 6, 16, 17, 18, 19, 20, 24, 39, 40, 48, 59, 78, 107.
Harris, 35. 104.
Lewis, 35.
Northern Pileated, 6, 18, 35, 104.
Red-bellied, 18, 19, 20, 24, 40, 48.
Red-headed, 18, 20, 24, 40, 48, 59, 78, 106.
Wren, Bewick, 20, 24, 59.
Carolina, 17, 18, 19, 20, 42, 44.
House, 8, 24, 39, 42, 50, 51, 59, 76, 80.
Long-billed Marsh, 13, 15, 21, 42, 50, 75, 80.
Northwest Bewick, 37, 103, 104.
Pacific House, 37.
Short-billed Marsh, 50, 57.
Rock, 36.
Western Winter, 37, 103, 104.
Tule. 37.
Winter, 16, 17, 20, 42, 50, 59.
Yellowhammer, 9.
Yellow-legs, 40, 48, 107.
Greater, 40, 84.
Yellow-throat, Northern, 8, 42, 50, 57, 59, 80, 99, 108.
Zamelodia ludoviciana, 7,
Zonotrichia albicollis, 7, 98.
coronata, 36.
lencophrys, 98.
gambeli, 36 105.
nuttalli, 36.

INDEX OF AUTHORS.

Abbott, Clinton G., New year bird census, 18, July Fourth Censo-horizon, 77.

Arrick, E. J., New year bird census, 19.

Adams, Reviews of his 'Post-glacial Origin of Migration of the Life of the Northeastern United States,' 'Southeastern United States as a Center of Geographical Distribution of Fauna and Flora,' 29; "The migration Route of Kirtland Warbler."

Antes, Frank T., July Fourth censo-horizon, 77.

Attwater, H. P., Review of his 'Boll Weevils and Birds', 28.

Baird, Robert L., A door-yard list of birds. 23, July Fourth censo-horizon, 77.

Blain, A. W., Jr , New year bird census, 19, July Fourth censo-horizon, 77.

Blake, Francis G., New year bird census, 17.

Blake, Maurice G., New year bird census, 20.

Barnett, Maud, Review of her 'Wisconsin Arbor and Bird Day Anual,' 63.

Burns, Frank L. Brewster Warbler (Helminthrophida leucobronchialis) in Chester Co.. Penn., 109.

Bruen, Frank, New year bird census, 16.

Comey, Arthur C., New year Bird census, 17: A partial list of the summer birds of Holderness, New Hampshire, 5.

Cooke, W. W., Review of his 'Some New Facts about the Migration of Birds," 94.

Craigmile, Esther A., The clock factory, 15, New year bird census, 20. A summer porch list at Hinsdale, Ill., 105.

Crosby, M. S., New year bird census, 17, July Fourth censo-horizon, 77.

Dawson, W. L., Oberlin all day with the birds, 43. An October all-day at Blaine, Wash., 100.

Deane, Ruthven, Review of his 'Unpublished Letters of John James Audubon, and Spencer F. Baird, '64.

Eifrig, Rev. G., Reviews of his 'Birds of Alleghany and Garrett Counties, Western Maryland,' 63, and 'Warbler Songs and Notes,' 6, 64.

Field, Irving A., Review of his 'The Birds of Licking County, Ohio,' 28; July Fourth censo-horizon, 77.

Forebush, Edward Howe, Review of his 'The Destruction of Birds by the Elements in 1903 04,' 94.

Henderson, Judge Junius, Notes from Boulder, Colorado, 27; Bobolinks increasing at Boulder Colorado, 92.

Galpin, A. W., July Fourth censo-horizon, 77.

Henninger, Rev. W. F., A disastrous trip, 21; The larks of Germany, 9; An ornithological reconnissance of the Grand Reservoir, Ohio, in 1904, 74; Note on the nesting of the Black Tern, 92; All day with the birds, 38. Some observations on a captive Red-tailed Hawk, 107. Brewster Warbler again in Ohio, 109.

Heilmann, Carl, All day with the birds, 38.

Hubel, Frederick C., A January Kingfisher, 92; All day with the birds, 39.

Hunt, Chreswell J., The Marsh Wren's midnight song, 14; New year bird census 17; Some Barn Swallow nests, 83; All day with the birds, 39; July Fourth censo-horizon, 77.

Hix, George E., New year bird census, 18; Lawrence Warbler (*Helminthophila lawrencei*) breeding in Bronx Park, New York City, 84; July Fourth censo-horizon, 77.

Jacobs, J. Warren, Review of his 'Gleanings No. III. The Haunts of the Golden-winged Warbler,' 62.

Jones, Lynds, The Cerulean Warbler (*Dendroica rara*), 3; A double nest of Red-eyed Vireo, 15; New year bird census, 19; An improvident Kingbird, 22; All day with the birds, 38; Oberlin all day with the birds, 43; Spring migration in Lorain county, Ohio, 1904, 55; July Fourth, censo-horizon. 77; An addition to the birds of Ohio, 85; Northern Ohio notes, 93.

Judd, Sylvester D., Review of his 'The Economic Value of Bob-white . 94.
Keck, Rev. J. M., Some birds of Olympia, Washington, 33.
Keim, Thos. D., July Fourth censo-horizon, 77.
Levering, Thomas H., New year bird census, 20.
McConnell, Harry B., New year bird census, 18; All day with the birds, 38.
McGregor, Richard C., Review of his 'Birds of Benguet Province, Luzon, and from the Islands of Lubang, Mindoro, Cuyo, and Cagayancillo,' 63.
Metcalf, C. L., New year bird census, 18.
Metcalf, Z. P., All day with the birds, 38.
Morris, C. H., New year bird census, 19.
Moseley, E. L., Notes from Sandusky, Ohio, 112.
Nye, D. E., New year bird census, 19.
Oberholser, Harry C., Reviews of his 'Revision of the American Great Horned owls', 'Reveiw of the Wrenns of the Genus Troglodytes', 27.
Osgood, Wilfred H., Review of his "A Biological reconnaisance of the base of the Alaska peninsular." 113.
Pease, O. H., Notes from Hope, Kansas, 27, July Fourth censo-horizon, 77.
Porter, E. H., Kearsarge Birds, 97.
Porter, Harold E., New year bird census, 16. Kearsarge birds, 97.
Parke, Louis T., New year bird census, 17.
Rogers, Charles H., New year bird census, 16; A two-day census, 39; July Fourth censo-horizon, 77.
Root, Francis M., New year bird census, 19.
Schantz, Orpheus M., A door-yard list from Morton Park, Ill., 57; All day with the birds, 39.
Shufeldt, Dr. R. W., Review of his 'Comparison of the Provisional Schemes of the classification of Birds', 94.
Sillowav, Prof. P. M., The song of the Dickcissel, Spiza americana, 52.
Sim, Robert. J., New year bird census, 18; Notes on the Holboell Grebe (*Colymbus holboellii*), 67; July Fourth censo-horizon, 77; Spring migration along lake Erie's shore, 80.
Smith, Elbert E., New year bird census, 16.
Strong, R. M., Review of his 'The Metallic Colors from the Sides of the Neck of the Domestic Pigeon'; 62.
Swales, Bradshaw H., Additions to list of the winter birds of Wayne County, Micbigan, 82.
Swarth, Harry S., Review of his 'Birds of the Huachuaa Mountains, Arizona' 62.
Taverner, P. A., Review of his "A discussion of the origin of bird migration," 113.
Todd, W. E. Clyde, Review of his 'The Birds of Erie and Presque Isle, Erie County, Pennsylvania, '95.
Wood, Norman A., Review of his 'Discovery of the Breeding Area of Kirtland Warbler', 63.
Walter, Herbert E. and Alice H., Review of their "Wild Birds in City Parks," 61.
Weed, Clarence M., and Dearborn, Ned, Review of their 'Bird Life Stories, '61.
Wilson, Burtis H. New Year bird census, 25; An all day bird study, 51, All day with the birds, 39.
Wolfe, F. B., New Year bird census, 19.

THE WARBLER

JOHN LEWIS CHILDS, Editor

WITH 1905 **The Warbler** begins a new series which will contain many superb Colored Plates of rare eggs such as Kirtland and Olive Warbler, Carolina Parroquet, Clark's Crow, Ipswich and Rufous Crowned Sparrow, Yellow and Black Rail, Calaveras Warbler, etc. Also splendid illustrations of Birds and Nests, and leading articles by well known authorities.

Published Quarterly, 32 Pages & Cover

SUBSCRIPTION PRICE
$1.00 PER YEAR. SINGLE COPIES 30c

Eggs of Kirtland Warbler will be figured in first issue (Jan. or Feb.) of the new series.

ADDRESS THE WARBLER
FLORAL PARK, N. Y.

Bird Papers

For Sale

Complete volumes as well as odd numbers of The Auk, Birds, Birds and Nature, etc., Osprey, and many others.

Write for list and prices.

LYNDS JONES,
Oberlin, O.

DAWSON'S BIRDS OF OHIO

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones 720 quarto pages. 322 birds accurately described, their habits interestingly given Combines ready, identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations to WILSON BULLETIN, Oberlin, O.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES.

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds o Ohio are treated. It does not attempt a description of the plumage. Price is 50 cents. Address, LYNDS JONES, Oberlin, Ohio.

Veterinary Elements.

What the Animal Husbandry Expert, Mr. Geo. M. Rommel, B. S. A., of the U. S. Dept. of Agriculture, has to say regarding "Veterinary Elements."

"I do not feel that I can add anything to that which has been already said in commendation . . . . but from my own personal experience I can say that if I could have had the training in veterinary work that the little book embodies, my course in animal husbandry would have been all the more valuable to me.

"You have pointed out the way for a splendid line of work, and I sincerely hope that the leaders in agricultural education will not be slow in following it."

The author is A. G. Hopkins, B. Agr., D. V. M.; Farmer and Agricultural Journalist; late instructor in Stock Judging and Veterinary Science, Wisconsin Agricultural College.

The above work is clearly and plainly written and easily understood by any one It can be secured, post-paid, for $1.50, from THE WILSON BULLETIN, Oberlin, Ohio.

New and Revised Edition Just Issued.

Principles of Plant Culture

BY E. S. GOFF.

Professor of Horticulture in the University of Wisconsin.

The text-book used in the classes in Plant Life and Horticulture, in the popular Short course in Agriculture, of the University of Wisconsin.

A Few of the Many Things that this Book Teaches.

How to select, test, plant, store and stratify seeds. How to promote flowering and fruiting. How to prevent sun-scald on fruit and other trees. How to promote wood maturity in autumn. How to water and care for greenhouse plants.

The Book Also Teaches

How to maintain the fertility of the soil. How to prevent damage from mice, gophers, rabbits, woodchucks and birds. How to spray plants to prevent insect and fungus ravages. How to prevent damage from chinch bugs, army worms and grasshoppers. How to treat oats to prevent smut. How to make and apply the Bordeaux mixture. How to grow plants from cuttings, layers and stolons. How to propagate plants by grafting and budding. How to make and take care of hot-beds and cold-frames. How to transplant trees, shrubs and herbaceous plants. How to prune fruit and ornamental trees. How to improve plants by selection and cross pollination, and many other things equally valuable.

While the work is written in plain and concise language, the subjects are treated in a thoroughly scientific manner and embody the results of the most recent experiment and research.

The book contains 285 printed pages with 179 illustrations. It is well printed on excellent paper and is neatly and substantially bound in cloth.

This book will prove a most valuable addition to the library of every school of agriculture, horticulture or forestry, and of every rural high or common school in this country, as well as to the library of every farmer and fruit-grower.

Price $1.00 per copy, postage paid, when ordered from

THE WILSON BULLETIN, Oberlin, O.

University
MAY
26
1905

Old Series Vol. XVII. New Series Vol. XII, No. 1.

The Wilson Bulletin

No. 50

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

March, 1905.

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.

CONTENTS.

Some Observations at Weedseed Inn. ALTHEA R. SHERMAN - 1
Breeding Habits of Parula Warbler (*Compsothlypis americana usneæ*) in New Jersey. MARK L. C. WILDE - - 4
Summer Birds of Summit, Union Co., N. J. LA RUE K. HOLMES 8
An Unusual Flight of Hawks in 1858. RUTHVEN DEANE - 13
A Trip to Gardiner's Island. FRANK BRUEN - - 14
Nesting Habits of the Grasshopper Sparrow in South Western Pennsylvania. J. WARREN JACOBS - - - 18
Some Notes on Michigan Warbler. J. CLAIRE WOOD - - 20
Some April and May Work Suggested - - - 22
Editorial - - - - - - - 23
Some Notes on Michigan Warblers. J. CLAIRE WOOD - - 20
Transplanting a Robin. FRANK BRUEN - - - 25
Moving a Wren's Home. FRANK BRUEN - - 26
Bird Builders at Fault. FRANK L. BURNS - - 26
Some New Jersey Records. FRANK L. BURNS - - 27
Book Reviews - - - - - - - 27
Publications Received - - - - - - 29
Roll of Members - - - - - - 30
Election of Members - - - - - - 32

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, O., edited by Lynds Jones.

Subscription: Fifty cents a year, including postagt, *strictly in advance.* Single numbers, 15 cents, unless they are "Special" numbers, when a special price is fixed. The BULLETIN, including all "Special" numbers, is free to all paid up members, either Active, Associate, or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for reviews, should be addressed to Lynds Jones, 5623 Drexel Ave., Chicago, Ill.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 15th of the month preceding publication.

THE WILSON BULLETIN

AN ILLUSTRATED QUARTERLY MAGAZINE DEVOTED TO THE STUDY OF BIRDS IN THE FIELD

Edited by LYNDS JONES

OFFICIAL ORGAN OF THE WILSON ORNITHOLOGICAL CLUB

OLD SERIES, VOLUME XVII. NEW SERIES, VOLUME XII.

1905

Published by the Club at Oberlin, Ohio

Index to Volume 17, 1905.

Acanthis linaria, 50, 118.
Accipiter cooperi, 9, 38, 97.
velox, 9, 38, 97, 120.
Actitis macularia, 9, 37, 51, 84, 86, 97, 120.
Æronympha prosantis, 134.
Actordromas maculatus, 37.
minutilla, 37.
Ægialitis meloda, 38.
semipalmata, 37.
Agelaius phoniceus, 11, 39, 85, 87, 90, 98, 120.
fortis, 90.
Aix sponsa, 8, 97, 120.
Alauda arvensis, 39.
Ammodramus caudacutus, 40.
henslowi, 111.
maritimus, 43.
sandwichensis savanna, 91.
savannarum passerinus, 87.
Ampelis cedrorum, 11, 41, 85, 87, 98, 118.
Anas boschas, 49, 119.
obscura, 37, 49, 97, 119.
Anhinga, 64.
Anhinga anhinga, 64.
Anthus pensilvanicus. 42.
Antrostomus vociferus, 10, 97, 120.
Ardea herodias, 37, 97, 119.
Ardetta exilis, 9, 37.
Asio accipitrinus, 38, 118.
wilsonianus, 10, 117.
Astragalinus, psaltria, 134.
tristis, 11, 40, 85, 87, 98, 118.
Aythya affinis, 49, 89.
americana, 49.
Bæolophus bicolor, 8, 12, 86, 88, 114, 115, 118.
Baldpate, 67, 76.
Bartramia longicauda, 9, 37.
Bittern, American, 8, 77, 120.
Least, 9, 37, 77, 99.
Blackbird, Red-winged, 4, 11, 17, 39, 60, 76, 85, 87, 90, 98, 100, 107, 120.
Rusty, 3, 39, 45, 46, 76, 119.
Thick-billed, Red-winged, 90.
Bluebird, 3, 4, 12, 16, 17, 26, 43, 44, 45, 46, 47, 50, 53, 54, 55, 60, 76, 81, 86, 88, 99, 100, 107, 122.
Bob-white, 3, 9, 17, 28, 52, 53, 54, 60, 80, 100, 107, 117, 133.
Bobolink, 4, 10, 16, 39, 77, 99, 120.
Bonasa umbellus, 9, 49, 97, 117.
Botaurus lentiginosus, 8, 120.
Branta canadensis, 37, 49, 119.
hutchinsii, 64.
Bubo virginianus, 10, 117.
Bufflehead, 49, 77, 80.
Bunting, Indigo, 11, 15, 41, 45, 46, 47, 53, 54, 60, 78, 85, 87, 98, 99, 100, 101, 106, 121, 130.
Buteo borealis, 9, 38, 49, 97, 119.
latissimus, 9.
lineatus, 9, 38, 49, 84, 120.
platypterus, 38, 89, 109, 120.
Butorides virescens, 9, 37, 84, 97, 120.
Cardinal, 11, 28, 40, 45, 46, 47, 52, 53, 54, 60, 81, 82, 85, 87, 100, 113, 118.
Cardinalis cardinalis, 11, 28, 40, 85, 87, 113, 118.
Carduelis carduelis, 40.
Carpodacus purpureus, 40, 110, 118.
Catbird, 3, 4, 12, 17, 21, 42, 45, 46, 53, 54, 59, 60, 78, 85, 87, 98, 99, 100, 101, 116, 121.
Cathartes aura, 28.
septentrionalis, 84, 86.
Ceophlœus pileatus, 108.
Certhia familiaris americana, 43, 118.
Ceryle alcyon, 10, 38, 84, 86, 97, 120.
Chætura pelagica, 10, 35, 85, 86, 97, 120.
Chaffinch, 39.
Charadrius dominicus, 49.
Charitonetta albeola, 49.
Chat, Yellow-breasted, 4, 12, 16, 17, 42, 53, 54, 60, 78, 85, 87, 92, 100, 116, 121, 131.
Chelidon erythrogaster, 11.

Chen cærulescens, 63.
Chewink, 85, 87, 106, 107.
Chebec, 4.
Chickadee, 1, 3, 7, 12, 16, 43, 44, 46, 50, 81, 99, 100, 118.
Carolina, 53, 54, 55, 60, 86, 88, 101, 106, 115.
Chordeiles virginianus, 38, 97, 119.
Chuck-will's-widow, 131.
Circus hudsonius, 9, 38, 120.
Cistothorus palustris, 12.
stellaris, 12, 43, 108, 114, 121.
Clangula clangula americana, 37, 49.
Coccyzus americanus, 10, 38, 116, 120.
erythrophthalmus, 10, 38, 120.
Colaptes auratus, 115.
luteus, 10, 38, 50, 84, 86, 97, 120.
Colinus virginianus, 9, 28, 119.
Colymbus auritus, 36, 49.
holbœllii, 36, 119.
Compsothlypis americana, 119.
usneæ, 4, 42, 85, 87, 108.
Contopus virens, 10, 98, 120.
Coot, American, 76, 79, 132.
Cormorant, Double-crested, 37.
Corvus americanus, 125.
brachyrhinchos, 10, 39, 50, 85, 86, 98, 118.
corax principalis, 108.
ossifragus, 39, 85.
Coturniculus savannarum passerinus, 18, 40, 91, 111, 121.
Cowbird, 4, 11, 17, 39, 61, 76, 82, 100, 120.
Creeper, Brown, 3 43, 44, 46, 54, 76, 80, 82, 118.
Crow, American, 10, 15, 16, 39, 47, 50, 53, 54, 55, 60, 76, 81, 85, 86, 98, 99, 100, 106, 118.
Fish, 39, 85, 100.
Crossbill, American, 28, 50, 67, 110, 118.
White-winged, 82, 111, 118.
Cryptoglaux acadica, 118.
Cuckoo, Black-billed, 4, 17, 38, 54, 78, 100, 120.
Yellow-billed, 21, 38, 47, 53, 79, 99, 100, 116, 120, 131.
Cyanospiza cyanea, 41, 85, 87, 98, 116, 121.
Dafila acuta, 119.
Dendroica æstiva, 12, 85, 87, 98, 121.
blackburniæ, 42, 92, 119.
cærulescens, 119.
castanea, 42.
coronata, 42, 119.
discolor, 92, 116.
dominica albilora, 92.
maculosa, 119.
palmarum, 42.
hypochrisea, 42, 119.
pensylvanica, 8, 9, 12, 42, 98, 121.
striata, 42, 85, 92, 119.
tigrina, 108, 114.
vigorsii, 42.
virens, 42, 119.
Dickcissel, 66, 78, 92.
Dolichonyx oryzivorus, 10, 39, 120.
Dove, Mourning, 4, 9, 53, 54, 60, 67, 76, 81, 84, 99, 100, 120.
Dryobates pubescens medianus, 10, 38, 50, 84, 86, 97, 117.
villosus, 10, 38, 50, 66, 97, 117.
Duck, Black, 18, 37, 49, 76, 80, 81, 97, 119.
Lesser Scaup, 49, 76, 81, 86, 89, 130.
Red-legged Black, 81, 89.
Ruddy, 77, 80.
Scaup, 76, 80.
Wood, 8, 15, 97, 120.
Eagle, Bald, 18, 49, 81, 107.
Golden, 80, 82.
Ectopistes migratorius, 108.
Elanoides forficatus, 90.
Empidonax flaviventris, 39, 80, 108, 110.
minimus, 10, 39, 98, 120.
traillii alnorum, 120.
virescens, 85, 86.
Euphagus carolinus, 39, 119.
Falco columbarius, 38.
peregrinus anatum, 27, 38, 108.
sparverius, 9, 38, 50, 120.
Finch, Purple, 16, 40, 53, 55, 78, 80, 82, 110, 118, 129.
Flicker, 115.
Northern, 2, 3, 10, 17, 38, 44, 46, 52, 53, 54, 55, 60, 76, 80, 84, 86, 97, 99, 100, 101, 106, 120.
Flycatcher, Acadian, 54.
Alder, 120.

Crested, 10, 17, 39, 53, 54, 60, 77, 85, 86, 98, 99, 100, 101, 116, 120, 131.
Green-crested, 21, 60, 78, 85, 86, 100.
Least, 4, 10, 16, 39, 45, 47, 60, 77, 82, 98, 100, 120.
Olive-sided, 4, 39, 119.
Scissor-tailed, 64.
Traill's, 60, 79.
Yellow-bellied, 39, 54, 60, 90, 110.
Fringilla cœlebs, 39.
Fuligula affinis, 86.
Galeoscoptes carolinensis, 12, 42, 85, 87, 98, 116, 121.
Gallinago delicata, 119.
Gallinule, Florida, 27.
Gallinula galeata, 27.
Gannet, 37.
Gavia imber, 36, 49.
Geothlypis agilis, 108, 109.
formosa, 92.
trichas, 85, 87.
brachydactila, 12, 42, 98, 121.
Gnatcatcher, Blue Gray, 53, 60, 67, 78, 86, 88.
Golden-eye, American, 37, 49, 76, 80.
Goldfinch, American, 4, 11, 16, 40, 47, 53, 54, 55, 60, 81, 85, 87, 98, 99, 100, 101, 118.
European, 40.
Goose, Blue, 63.
Canada, 37, 49, 76, 119.
Hutchins', 64.
Goshawk, 14.
Grackle, Bronzed, 39, 44, 46, 47, 52, 53, 54, 76, 99.
Purple, 11, 15, 16, 17, 39, 61, 85, 87, 100, 101, 121.
Grebe, Holbœll's, 28, 36, 119.
Horned, 36, 49, 77, 80, 81.
Pied-billed, 36, 49, 77, 79, 99, 119.
Grosbeak, Blue, 60, 131.
Canadian Pine, 50, 80, 82.
Northern Pine, 50,
Pine, 28.
Rose-breasted, 11, 21, 40, 45, 46, 47, 53, 78, 98, 99, 100, 101, 102, 121, 131.
Grouse, Ruffed, 9, 49, 80, 94, 97, 107, 117, 131.
Gull, Bonaparte, 49, 77, 79, 89.
Great Black-backed, 36.
Herring, 37, 49, 79, 80, 81, 118.
Kittiwake, 36.
Laughing, 74.
Ring-billed, 37, 49.
Habia ludoviciana, 8.
Haliæetus leucocephalus, 49.
Hawk, American Rough-legged, 80.
American Sparrow, 9, 38, 50, 60, 76, 81, 82, 99, 100, 120.
Broad-winged, 9, 38, 53, 61, 76, 89, 100, 109, 120.
Cooper, 9, 38, 76, 82, 97.
Duck, 27, 38, 108.
Fish, 84.
Marsh, 9, 38, 76.
Pigeon, 38, 81, 100.
Red-shouldered, 9, 38, 49, 84, 120.
Red-tailed, 9, 15, 17, 38, 49, 81, 97, 119.
Sharp-shinned, 9, 38, 60, 81, 97, 120.
Helminthrophila celata, 108, 113.
chrysoptera, 41, 89, 98.
lawrencei, 41.
leucobronchialis, 67.
peregrina, 42.
pinus, 11, 41, 121.
rubricapilla, 41.
Hemitheros vermivorus, 11, 41, 87, 92, 98, 121.
Hen, Heath, 107.
Helodromas solitarius, 37.
Heron, Black-crowned Night, 9, 17, 37, 84, 100, 120.
Great Blue, 15, 16, 18, 37, 77, 81, 97, 119.
Green, 9, 17, 37, 61, 77, 84, 97, 100, 107, 120.
Hirundo erythrogaster, 41, 85, 98, 121.
Horizopus virens, 85, 86.
Hummingbird, Ruby-throated, 4, 10, 17, 38, 45, 46, 47, 53, 61, 78, 86, 97, 100, 102, 120, 131.
Hylocichla aliciæ, 43.
fuscescens, 43, 121.
guttata palasii, 43, 119.
mustelina, 43, 86, 88, 99, 116, 121.
swainsonii, 43.
ustulata swainsonii, 119.
Icteria virens, 8, 12, 42, 85, 87, 92, 108, 116, 121.

Icterus galbula, 11, 39, 87, 91, 98, 121.
spurius, 11, 39, 85, 87, 121.
Jaeger, Pomarine, 36.
Jay, Blue, 1, 3, 4, 10, 16, 39, 44, 46, 47, 50, 53, 54, 55, 61, 80, 85, 86, 98,, 99, 100.
Canada, 50.
Junco hyemalis, 40, 50, 118.
Junco, Slate-colored, 1, 40, 44, 47, 50, 52, 54, 55, 80, 118.
Killdeer, 60, 76, 100.
Kingbird, 4, 10, 17, 47, 53, 54, 60, 77, 85, 86, 98, 99, 100, 120, 131.
Kingfisher, Belted, 10, 17, 38, 52, 61, 78, 81, 82, 84, 86, 97, 99, 100, 120.
47, 53, 55, 66, 77, 79, 80, 82, 110, 120.
Ruby-crowned, 43, 44, 45, 46, 50, 53, 66, 77, 79, 119, 131.
Kite, Swallow-tailed, 90.
Lanius borealis, 41, 50, 118.
ludovicianus migrans, 41, 50.
Kinglet, Golden-crowned, 43, 46,
Lanivireo flavifrons, 85, 87.
Lark, Horned, 39, 80, 118, 133.
Hoyt Horned, 80.
Prairie Horned, 2, 53, 54, 55, 81, 91, 130.
Larus argentatus, 37, 49, 118.
atricilla, 74.
delawarensis, 37, 49.
marinus, 36.
philadelphia, 49.
Loon, 36, 49, 77, 80.
Longspur, Lapland, 80.
Lophodytes cucullatus, 49.
Loxia curvirostra minor, 28, 50, 110, 118.
Mallard, 49, 76, 80, 117.
Martin, Purple, 26, 60, 77, 85, 87, 99, 100, 101, 106, 107, 121.
Meadowlark, 3, 4, 11, 17, 39, 47, 53, 54, 55, 58, 60, 81, 85, 99, 100, 120, 131.
Megascops asio, 10, 115, 117.
Melanerpes carolinus, 110.
erythrocephalus, 10, 38, 86, 118.
Meleagris gallopavo fera, 90.
sylvestris, 108.
Melospiza cinerea melodia, 11, 40, 50, 85, 87, 91, 98, 118.
georgiana, 11, 40, 98, 112, 121.
lincolnii, 40, 112.
Merganser americanus, 49, 84.
serrator, 49.
Merganser, American, 49, 76, 80, 81, 84.
Hooded, 49, 77, 80.
Red-breasted, 49, 76, 80.
Merula migratoria, 12, 43, 50, 86, 88, 99, 112.
Mimus polyglottos, 115.
Mniotilta varia, 11, 41, 85, 87, 98.
Mockingbird, 53, 54, 55, 59, 60, 115.
Molothrus ater, 11, 39, 120.
Myiarchus crinitus, 10, 39, 85, 86, 98, 116, 120.
Nighthawk, 17, 54, 60, 78, 97, 100, 119, 130, 131.
Nettion carolinensis, 119.
Nuthatch, Red-breasted, 43, 50, 53, 79, 119.
White-breasted, 4, 12, 43, 44, 50, 60, 81, 88, 99, 101.
Nuttalornis borealis, 39, 119.
Nycticorax nycticorax nævius, 9, 37, 84, 120.
Oidemia americana, 37, 49.
deglandi, 37.
Olbiorchilus hiemalis, 43, 118.
Old-squaw, 80.
Olor buccinator, 28.
columbianus, 28.
Otocoris alpestris, 39, 118.
actia, 133.
praticola, 91.
Oriole, Baltimore, 4, 7, 11, 17, 39, 46, 47, 53, 54, 59, 61, 77, 87, 91, 98, 100, 121.
Orchard, 11, 18, 39, 53, 54, 60, 78, 85, 87, 99, 101, 106, 121, 131.
Osprey, American, 15, 16, 17, 38, 77, 100, 119.
Oven-bird, 12, 16, 17, 21, 42, 45, 46, 53, 54, 60, 78, 85, 87, 98, 100, 101, 102, 106, 121.
Owl, American Barn, 9, 109, 117, 132.
American Long-eared, 81, 117.
Barred, 10, 117.
Great Horned, 10, 81, 106, 117.
Saw-whet, 118.

Screech, 10, 81, 115, 117.
Short-eared, 38, 118.
Snowy, 80.
Pandion haliætus carolinensis, 38, 84, 119.
Passer domesticus, 11, 39, 85, 87, 98, 118.
Passerculus princeps, 40.
sandwichensis savanna, 40, 111, 119.
Passerella iliaca, 40, 112, 119.
Passerina cyanea, 11.
nivalis, 28, 50, 118.
Pavoncella pugnax, 134.
Penthestes aurocapillus, 99.
carolinensis, 86, 88.
Perisoreus canadensis, 50.
Petrochelidon lunifrons, 11, 98, 121.
Peucæa bachmanii, 91, 116.
Pewee, Wood, 10, 17, 53, 54, 60, 78, 85, 86, 98, 99, 100, 106, 120, 130.
Phalacrocorax dilophus, 37.
Pheasant, English, 18.
Philohela minor, 9, 120.
Phœbe, 2, 3, 4, 10, 16, 26, 39, 47, 53, 54, 60, 76, 86, 98, 100, 120.
Picoides arcticus, 50.
Pigeon, Passenger, 13, 48.
Pinicola enucleator leucura, 28, 50.
Pintail, 76, 79, 119.
Pipilo erythrophthalmus, 11, 40, 85, 87, 98, 113, 121.
Pipit, American, 77, 129.
Piranga erythromelas, 11, 41, 87, 98, 121.
rubra, 87, 116.
Plover, American Golden, 49.
Piping, 18, 138.
Semipalmated, 37.
Podilymbus podiceps, 36, 49, 119.
Polioptila cærulea, 86, 88.
Poœcetes gramineus, 40, 119.
Porzana carolina, 37.
Progne subis, 85, 87, 121.
Protonotaria citrea, 113.
Quail, Gambel's, 133.
Querguedula discors, 119.
Quiscalus quiscula, 11, 39, 85, 87, 121.
æneus, 39.
Rail, Clapper, 37.
King, 9, 78, 82, 120.
Virginia, 78, 82, 100, 120.
Rallus crepitans, 37.
elegans, 9, 120.
virginianus, 120.
Raven, Northern, 48.
Red-head, 49.
Redpoll, 1, 50, 80, 118.
Red-wing, Thick-billed, 90.
Redstart, American, 4, 12, 16, 17, 21, 42, 46, 47, 53, 60, 78, 85, 87, 98, 100, 101, 102, 121, 131.
Regulus calendula, 43, 50, 66, 119.
satrapa, 43, 66, 118.
Riparia riparia, 11, 41, 85, 121.
Rissa tridactyla, 36.
Robin, American, 2, 3, 4, 12, 17, 25, 26, 43, 44, 46, 47, 50, 53, 54, 55, 58, 60, 76, 81, 86, 88, 99, 100, 101, 107, 122, 130, 131.
Ruff, 134.
Sandpiper, Bartramian, 9, 37, 77.
Least, 37, 79.
Pectoral, 37, 77.
Solitary, 37, 54, 77, 82.
Spotted, 9, 17, 37, 51, 52, 61, 77, 84, 86, 97, 100, 120.
Sapsucker, Yellow-bellied, 38, 46, 55, 56, 57, 77, 80, 82, 119, 131.
Sayornis phœbe, 10, 39, 86, 98, 120.
Scoter, American, 37, 49.
White-winged, 37.
Seiurus aurocapillus, 12, 42, 85, 87, 98, 121.
motacilla, 42, 85, 98, 102, 119.
noveboracensis, 42, 87, 114, 119.
Setophaga ruticilla, 12, 42, 87, 98, 121.
Shoveller, 27.
Shrike, Loggerhead, 55, 60.
Migrant, 47, 50, 76.
Northern, 1, 50, 80.
Sialia sialis, 12, 43, 86, 88, 99, 122.
Siskin, Pine, 28, 40, 50.
Sitta canadensis, 43, 50, 119.
carolinensis, 12, 43, 50, 88, 99, 118.
Skylark, 39.
Snipe, Wilson, 3, 52, 76, 80, 119.
Snowflake, 28, 50, 80, 118, 129.
Sora, 37, 45, 47, 78.

Sparrow, Bachman's, 53, 54, 58, 60, 91, 116, 131.
Chipping, 4, 11, 17, 46, 47, 53, 54, 60, 76, 77, 98, 100, 101, 115, 121.
English, 3, 4, 11, 16, 17, 39, 85, 87, 98, 118.
Field, 11, 17, 40, 47, 52, 53, 54, 55, 60, 76, 85, 87, 98, 99, 100, 101, 107, 116, 121.
Fox, 40, 76, 80, 112, 114.
Grasshopper, 17, 18, 19, 40, 54, 60, 77, 91, 100, 101, 111, 121.
Henslow's, 111.
Ipswich, 40.
Lark, 46, 47, 77, 91.
Leconte's, 52, 54.
Lincoln's, 40, 53, 54, 78, 112.
Savanna, 17, 40, 77, 91, 111, 119.
Seaside, 35, 43, 100.
Sharp-tailed, 35, 40, 100.
Song, 3, 4, 11, 17, 40, 45, 47, 50, 52, 54, 55, 60, 76, 81, 85, 87, 91, 98, 99, 100, 101, 112, 118.
Swamp, 11, 40, 52, 53, 76, 98, 112, 121.
Tree, 40, 50, 54, 55, 80, 118.
Vesper, 3, 40, 52, 60, 77, 81, 82, 100, 101, 119.
White-crowned, 40, 52, 54, 61, 77, 79, 119.
White-throated, 3, 40, 44, 45, 52, 53, 54, 61, 77, 79, 112, 119.
Spatula clypeata, 27.
Sphyrapicus varius, 38, 56, 119.
Spinus pinus, 28, 40, 50, 119.
Spiza americana, 66, 92.
Spizella monticola, 40, 50, 118.
pusilla, 11, 40, 85, 87, 98, 116, 121.
socialis, 11, 40, 87, 98, 115, 121.
Starling, 39, 120.
Stelgodopteryx serripennis, 11, 41, 85, 87, 113, 121.
Stercorarius pomarinus, 36.
Sterna hirundo, 34, 37, 55, 88.
Strix pratincola, 9, 109.
Sturnella magna, 11, 39, 85, 98, 120.
Sturnus vulgaris, 39, 120.
Sula bassana, 37.
Swallow, Bank, 11, 17, 41, 53, 60, 85, 99, 101, 121.
Barn, 4, 11, 17, 41, 47, 61, 77, 85, 98, 99, 100, 101, 106, 107, 121.
Cliff, 11, 53, 77, 98, 100, 121, 131.
Rough-winged, 11, 41, 53, 60, 77, 95, 98, 101, 113, 121.
Tree, 98, 119.
Swan, Trumpeter, 28.
Whistling, 28.
Swift, Chimney, 10, 17, 38, 46, 47, 53, 54, 60, 77, 86, 97, 99, 100, 101, 107, 120, 131.
Symphemia semipalmata, 27.
Syrnium varium, 10, 117.
Tanager, Scarlet, 11, 17, 21, 41, 48, 53, 59, 61, 78, 98, 100, 101, 102, 106, 121, 131.
Summer, 53, 54, 60, 116, 130.
Teal, Blue-winged, 119.
Green-winged, 119.
Telmatodytes palustris, 43, 86, 121.
thryophilus, 134.
Tern, Black, 27, 78, 81, 126.
Common, 18, 34, 37, 55, 78. 88.
Thrasher, Brown, 2, 3, 4, 12, 17, 43, 44, 45, 46, 47, 53, 54, 59, 60, 77, 87, 98, 99, 100, 101, 107, 115, 121.
Thrush, Gray-cheeked, 21, 43, 78.
Hermit, 3, 43, 54, 59, 61, 77, 80, 81, 82, 119, 129.
Olive-backed, 21, 43, 78, 79, 119.
Wilson's, 12, 43, 46, 47, 53, 54, 60, 78, 121.
Wood, 12, 17, 43, 44, 45, 46, 47, 54, 59, 60, 78, 86, 88, 99, 100, 101, 102, 106, 116, 121, 130.
Titmouse, Tufted, 12, 53, 54, 55, 60, 67, 81, 86, 88, 100, 101, 106, 114, 118.
Totanus melanoleucus, 37, 49.
Towhee, 11, 17, 40, 44, 45, 46, 47, 53, 54, 55, 60, 76, 81, 82, 85, 87, 98, 100, 101, 106, 113, 115, 121, 129.
Toxostoma rufum, 12, 43, 87, 98, 115, 121.
Trochilus culubris, 10, 38, 86, 97, 120.
Troglodytes äedon, 12, 43, 87, 98, 115, 121.

Thryothorus ludovicianus, 8, 12, 43, 85, 87, 121.
Turdus fuscescens, 12.
mustelinus, 12.
Turkey, Wild, 90.
Turnstone, 79.
Tyrannus tyrannus, 10, 39, 86, 99, 120.
Veery, 44.
Vermivora pinus, 85.
Vireo bellii medius, 134.
flavifrons, 11, 41; 116, 121.
gilvus, 41, 91, 121.
noveboracensis, 11, 41, 85, 87, 92, 121.
olivaceus, 11, 41, 98, 121.
philadelphicus, 92.
solitarius, 41, 119.
Vireo, Bell's, 3.
Blue-headed, 21, 41, 77, 79, 119.
Philadelphia. 3, 59, 60, 92.
Red-eyed, 11, 17, 21, 41, 53, 54, 60, 77, 85, 87, 98, 100, 101, 106, 121, 130.
Warbling, 16, 41, 53, 54, 59, 60, 77, 85, 87, 91, 100, 101, 121.
White-eyed, 4, 11, 17, 41, 53, 54, 60, 85, 87, 92, 101, 106, 121, 130.
Yellow-throated, 11, 16, 41, 53, 54, 60, 77, 85, 87, 100, 101, 116, 120, 130, 131.
Vireosylvia gilva, 85, 87.
olivacea, 85, 87.
Vulture, Turkey, 28, 52, 53, 54, 55, 58, 60, 76, 84, 86, 100, 106.
Warbler, Bay-breasted, 21, 42, 60, 67, 79.
Black and White, 11, 17, 21, 41, 44, 45, 46, 53, 54, 60, 77, 85, 87, 98, 101, 121, 130.
Blackburnian, 21, 42, 59, 61, 78, 92, 119.
Black-poll, 21, 42, 53, 54, 60, 79, 85, 92, 119, 131.
Black-throated, Blue, 21, 60, 78, 79, 100, 119.
Black-throated, Green, 21, 42, 53, 54, 60, 78, 79, 82, 119.
Blue-winged, 11, 41, 60, 78, 85, 106, 121.
Brewster's, 67, 78.
Canadian, 46, 47, 60, 79, 98.
Cape May, 53, 60, 79, 114, 131.
Cerulean, 45, 47, 53, 61, 78, 100.
Chestnut-sided, 12, 42, 46, 47, 54, 59, 60, 78, 79, 98, 102, 121, 130, 131.
Connecticut, 21, 60, 79, 109, 130.
Golden-winged, 60, 79, 98.
Hooded, 42, 79, 130.
Kentucky, 78, 92, 100, 130.
Kirtland's, 68, 79.
Lawrence's, 35, 41.
Magnolia, 46, 47, 54, 60, 78, 79, 119, 131.
Mourning, 21, 79.
Myrtle, 3, 21, 22, 42, 53, 54, 55, 60, 78, 79, 119, 130, 131.
Nashville, 21, 41, 60, 78, 79.
Northern Parula, 4, 5, 6, 7, 17, 42, 78, 108.
Orange-crowned, 78, 113.
Palm, 3, 42, 78, 79, 131.
Parula, 59, 60, 85, 87, 106, 119, 131.
Pine, 42, 60, 78, 106.
Prairie, 42, 53, 54, 58, 59, 60, 79, 92, 106, 107, 116.
Prothonotary, 78, 113.
Sycamore, 92, 130.
Tennessee, 42, 60, 79, 130, 131.
Wilson, 42, 45, 47, 60, 79, 114, 119.
Western Parula, 21.
Worm-eating, 11, 41, 53, 59, 60, 87, 92, 98, 121.
Yellow, 12, 17, 54, 60, 78, 85, 97, 98, 99, 100, 101, 121, 130.
Yellow Palm, 42, 53, 79, 119.
Water-Thrush, 21, 42, 78, 79, 114, 119.
Louisiana, 60, 77, 85, 87, 98, 100, 102, 119, 130.
Waxwing, Cedar, 11, 17, 41, 54, 55, 82, 85, 87, 98, 99, 118.
Whippoorwill, 10, 16, 59, 61, 78, 97, 106, 107, 120, 131.
Willet, 27.
Wilsonia canadensis, 42, 98.
mitrata, 42.
pusilla, 42, 108, 119.
Woodcock, American, 9, 15, 27, 48, 76, 81, 120, 129.

Woodpecker, Downy, 10, 16, 38, 46, 50, 53, 54, 55, 57, 60, 81, 84, 86, 97, 100, 106, 117.
Arctic, Three-toed, 50.
Hairy, 1, 10, 38, 44, 46, 50, 54, 61, 66, 81, 97, 99, 100, 117.
Northern Pileated, 50, 81.
Pileated, 53, 54, 80.
Red-bellied, 52, 54, 81, 110.
Red-headed, 4, 10, 38, 53, 54, 60, 77, 81, 86, 99, 100, 118.
Wren, Bewick's, 54, 55, 58, 60.
Carolina, 12, 18, 43, 53, 54, 55, 59, 60, 81, 83, 87, 100, 101, 121.
House, 3, 12, 16, 26, 44, 46, 47, 60, 78, 85, 88, 99, 100, 101, 121.
Long-billed Marsh, 12, 43, 78, 86, 100, 121.
Short-billed Marsh, 12, 43, 114, 121.
Winter, 43, 78, 79.
Yellow-legs, 61, 77, 80.
Greater, 37, 49, 77, 79.
Yellow-throat, Maryland, 53, 60, 85, 87, 106, 107, 130.
Northern, 4, 12, 17, 21, 42, 78, 98, 100, 101, 121.
Zamelodia ludoviciana, 11, 40, 98, 121.
Zenaidura macroura, 9, 84, 120.
Zonotrichia albicollis, 40, 119.
coronata, 40.
leucophrys, 119.

INDEX OF AUTHORS.

Baird, Rob't L.—Bird Migration at Oberlin, Ohio, 75.

Bruen, Frank.—A Trip to Gardiner's Island, 14; Transplanting a Robin, 25; Moving a Wren's Home, 26; Widow Hummer, 102.

Burns, Frank L.—Bird Builders at Fault, 26; Some New Jersey Records, 27.

Deane, Ruthven.—An Unusual Flight of Hawks in 1858, 13.

Embody, G. C.—Bird Horizons from Russellville, Ky., 52.

Hann, H. H.—A Preliminary List of the Birds of Summit County, N. J., 117.

Henninger, W. F.—All Day with the Birds at Durmid, Va., 58; Two Goose Records from Ohio, 63; Further Notes on the Birds of Middle Southern, O., 89; Three Hitherto Unknown Pelican Records from Ohio, 126 ;An Addition to the Birds of Middle Southern Ohio, 126; A Fourth Record of the Barn Owl for Seneca County, Ohio, 129.

Hix, George E.—A Year with the Birds in New York City, 35.

Holmes, LaRue K.—Summer Birds of Summit, Union Co., N. J., 8.

Hunt, Chreswell J.—A-Birding Among the New Jersey Pines, 105; The Spotted Sandpiper, 51.

Jacobs, J. Warren—Nesting Habits of the Grasshopper Sparrow in South-western Pennsylvania, 18.

Jones, Lynds.—Some Further Suggestions for Bird Study, 61; Additions to the Birds of Ohio, 64; The Drumming of the Ruffed Grouse, 94.

Metcalf, Zeno.—Ruby-crowned Kinglet in Winter, 66.

Mills, Wm. J.—Some Breeding Records from East Point, Ga., 115.

Morris, C. H.—Winter Notes on Yellow-bellied Sapsucker (*Sphyrapicus varius*), 56.

Oberholser, Harry C.—Two Bird Days near Washington, D. C., 84.

Rogers, Charles H.—Louisiana Water-Thrush in Philadelphia in Summer, 103.

Schaller, Carleton.—Two Weeks with the Birds in Kent, Conn., 97; Some Spring Records from the Vicinity of New York City, 67.

Sherman, Althea R.—Some Observations at Weedseed Inn, 1.

Smith, Robert Windsor.—Bird Migration at Kirkwood, Ga., 129.

Sparks, Marion E.—Birds vs. Street Cars, 44.

Swales, Bradshaw H.—Additions and Additional Data to a Preliminary List of the Land Birds of South-eastern Michigan, 108.

Taverner, P. A.—Ornithology a Science, 123; Priority, 125; Bird Casualties, 129.

Wilde, Mark L. C.—Breeding Habits of Parula Warbler (*Compsothlypis americana usneae*), in New Jersey, 4.

Wood, J. Claire.—Some Notes on Michigan Warblers, 20; Nesting of the Yellow-bellied Sapsucker, 57; Some Nesting Sites of the Hairy Woodpecker, 66; Extracts from My Note-book, 1905, 129; The Dickcissel in Wayne Co., Mich., 67.

Wood, Walter C.—Autumn Birds of the Les Cheneaux Islands, 48.

ERRATA FOR VOL. XII.

Bulletin No. 50. Contents. For "Some Notes on Michigan Warblers. J. Claire Wood—20," line 12, read "Notes. The Robin's Return. Chreswell J. Hunt."

Page 7, line 10, for "Orioles" read "Oriole's."

Page 22, line 6, for "give" read "gave."

Page 28, line 37, for "specius" read "species," last line, for "speciments" read "specimens."

June number. For No. "52" on front cover read "51."

Page 36, line 13. For the period substitute a comma. Line 35, for "Pomrine" read "Pomarine."

Page 45, line 20, for "Rusty Blackbird" read "Bronzed Grackle."

Page 55, line 3, for "Downey" read "Downy."

Page 62, line 19, for "appears" read "appear."

Bulletin No. 52, the page numbers are thrown forward because in making up the forms, the printer counted in the two pages of advertising beyond blank page 70.

Page 75, line 4, for "presnt" read "present."

Page 75, line 16, for "are from" read "for 1905."

Page 80, line 36, for "Ruffled" read "Ruffed." Line 39 add "northern."

Page 84, line 19, for "Mergauser" read "Merganser."

Page 85, line 1, for "Chactura" read "Chætura." Line 13, for "quiscalus" read "quiscula."

Page 92, for paragraph 22 substitute the following: Icteria virens. Yellow-breasted Chat. On May 22, 1906 (Scioto Co.) I found a set of 2 heavily incubated eggs. On May 17, 1901 (Pike Co.) I found another heavily incubated set. On May 31, 1905, I found a set of 4 fresh eggs in Scioto Co. What is the reason for this peculiarity in breeding habits?

Page 91, line 2 from bottom, for "frequently" read "frequents."

Page 94, line 21, for "intirely" read "entirely."

Page 99, line 8, after "fectly" insert "certain."

Page 102, line 3, for "store" read "stone."

Page 103, line 9 from bottom, for "to" read "too," and for "too" read "to."

Page 106, line 24, for "Priarie" read "Prairie."

Page 109, line 24, for "any" read "my."

Page 112, line 22, after "found" insert "it."

Page 121, bottom line, for Hyocichla" read "Hylocichla."

Page 123, line 6 from bottom, for "objection" read "objective."

Page 132, line 16, for "Hemingway" read "Henninger."

THE
WILSON BULLETIN
NO. 50.
A QUARTERLY JOURNAL OF ORNITHOLOGY.

VOL. XVII. MARCH, 1905. NO. 1

SOME OBSERVATIONS AT WEEDSEED INN.

BY ALTHEA R. SHERMAN.

Not every person who enjoys the companionship of birds is permitted to go a-field every day, or even once each week. The majority of us must be content the greater part of the year with the bird study obtained in our own dooryards. It is the purpose of this article to refer to some of the bird items noted at Weedseed Inn. The latitude of this inn is that of 43° north. Its longitude is nearly that of 91° west from Greenwich, which locates a spot of prairie land six miles from the Mississippi River and one mile and a half from the timber belt that borders that river. This bird hostelry covers a space of three acres, which embraces an orchard, meadow-land, and a portion of a ravine. The latter in spring and autumn is wet enough to entertain some of the water fowls.

Beginning with January the weather for six weeks is usually too severe and the location too exposed for the inn to entertain many birds. Until the last of December Blue Jays have usually called almost every day, and occasionally a Hairy Woodpecker has stopped for a meager lunch, and a mild spell has brought out a few Juncos. But the winter of 1903–4 was an exception. Chickadees came often and pecked at some suet hung out for them. Flocks of Redpolls came several times and bird music was furnished by a Northern Shrike that called often and announced his presence with one of his squeaky airs. Plum trees full of thorns, and a meadow full of mice with but little snow on the ground made Weedseed Inn a favorite caravansary for him. He was often seen impaling a mouse on a

thorn, and the hindquarters of his prey were some times found in the trees after he left.

Toward the last of February the call of the Prairie Horned Lark comes up from the adjoining fields, and it is not long before our Robins and Phœbes return to us. None of our birds have ever been marked, but a neighbor had a male Robin that "the fool with a gun" deprived of a foot. This bird with his mate for two years has nested in her yard. Will another springtime bring him back is a problem soon to be solved.

We feel very certain that some of our birds return year after year. Several Phœbes stay about for four or five weeks, but at nesting-time it is but one pair that remains to occupy the old nest in the barn that has been the birthplace of so many Phœbes. A crack in the flooring above the nest affords a fine opportunity for observing Dame Phœbe's method of raising a family. She does a little refitting of the nest every spring, and on an early day in May lays her first egg. Every morning thereafter between five and nine o'clock an egg is added to the clutch until five jewel-like treasures are to be found. It has been a source of interest to notice the variation in the periods of incubation; that some broods remain longer in the nest than others; that sometimes eight and again ten days elapse from the time the nest is deserted by the first brood before Phœbe lays the first egg of her second set. To watch her five little ones go to bed night after night on the lowest branch of an apple tree has been an interesting experience. Mr. Phœbe must have taken these young ones to another summer resort, since they disappeared from Weedseed Inn about the time Mother Phœbe began her second sitting.

For several years this hostelry has been the home of a pair of Brown Thrashers. They arrive very early in May, and about the middle of that month the first egg is laid. For the past two seasons their first nest for the year has been in a lilac bush about fifteen feet from the house; the same nest being occupied both years. Possibly they might feel that "the world is out of joint" if no human being peeped into their nest each day.

Then there is the Flicker's hole in the barn. It has been there a long time, for it is remembered that he is now a large lad who as a little fellow once asked, "Doctor, where do the

Woodpeckers go when they go in the barn?" The hole gives entrance to a nesting-place four by fifteen inches with a depth of eighteen inches. This space is covered by a removable board in which is a peep-hole commanding a good view of the interior, a rare place for observation. The question of each year is "What bird will take possession of the Flicker's hole?" English Sparrows certainly will if not watched and routed. Sometimes a Flicker raises a brood there, another year a Wren will raise two broods in one summer. The favorably located nests of Phœbe, Flicker, Wren, and Brown Thrasher have each yielded sufficient topics of interest for a story by itself.

Many of the common birds nest about the place, but their nests are not always found. One year a Bob-white had a nest containing fourteen eggs not more than ten feet from the plowed ground of the garden in which people worked daily. Unfortunately a mowing machine ran over the spot and destroyed the nest before mankind discovered it.

During migration days Weedseed Inn entertains its share of guests. Early in the spring one may look for a day now and then when a pair of Hermit Thrushes spends the whole day here, moving about until the gathering gloom of night hides them from view, but they are never seen the following day. On other days the maple trees are covered with Rusty Blackbirds, enough to fill four and twenty pies, and the air is stirred with music, enough to fill a whole country side. For about a week in both spring and fall a dozen or more of Wilson's Snipe prod and paddle in the marshy spots of the ravine, and the question arises, "Are they not the same birds that tarry each season?" One spring day comes to mind when the trees fairly swarmed with warblers. I then had little experience in naming the birds and could identify but few of them.

The hour in which I identified the greatest number of birds was from 7:30 to 8:30 o'clock on September 24, 1904. Almost all of them were seen from one window. The birds observed there were the Flicker, Phœbe, Blue Jay, English Sparrow, White-throated Sparrow, Song Sparrow, Meadowlark, Philadelphia Vireo, Bell's Vireo, Myrtle Warbler, Palm Warbler, Brown Thrasher, House Wren, Chickadee, Bluebird and Robin. A few minutes after half past eight Vesper Sparrows and a Brown Creeper were seen, but a Catbird that nested in

the yard during the summer did not appear until later in the day. Add to this list of nineteen the birds seen during the six previous days of the week which were Black-billed Cuckoo, Olive-sided Flycatcher, Baltimore Oriole, Goldfinch, American Redstart and White-breasted Nuthatch. This lot for one week was equalled in number on May 25, 1904, when twenty-six species of birds were observed at Weedseed Inn. They were Mourning Dove, Flicker, Red-headed Woodpecker, Ruby-throated Hummingbird, Kingbird, Phœbe, Chebec, Blue Jay, Bobolink, Cowbird, Red-winged Blackbird, Meadowlark, Baltimore Oriole, English Sparrow, Chipping Sparrow, Song Sparrow, Swallow, White-eyed Vireo, Maryland Yellow-throat, Yellow-breasted Chat, Brown Thrasher, House Wren, Catbird, Bluebird, Robin, and another which could not be satisfactorily named.

Early rising and a day devoted to observing the birds would, no doubt, secure a much longer list of bird guests for one day at this bird hostelry. A list of seventy-nine species named and many others that were not identified suggests in a limited degree what has been seen in one yard by a tyro in the dooryard study of birds.

BREEDING HABITS OF PARULA WARBLER (*Compsothlypis americana usneæ*) IN NEW JERSEY.

BY MARK L. C. WILDE.

Parula Warblers are very common during the breeding season, in suitable localities, throughout the lower half of the state of New Jersey. Commencing at Brown's Mills, on the Rancocas Creek, situated in Burlington county some fourteen miles east of Mount Holly, and journeying southward to the Delaware Bay and the Atlantic Ocean, these birds can be found breeding on the edge of all swamps, streams, lakes, ponds, and mill dams, where there is a fairly good growth of that bearded lichen (Usnea barbata), which many of the south Jerseymen deign to call "Beard-Moss."

While the climatic conditions, to a very large extent, may be responsible for the presence and growth of this so-called "Beard-Moss," one thing is certain, and that is, this lichen absolutely controls the distribution of the Parula Warbler, as

far as the state of New Jersey is concerned. Wherever there is an abundant growth of Usnea barbata a colony of these beautiful warblers will be found breeding, for, strange as it may seem, they construct their nests in it, build their nests of it, and they search in and around it for insects, on all the trees, bushes, and vines, where it grows.

Before proceeding further I wish to say a few things relative to this lichen (Usnea barbata). I shall hereafter call it "Beard-Moss," only because that name has been associated in my mind for so many years, in connection with Parula Warblers. It therefore comes to me more naturally than any other name possibly could. "Beard-Moss" must be seen to be appreciated.

It is a long, stringy, or hairlike-growth, grayish-green in color, and when covered with dew or moistened by the rain, it feels quite soft and is pleasant to the touch. In the dampened condition it gives you the impression of elasticity; so much so that I have often tried to make it stretch, as it should were it composed of rubber. It grows in a tangled mass the lower strands only being combed out, as it were, and left to float in the passing breeze. It grows on the tree limbs, and is thick in proportion to the thickness of the limb from which it hangs, excepting when it grows on vines or bushes where the twigs are close together; then the whole growth is a solid mass of "Beard-Moss." In New Jersey it thrives best on the lower limbs of the trees and on small bushes and vines near the surface water of the swamps, streams, or mill ponds, and in the lakes or ponds it mostly grows at the "head," or upper ends, where the water, coming down, empties into these bodies of water.

Imagine being at the "head" of one of these ponds, where all the trees and bushes are draped and festooned with this beautiful growth, and here, on the warm summer days, you are amid the Parula Warblers on their ideal breeding grounds.

The growth of the "Beard-Moss" at Brown's Mills is very scant as compared with the growth along the Maurice River up to about Willow Grove pond, located in eastern Salem county, and even here it is not nearly so abundant as in lower Cumberland, Atlantic, and Cape May counties. The farther

south you go the thicker will be the growth of the "Beard-Moss," hence the Parulas are more abundant.

Parula Warblers breed in colonies, owing to the "Beard-Moss" occurring in separate and distinct patches, where conditions are most favorable to its growth. Some of the ponds where I once found many of their nests, are now almost deserted, simply because the mill-dams have broken, or the water has been drained off, causing the "Beard-Moss" to dry up and die, and the Parulas have taken up quarters in some other neighboring locality where Usnea barbata thrives more luxuriantly.

Parula Warblers arrive from the South in the lower portion of the state about the first of May, apparently already paired, and within about ten days or two weeks nest building has begun. Full sets of fresh eggs are usualy deposited in Cape May county by May 20th, while further north, at Brown's Mills, Burlington county, the date would be about June 1st.

Almost all of the many nests I have examined contained four eggs. I have never found over four, but occasionally some nests contain but three. It will therefore be seen that four eggs constitute a full set under normal conditions.

On several occasions I have spent from seven to ten days at a time right among these Warblers on their breeding ground, from early morning until dusk, and by not distrubing their nests, they have become so accustomed to my presence as to take little notice of me. I have moved along slowly and quietly in my boat, passing in and around the bushes and trees containing their nests, often within three or four feet of the sitting birds. The feeding birds were likewise just as unconcerned as those which were incubating their eggs.

Nests can be found from the border to the middle of the mill-ponds and open swamps, and may be looked for anywhere from under the tip of an outstretched or drooping branch, to against the tree trunk, or in the smaller bushes; and from one foot above the water to twenty feet high. Generally, however, on account of the "Beard-Moss" growing more abundantly on the lower branches of the trees and on the bushes, five feet may be considered the average height.

The females alone attended to the construction of the nest, while the males were leisurely feeding in and around the tan-

gled moss-covered branches, often clinging to them upside down in Chickadee fashion, reaching here and there for lurking insects, and flying a short distance, they would pause for a moment to emit their sweet song.

The nest is invariably placed in a hanging position. The female usually selects a tree or bush in which the "Beard-Moss" grows quite thickly, and here, within the tufts, she loops and weaves together the inside particles of moss, forming a beautiful nest, much resembling the style of the Baltimore Orioles, only of course on a very much smaller scale. The bird is careful that the moss shall be left hanging in its natural way from the bottom and sides of the nest, and often so conceals it that it can only be found by close and careful searching. Into the structure the bird then carries thread-like particles of the moss collected from some near-by tree. I have never known them to use moss from the tree or bush in which their nest is built for lining purposes.

"Beard-Moss" is used exclusively by some Parulas in lining their nests, while others add a few horse-hairs and a yellow down which is taken from the stems of swamp ferns. The nest is very compact and closely woven, occasionally having a few pine-needles stuck into it around the outside, probably to help support and pin it to the hanging particles of moss.

The entrance, which is always on a level with the top of the bowl, is made through the moss on the side, very often directly under the limb where the moss is parted. The walls of the bowl, being at least half an inch in thickness, form a platform which is sometimes flattened out, resembling a small mat, on which the bird rests when entering or leaving the nest. Some nests have two or more entrances, either left as peep windows for escape, or unintentionally caused by the thinness of the moss above the bowl. I have examined a few nests where the entrance was made from the top, the nest having been suspended either between two twigs, or between the trunk of a tree and an adjoining tuft; but such cases as these are rare, and may be considered departures from their regular style of building. From a very large number of nests of this species examined during the past ten years, nearly all were entirely or partly roofed over, with the entrances from the sides, as previously described. One nest collected some years

ago, was suspended from a two-inch limb, containing little or no moss outside of that of which the nest was constructed; but this I do not consider a typical nest.

The inside measurements of the nests vary, ranging from about one and a half to two inches both in depth and diameter.

The eggs show a remarkable variation both in size and shape. They have a white ground color, and are more heavily dotted with reddish brown and lilac at the larger ends, often forming a ring around them.

Camden, N. J.

SUMMER BIRDS OF SUMMIT (UNION COUNTY), NEW JERSEY, AND VICINITY.

BY LA RUE K. HOLMES.

Summit is located on the crest of the Second Mountain, at an elevation varying from 300 to 520 feet above sea level. The surrounding hills were at one time well wooded, but most of the timber now standing is of recent growth, only a few small tracts of virgin forest now remain and these are being rapidly cut. The cover is for the most part deciduous, there being but few groves of coniferous trees in this neighborhood that I know of. The ground is furrowed in every direction by streams, mostly of a small size and a river (the Passaic), runs through the center of the territory covered by this list. Several fresh-water swamps of considerable area are within walking distance of Summit and are frequently visited.

As will be seen from the list, Summit is located between the Carolinian and Alleghanian zones; and such Carolinian birds as Bæolophus bicolor, Thryothorus ludovicianus, Icteria virens; and Alleghanian birds as Dendroica pensylvanica, Habia ludoviciana, etc., are found breeding here.

1. **Aix sponsa.** Wood Duck.—A rare summer resident, formerly more common. A pair succeeded in rearing a brood of 8, this last summer (1904) within three miles of Summit. The parent birds with the young were seen daily by many people and were apparently devoid of fear.

2. **Botaurus lentiginosus.** American Bittern.—A rare summer resident in the large fresh-water marshes. I only know of three nests having been found.

3. **Ardetta exilis.** Least Bittern. A summer resident, probably rare. I have no nesting record.

4. **Butorides virescens.** Green Heron. A rather common summer resident in all suitable localities. Never more than one pair breeding in a woods (?).

5 **Nycticorax n. nævius.** Black-crowned Night Heron. A large colony of these birds was formerly located in this vicinity; but after having been attacked many times by plume hunters and eggers, was finally broken up. I have no recent record of its having bred in this section.

6. **Rallus elegans.** King Rail. A nest containing nine eggs of this species was found in the Great Swamp several years ago. It may breed there regularly.

7. **Philohela minor.** Woodcock. A rather rare summer resident. Breeds early in the spring. Nests have been found while snow was yet on the ground.

8. **Bartramia longicauda.** Bartramian Sandpiper. A very rare summer resident. I know of but one field inhabited by this bird and as a rule two pairs of birds breed there each season.

9. **Actitis macularia.** Spotted Sandpiper. A rather rare summer resident; breeding chiefly in low-lying corn fields; at least that is where nests have most often been found.

10. **Colinus virginianus.** Bob-white. Formerly common. Now a rare summer resident, but seems to be increasing in the last year or two.

11. **Bonasa umbellus.** Ruffed Grouse. The same may be said of this, as of the preceding species.

12. **Zenaidura macroura.** Mourning Dove. Rare. Some years more common than others.

13. **Circus hudsonius.** Marsh Hawk. A few pairs breed in the larger fresh-water marshes.

14. **Accipiter velox.** Sharp-shinned Hawk. A very rare summer resident.

15. **Accipiter cooperi.** Cooper's Hawk. The nest of this species is met with more often than that of *A. velox*, and judging from individuals seen I should say it was much more common.

16. **Buteo borealis.** Red-tailed Hawk. A very rare summer resident and becoming rarer with the cutting of the virgin forests on the higher hills.

17. **Buteo lineatus.** Red-shouldered Hawk. Our commonest breeding hawk.

18. **Buteo latissimus.** Broad-winged Hawk. A very rare summer resident.

19. **Falco sparverius.** Am. Sparrow Hawk. A not uncommon summer resident; next to *B. lineatus* in point of numbers.

20. **Strix pratincola.** Am. Barn Owl. An extremely rare summer resident. I know of but one nest having been found and have

a record of five young birds not fully feathered having been caught in a trap set for hawks and owls in a pasture.

21. **Asio wilsonianus.** Am. Long-eared Owl. A rare summer resident.

22. **Syrnium varium.** Barred Owl. Formerly a rather common resident but now rapidly disappearing.

23. **Megascops asio.** Screech Owl. A not uncommon resident; rapidly decreasing in numbers.

24. **Bubo virginianus.** Great Horned Owl. Never common, but now almost entirely exterminated locally. I doubt if a pair breeds nearer than ten miles of Summit. It has, however, been captured, during the early winter (December and January) in steel traps set on posts in meadows near Summit within the last year or two, and it may breed in the wilder portion of the county.

25. **Coccyzus americanus.** Yellow-billed Cuckoo. A rather common summer resident; more abundant some years than others.

26. **Coccyzus erythrophthalmus.** Black-billed Cuckoo. Not as common as the preceeding, and equally erratic.

27. **Ceryle alcyon.** Belted Kingfisher. A rather common summer resident; nesting in all suitable localities.

28. **Dryobates villosus.** Hairy Woodpecker. A rather rare summer resident.

29. **Dryobates pubescens medianus.** Downy Woodpecker. A common summer resident.

30. **Melanerpes erythrocephalus.** Red-headed Woodpecker. Locally common.

31. **Colaptes auratus luteus.** Northern Flicker. The commonest breeding woodpecker.

32. **Antrotosmus vociferus.** Whip-poor-will. Formerly a common summer resident, but I have no record of any having been seen in the summer for the past fifteen years.

33. **Chætura pelagica.** Chimney Swift. An abundant summer resident.

34. **Trochilus colubris.** Ruby-throated Hummingbird. A common summer resident.

35. **Tyrannus tyrannus.** Kingbird. A common summer resident.

36. **Myiarchus crinitus.** Crested Flycatcher. Rather common in woodland.

37. **Sayornis phœbe.** Phœbe. An abundant summer resident.

38. **Contopus virens.** Wood Pewee. A common summer resident in woodland.

39. **Empidonax minimus.** Least Flycatcher. A rather rare summer resident.

40. **Cyanocitta cristata.** Blue Jay. Abundant.

41. **Corvus brachyrhynchos.** American Crow. Abundant.

42. **Dolichonyx oryzivorus.** Bobolink. Locally common. Apparently decreasing in numbers.

43. **Molothrus ater.** Cowbird. Common.

44. **Agelaius phœniceus.**. Red-winged Blackbird. Abundant in the fresh-water marshes.

45. **Sturnella magna.** Meadowlark. Common.

46. **Icterius spurius.** Orchard Oriole. Locally rare.

47. **Icterus galbula.** Baltimore Oriole. Common; breeding even in the heart of the city.

48. **Quiscalus quiscula.** Purple Grackle. Common. Breeds in colonies in conifers.

49. **Passer domesticus.** English Sparrow. Abundant everywhere.

50. **Astragalinus tristis.** American Goldfinch. Common.

51. **Spizella socialis.** Chipping Sparrow. Abundant in the settled portions; rare in rural districts.

52. **Spizella pusilla.** Field Sparrow. Abundant summer resident.

53. **Melospiza cinerea melodia.** Song Sparrow. Abundant summer resident.

54. **Melospiza georgiana.** Swamp Sparrow. Abundant summer resident.

55. **Pipilo erythrophthalmus.** Towhee. Common summer resident.

56. **Cardinalis cardinalis.** Cardinal. A rare summer resident; only two or three records.

57. **Zamelodia ludoviciana.** Rose-breasted Grosbeak. A common summer resident.

58. **Passerina cyanea.** Indigo Bunting. A rather common summer resident.

59. **Piranga erythromelas.** Scarlet Tanager. A common summer resident.

60. **Petrochelidon lunifrons.** Cliff Swallow. Formerly common; has now entirely disappeared.

61. **Chelidon erythrogaster.** Barn Swallow An abundant summer resident.

62. **Riparia ripara.** Bank Swallow. Never very common, but now entirely extirpated.

63. **Stelgidopteryx serripennis.** Rough-winged Swallow. A very rare summer resident.

64. **Ampelis cedrorum.** Cedar Waxwing. A common summer resident.

65. **Vireo olivaceus.** Red-eyed Vireo. Abundant summer resident.

66. **Vireo flavifrons.** Yellow-throated Vireo. Rather rare.

67. **Vireo noveboracensis.** White-eyed Vireo. Common along streams.

68. **Mniotilta varia.** Black-and-white Warbler. Rather rare.

69. **Helmitherus vermivorus.** Worm-eating Warbler. An extremely rare summer resident.

70. **Helminthophila pinus.** Blue-winged Warbler. A common summer resident.

71. **Dendroica æstiva.** Yellow Warbler. Common summer resident.

72. **Dendroica pensylvanica.** Chestnut-sided Warbler. Rare summer resident; but apparently increasing as a summer resident.

73. **Seiurus aurocapillus.** Oven-bird. Commonest breeding warbler.

74. **Geothlypis trichas brachidactyla.** Northern Yellow-throat. Common summer resident.

75. **Icteria virens.** Yellow-breasted Chat. Rather common summer resident.

76. **Setophaga ruticilla.** American Redstart. A very rare summer resident; but one record.

77. **Galeoscoptes carolinensis.** Catbird. Abundant summer resident.

78. **Toxostoma rufum.** Brown Thrasher. Common summer resident.

79. **Thryothorus ludovicianus.** Carolina Wren. A very rare summer resident; but two records.

80. **Troglodytes aëdon.** House Wren. Rather common.

81. **Cistothorus stellaris.** Short-billed Marsh Wren. Locally common; breeds in a very restricted area.

82. **Cistothorus palustris.** Long-billed Marsh Wren. Locally common in fresh-water marshes and along Passaic river.

83. **Sitta carolinensis.** White-breasted Nuthatch. Common.

84. **Bæolophus bicolor.** Tufted Titmouse. Rare; becoming more common.

85. **Parus atricapillus.** Chickadee. Common.

86. **Turdus mustelinus.** Wood Thrush. Abundant.

87. **Turdus fuscescens.** Wilson's Thrush. Rather common.

88. **Merula migratoria.** Robin. Abundant.

89. **Sialia sialis.** Bluebird. Abundant.

The foregoing list I believe to be fairly complete; there are however, several birds whose status, as far as this locality is concerned, I am in doubt of, and have thought it best to eliminate them from the list.

It will of course be understood that the relative abundance of birds mentioned in this list is applicable to the breeding season only.

AN UNUSUAL FLIGHT OF HAWKS IN 1858.

BY RUTHVEN DEANE.

The following letter is not only of interest in itself, but the fact that it is addressed to Audubon's two sons, John Woodhouse and Victor Gifford, adds to its historical value. While it is well known that there are conspicuously large migrations of hawks every spring and fall, when birds are almost continually in sight and at times lasting several consecutive days, yet such a one as here described quite reminds us of the enormous flights of the Passenger Pigeon in the early days, as related by Audubon and Wilson.

As to the thirteen specimens shot being those of the Goshawk, would seem most unlikely, as all authorities for the state quote the species, not only as a winter visitor, but one of quite rare occurrence.

I am much indebted to Miss M. R. Audubon, who has recently presented me with this letter:

"CINCTI, Sept. 27th, 1858.

"My Dear Sirs,

I wish to communicate an important fact in natural history, never known to me before, and which I would hardly have believed, had it not been witnessed by an old friend, who is a good observed of nature—Mr. G. C. Coney [1]—in whose veracity I can place the most implicit confidence.

"On the 17th inst., at 8 o'clock A. M., he went out with his gun, when he happened to see, what he considered a rather large flock of hawks, but upon casting his eyes around, he saw as far as he could see a continuous stream of them coming.

"This flock was about 40 yards in width, and took by his watch one and a half hours to pass.

"They came from N. N. E. and flew S. S. W. They sailed in a direct line with extended wings, but at certain distances some of them kept wheeling around and around in a circle, and were joined by others, while some of them floated off again with the main stream.

"He shot two of them which were a little lower than the rest. As it was such an unusual sight, when he saw them he immediately ran and shouted for some of the neighbors, who all witnessed the facts.

[1] George C. Coney, St. Marys, Ohio. An enthusiastic and well known sportsman, as well as a keen observer of birds. Died several years ago.

"The place where this occurred was on a great reservoir in Mercer County, Ohio, midway between St. Mary's and Celina. This body of water feeds the Miami Canal, is ten miles long, and from three to five miles in width. During the summer season there are always some of these hawks—a smaller species—and also a few of a much larger kind—are always about there. Owing to the facility of procuring their food, as there are a great many young ducks, squirrels, etc., they also do great damage to the farmers' chickens. One friend had upwards of one hundred young chickens and only forty-five are now remaining.

"He shot thirteen of the species in controversy, which according to his description must have been the Goshawk (Astur palumbarius) and also shot many of the other two kinds during this summer, but never saw more than from two to ten (the highest number) in one flock together before.

"Such migration of hawks I believe was never heard of before, and the question is will they do so annually? and in day or night time. If the first, I should think they would have been seen before.

"I was on the prairies in August, grouse were scarce, owing to the first brood being nearly all destroyed by the heavy rains in May and June,, when these level prairies were all under water. The young ducks, however, were very plenty. When will Agassiz be out with his fish?

Truly yours,

J. S. UNZICKER.[1]

"Messrs. J. W. and V. G. Audubon, N. Y."

[1] Dr. J. S. Unzicker, Cincinnati, Ohio. Born August 11, 1812, died April 18, 1876. A prominent physician, an ardent sportsman, and lover of birds, and had been personally acquainted with Audubon and his sons.

A TRIP TO GARDINER'S ISLAND.

that my wish was realized.

BY FRANK BRUEN.

It would be presumptuous for me to try to write up the birds of Gardiner's Island after the admirable article contributed by Mr. Frank M. Chapman to the Nov.-Dec. (1903) number of Bird-Lore; but it would be nearly impossible to visit Gardiner's Island and write nothing at all. Ever since reading the above-mentioned article, I have had a consuming desire to visit the island; but it was not until June 17, 1904,

The approach, of Mr. Ford and myself, to the island was not in the romantic fashion described by Mr. Chapman, but in an *intermittent* gasoline boat from Saybrook Point, Conn.

Beaching our boat we found what proved to be the best camping spot on the island; it was also richest in the number of bird species. Although three o'clock in the afternoon when we landed, nevertheless, thirty-four (34) species of birds were noted in the immediate vicinity.

Our stay from June 17th to 21st was all too short to properly cover the island in detail, but still we felt that we had seen a great deal. Most of the birds mentioned by Mr. Chapman were seen; all, I believe, excepting the Wood Duck, Woodcock, and Indigo Bunting; and in addition to his list were the Great Blue Heron and Red-tailed Hawk. The latter was seen three different days—an Ishmaelite among birds, for every feathered creature was against him. When first seen our attention was attracted by his "exhaust steam" scream as he went off, chased by Purple Grackles. The next day at another part of the island some Grackles and Ospreys were after him, and the third day some Crows were after him. He kept in the virgin forests as much as possible. It was a question in our minds whether he was trying to live there or wished to leave the island but could not owing to the vigilance of the Ospreys.

The Osprey display was simply amazing to us who had never seen them at such close quarters nor in such abundance. Most of the nests examined had young birds from one to ten days old as nearly as we could judge; a few had eggs only or eggs and just hatched young. On the beach at one place were four nests, placed on the ground, in a distance of 700 feet; one nest had so *little material* in it that it compared closely, except in size, with the *most elaborate*(?) of the terns' nests. We noted 78 nests of the Osprey in use, but undoubtedly there were as many more on the island. At one time 38 Ospreys were in sight, counting those easily seen on their nests and those in the air.

In the two tern colonies 35 nests containing 89 eggs were found: without making an exhaustive search. The south colony is really two, being divided by an inlet in which the water was too deep to wade at the time of our visit, or more nests would have been recorded. One nest had four eggs, but most

of them had three eggs; others of one or two eggs each were probably not yet complete.

We found the English Sparrows and Purple Grackles nesting in the crannies of the Ospreys' nests. The sight of the Britishers in the depths of the virgin forests was a most unwelcome one to us.

To us the Grackles, which were abundant, formed the one bad feature of the island bird life; for they seemed to be continually upon the watch to prey upon the young and eggs of the smaller birds. Many times as we passed through the woodland paths (one cannot go outside of them owing to the dense growth of green or cat briers), Ovenbirds, Redstarts ,etc., would scold us for coming near their nests or young; and almost immediately the Grackles would be seen slipping up to see what they could find.

A storm the last day of our stay on the island prevented our looking for the Great Blue Herons' nests.

A novelty to us was the constant chatter of the Yellow-breasted Chat at night. Almost the first bird to greet us after we landed, he kept it up all night excepting an interval between 8 and 10 o'clock; whenever Mr. Ford or I awoke during the night he was going full blast.

We were disappointed in the length of our list for the island, for we could have found a larger number of species with the same effort, at home; but the special features of the list more than made up for its shortness. We missed our Yellow-throated and Warbling Vireos, Purple Finches, Whip-poor-wills, Blue Jays, Bobolinks, Phœbes, and House Wrens, and some others,—to be sure some of them may have been there but missed by us. But one pair of Bluebirds was seen. No Least Flycatcher was seen or heard, but he seems to vanish or lose his voice at about this time even in Connecticut

We feel under many obligations to Mr. Gardiner, the owner of the island, for his courtesy in allowing us to camp there.

Subjoined is a list of the birds noted:—

1. Downy Woodpecker, 4.
2. White-breasted Nuthatch; few seen.
3. Chickadee; few seen.
4. American Goldfinch; few seen.
5. American Crow; abundant.

6. Song Sparrow; common.
7. Red-tailed Hawk, 1.
8. Bobwhite; common.
9. Herring Gull; 12 or 13 seen.
10. Meadowlark; 6 seen.
11. Bluebird, 2.
12. Robin; common about the homestead only.
13. Red-winged Blackbird; common.
14. Purple Grackle; abundant.
15. Cedar Waxwing, 1.
16. Northern Flicker, 4.
17. Field Sparrow; few seen. Sheep graze too close for cover,
18. Belted Kingfisher. rather common.
19. Cowbird, 3.
20. Savanna Sparrow; 10 seen.
21. Chipping Sparrow; 2 seen.
22. Barn Swallow; few near barns.
23. Chimney Swift, 3.
24. Towhee; fairly common.
25. Black and White Warbler; 3 seen.
26. Brown Thrasher; several.
27. Spotted Sandpiper; fairly common. Saw crow carry off a young one.
28. Bank Swallow; several large colonies.
29. Catbird; fairly common.
30. Northern Yellow-throat; common.
31. Oven-bird; common.
32. Yellow Warbler; common.
33. Wood Thrush; fairly common.
34. Red-eyed Vireo; common.
35. Baltimore Oriole; 2 seen.
36. Kingbird; few seen.
37. American Redstart; fairly common.
38. Yellow-breasted Chat; 2 seen.
39. Black-billed Cuckoo; few seen.
40. Scarlet Tanager, 1.
41. Grasshopper Sparrow, 9.
42. Crested Flycatcher; fairly common.
43. Ruby-throated Hummingbird, 1.
44. White-eyed Vireo, 2.
45. Northern Parula Warbler; 7 noted where usnea moss grew.
46. Wood Pewee; common.
47. Nighthawk, 4.
48. Green Heron; fairly common.
49. Black-crowned Night Heron; common.
50. English Sparrow; common.
51. Osprey; abundant.

52. **Orchard Oriole, 1.**
53. **Common Tern, 200.**
54. **Black Duck, 2.**
55. **Carolina Wren, 3.**
56. **Great Blue Heron, 5.**
57. **Bald Eagle, 1.**
58. **English Pheasant, 20.**
59. **Piping Plover, 5.**

NESTING OF THE GRASSHOPPER SPARROW IN SOUTHWESTERN PENNSYLVANIA.

BY J. WARREN JACOBS.

The Yellow-winged Sparrow *(Conturniculus savannarum passerinus)* is distributed in limited numbers, during the nesting season, throughout Greene county, in the extreme southwestern corner of Pennsylvania. Their favorite resorts are in the hillside fields and along the low ridges. In pasture fields, not too cleanly kept, and where the wild "sink field" mats its frail vines through the carpet of blue grass, the birds choose a site for a nest. Not every apparently good field has its pair of birds, and indeed one may pass through several such fields without hearing the song of this bird or flushing the female from her nest. However, it must not be inferred from this that birds have not escaped notice, or that a nest has not been passed without the sitting bird taking flight.

The bird itself is very shy; its song pleasing, but not distinguishable a very great distance, being easily drowned by the rattle of numerous ever-singing chats and the medley of a dozen other species which haunt the neglected fields. The female is a close sitter, not leaving her post until almost trampled upon. This makes nest-seeking very tedious; and the apparent scarcity of this species renders uncertain the result of a careful search.

One of the very first nests new to me was of this species, and stumbled upon accidentally on top of the ridge overlooking my home town, away back in the '80s, in about the second year of my bird studies. The old bird fluttered from under my feet and darted down over the hill to a brier clump. It was a new bird to me then, but I got a fairly good look at it; and a year

or two later, when the little first Edition of "Davies' Nest and Eggs" came out, I cleared up the identity, which, by the way, was made easier by the fact that no other sparrow, having similar nests and eggs, was likely to be found here. The eggs, I discovered, after getting home, were badly incubated, and I lost them at the end of the blow-pipe.

Several years went by without this bird or its nest coming under my observation; and when, in 1893, I was preparing a collection of Pennsylvania eggs for exhibition at the Chicago World's Fair, I almost grieved over the loss of this set, but the species was represented by a set of three eggs from New Jersey. A bird apparently so rare, I thought, could be found nesting only by chance, and after several fruitless hunts, I gave up hope of ever replacing the lost set. Thus the matter stood for some years, when a friend brought me a nest and five eggs of a sparrow, unknown to him, which he had found by accidentally stepping against a bunch of grass, flushing the bird. These proved to be eggs of the Yellow-winged. The eggs were fresh, and were found on July 8, 1898, in a pasture field about six miles west of Waynesburg. The composition of the nest was of grass and grass-rootlets, lined with fine grass. Measurements: Outside diameter, 4.5 in.; inside, 2.5 in. Outside depth, 2.0 in.; inside, 1.3 in. It was sunken in the ground and well concealed by the tuft of grass.

Comparing these eggs with two sets taken lately by myself, I find them to be much smaller, and exhibiting a less mottled appearance where the markings are thickest, which is in the form of a broken wreath around the larger end of each. The ground color is pure white, which is the case with the other two sets mentioned, and the predominating color of the markings is burnt sienna, intermixed with fewer blotches of lavender and heliotrope purple. In shape they are broad or rounded ovate, and measure .73x.56, .72x.56, .71x.56, .67x.54, and .69x .56 inch.

The finding of this set of Yellow-winged Sparrow revived my old enthusiasm to find the bird nesting. Therefore, during the next few years, I spent many hours in fruitless search, until June 21, 1903, when, as on my first discovery, I accidentally flushed a bird from her nest sunken in the ground and well concealed by "sink field" vines, in a neglected field

on the side of a hill. I was returning home in the evening, after a day spent in the woods, and in crossing this field I stepped entirely over the nest before the old bird fluttered out and took refuge in a brier clump. The eggs, four in number, were fresh, and are rather heavily marked, the blotches forming a solid wreath around the larger ends of three and a mottled cap on the fourth; the colors being vinaceous-cinnamon, vinaceous, lavender, heliotrope purple, and burnt sienna—the first predominating. Elliptical-ovate in shape, and measure .78x.58, .79x.58, .79x.60, and .78x.57 inch.

The last nest found by me was on July 21, 1903, and like others, was stumbled upon quite accidentally. I was out berry-picking with my nephew and some of his young friends, when one of the party called my attention to a bird he had just flushed at his feet. Seeing it was a Yellow-winged Sparrow, I began a search for its nest, and soon found it quite near to where the boy was standing. It was composed wholly of grass and was placed in a depression five inches deep. It contained four eggs in which embryos were beginning to form. The same description of coloring given for the last will answer for this set, except that the mottled cap is lacking. They are also slightly more pointed than the last. Size: .81x.61, .81x.62, .80x.60, and .78x.61 inch.

Taking the dates into consideration, this bird probably raises but one brood in this part of its range, and its nesting period extends throughout June and July, with care of young sometimes reaching into August.

SOME NOTES ON MICHIGAN WARBLERS.

BY J. CLAIRE WOOD.

There is a certain piece of thick woods here covering about twenty acres of ground. Its exact location is Private Claim 49, Ecore township, Wayne county. The more dense portions are free of undergrowth, but in places the forest floor is concealed by the thickest kind of brush tangle. Elm, red oak, maple, beech, butternut, chestnut and sycamore abound in the order named. A luxuriant growth of wild grape vines is a characteristic feature of the butternut section and near the

easterly end is a second growth of paw paw trees while beyond the easterly margin lie low swampy bush lands with openings grown to marsh grass. The woods is long and narrow and divided its entire length by a county drain six feet deep and twenty broad at the top. As no other timber of a similar nature exists in the neighborhood this is a favorite resting place and resort for the woodland migrants.

For the purpose of ascertaining the approximate time of departure and relative abundance of the late warblers I devoted what days I could spare to this woods and worked it so thoroughly that it is doubtful if any species escaped notice. October 23 was the last day I searched this woods, but the warblers were gone except the Myrtle and one Northern Yellow-throat. As, in the question of identification, the greater importance is attached to specimens secured a * indicates that one or more were taken on the date to which it is prefixed. The numbers exceeding 25 were estimated, but all less than that were counted.

1904	Sept. 25	Sept. 28	Oct. 2	Oct. 6	Oct. 16
Black and White Warbler	*1		*1		
Nashville Warbler		*1			
Western Parula Warbler		*12			
Black-throated Blue Warbler	*6		*15	*3	
Myrtle Warbler		*6	*24	*6	*125
Bay-breasted Warbler	*7	*6	*3		3
Black-poll Warbler	*3	*125	*3		*3
Blackburnian Warbler	*2				
Black-throated Green Warbler	*3	*1	*75		
Connecticut Warbler				*1	
Mourning Warbler	1			1	
Northern Yellow-throat			*3		
American Redstart	*2	6	*2		
Oven-bird	15	7			
Water-Thrush	1				

Two Rose-breasted Grosbeaks* were noted on the 25th. A few Olive-backed and about 50 Grey-cheeked Thrushes were observed on the 28th. Green-crested Flycatchers* were last seen on the 2d; also Catbirds, Red-eyed Vireos* and Scarlet Tanagers.* The Blue-headed Vireos were common inclusive of the 2d, but only one on the 6th. Last Yellow-billed Cuckoo on latter date.

About three o'clock one foggy morning early in September, 1887, large numbers of the Myrtle Warblers were congregated about the electric light tower at the corner of Woodward and

Adams avenues of this city. There were also smaller numbers at the next tower but, although they appeared to fly against the glass, none were injured. At the other tower, however, the birds kept falling at the rate of about one per minute. I returned at eight o'clock and started to count the dead but soon give it up. It is sufficient to say that there were five or six bushels of them. They were all Myrtles with the exception of one Field Sparrow.

SOME APRIL AND MAY WORK SUGGESTED.

During the season of migration we are so occupied with the movements of the birds northward that we generally forget that there are other sides to bird study. When do the different species of birds begin to build their nests? How long does it take a pair to complete a nest? Are the eggs deposited on successive days? How long is the period of incubation? Do both birds take part in the incubation, or does one sit continuously while the other feeds her, or how is it? How rapidly do the young grow, and when do they leave the nest? When do the feathers first appear, and how are the downs attached to them? Do the birds use the old nest a second time, and if so do they remodel it, clean it, or use it as the young left it? How do the old birds feed the young during the first few days after hatching? These, and a host of other questions can be answered by anybody who can and is willing to give some time to watching nests that may be so placed that they can be seen at close range. Have an eye to your immediate surroundings in addition to specially favored places where you love to go. Prove that the slur often aimed at amateur field work is not applicable in your case at least. Such work needs to be done.

THE WILSON BULLETIN.

A Quarterly Magazine Devoted to the Study of Living Birds.
Official Organ of the Wilson Ornithological Club.

Edited by LYNDS JONES.

PUBLISHED BY THE CLUB, AT OBERLIN, OHIO.

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number, postpaid.
Price in all countriss in the International Postal Union, 65 cents a year, 20 cents a number.
Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr, Frank L. Burns, Berwyn, Penn., or to Mr. John W. Daniel, Jr., Washington, D. C.

EDITORIAL.

The editor has Mr. Frank L. Burns to thank for collecting a considerable part of the copy for this issue.

The next number will contain two local lists of more than usual interest to readers in general. Carefully prepared local lists are always in demand.

The editor's headquarters will remain at 5623 Drexel avenue, Chicago, Ill., until further notice in the June number. Correspondence relating to advertising, subscriptions, and articles intended for print should be addressed to him there, but letters of request for sample copies or back numbers will receive prompt attention if addressed to Oberlin, Ohio.

It is not too early to be thinking about the May migration work. In spite of the splendid record made last year there is plenty of room for a better record yet. Study the weather map, if you have opportunity, and with its help select the best day and then work your field to your fullest capacity. Put your locality and your work in the first rank for wealth of bird life during the migrations. The editor cannot promise to print all "All Day" lists, but he will gladly print a list of those who make such lists with the number of species recorded if each person will send his list for inspection and possible tabulation.

In *Bird Lore* for November-December, Mr. Ernest Tompson Seton in introducing an excellent outline for the study of birds, writes in substance that the time has gone by when adequate good can result from ordinary collecting in well-known regions and that the experts of our museums, knowing the value of birds as they do, better than any other class of men, are the only ones who should be allowed to collect bird skins to-day. Just what is meant by ordinary collecting is not apparent, yet the student seeking to faithfully carry out his suggestions will find it expedient if not absolutely necessary to good work in fully one-quarter of the thirty-one sub-

heads offered, to take life occasionally. Were the museums many times as numerous as they are at present, there would still be many individuals unable to reach them through lack of time, distance or other causes. Would it be right to exclude such persons from collecting? In what manner have a certain coterie attained superior knowledge of the value of kinds over that of another class also studying at first hand? All honor to the museum expert! He is usually a hard and conscientious worker, who, not for value received, but perhaps for a consideration, collects, preserves and studies birds for the benefit of the public. Likewise the humble amateur—without the consideration. No one has, or should have, a monopoly of the knowledge of the value of a single species. With all respect for the writer, such sentiments are pure rot, unworthy of him and unjust to those at which they are aimed. F. L. B.

In *American Ornithology* for January, Mr. Reed has given us much excellent matter relating to the Warblers of the genus *Geothlypis* together with a plate showing the members in color, one-half natural size. F. L. B.

Prof. H. A. Surface, Economic Zoologist of Pennsylvania, has made a strong point of bird protection since he entered the office, in order that the people could understand why the birds should be preserved from an economic standpoint. The result has been most gratifying. Never before has there been such a strong sentiment for the protection of birds in the State. F. L. B.

Mr. Frank Bruen has an interesting article on the winter birds of Bristol, Conn., in a recent issue of a local paper. F. L. B.

Mr. W. H. Brownson, of the Portland *Adveriser*, is now the editor of the *Journal of the Maine Ornithological Society*, and publishes in the former for January 14, "A List of the Birds Observed in Cumberland County (Maine) in 1904," including 146 species; also on January 21, "A visitor from the North Pole," an account of the capture of a Dovekie, blown in by a storm. F. L. B.

The editor is sorry that space did not permit the printing of the New Year Censo-Horizons this time, because other matter of more importance was received. He is glad to be able to report, however, that there was more material sent in than could be printed in this number. If this were always true the editor would not only be saved a considerable amount of worry, but a better Bulletin would always be insured. Original field work which has for its object the increase of our knowledge of the birds is always in great demand, and earnestly solicited. We have scarcely more than made a beginning in the study of bird habits.

NOTES.

THE ROBIN'S RETURN—Much has been said of late concerning the intelligence of birds. It is evident that birds are more or less intelligent. Whether they act entirely by instinct or whether they have some mental faculties developed to a more or less degree we cannot say with certainty, but some of their actions show an amount of intelligence.

In the spring of 1903 a young lady, living in a suburb of Philadelphia, Pa., found a young Robin which had evidently fallen from a nest and was starving to death. She took it into the house and fed it and as it grew she became very much attached to it and it became very tame. After it was full grown it was allowed entire freedom. Every day it would go out and forage for itself but would always return at night to the house. A window was left open a few inches for it and every evening it would come into the house to roost, leaving again early in the morning.

In the late fall when the Robins were migrating it too disappeared, presumably following its kin to their winter residence.

This spring (1904) the lady was attracted by a Robin chirping noisily in a tree near the house. She hardly thought it possible that her Robin had returned but she went into the yard and spoke to the bird in the tree. The bird at first seemed a little shy but would let her approach quite close to it. At last it seemed to get more confidence and finally flew to her outstretched hand and alighted upon it.

This summer it has again been roosting within the house every night and spending the day abroad.

This not only proves that the bird returned to its haunts of the preceding summer but it proves also that it was able to recognize a human face. It apparently knew its friend when it saw her after an absence of several months.—Chreswell J. Hunt.

TRANSPLANTING A ROBIN.—Happening to be in the foundry flask yard July 4, 1904, I noticed a large cope leaning against a pile of flasks, the baffle boards of which made a series of shelves upon which were several robin's nests; the arrangement being as in the subjoined sketch. Nests Nos. 1, 2 and 3 were complete; Nos. 4 and 5 being merely foundations. Nos. 1 and 2 contained one egg each partly incubated I supposed. Visiting the nests July 6, two eggs were found in nest No. 1. I called the attention of the yard foreman to the nests and asked him if he would try to protect them from harm. He exclaimed, "Well, that is hard luck, for we need that cope today." After talking it over a little, we



concluded to take another cope as nearly like the first as could be had, putting it in the same position as the first and moving the nests to the same relative positions and await results.

This plan was carried out excepting the arangement of the nests; for on July 7, when I again visited the nests, they were placed as shown below—the foundations Nos. 4 and 5 not having been moved.



However, when I approached, Mrs. Robin was seen sitting on nest No. 1. On July 8, she was on nest No. 2, but on the 9th, 11th, 12th, 13th and 16th she was on No. 1. On July 20 two young birds were found and the nest was deserted on the 29th. Nest No. 3 was completed but did not seem to have been used.

Mrs. Robin will no doubt remember her hard trials but eventual triumph over difficulties, a long time and steer clear of the flask yard; but I felt abundantly repaid for the experiment. Another time I should put all the eggs in one nest.—FRANK BRUEN, Bristol, Conn., December, 1904.

MOVING A WREN'S HOME.—For the past two seasons a pair of House Wrens have built their nest in a crevice formed at the joint where brace and post meet on the veranda of the Bristol Golf Club Tea House. From May 11 to 27, this year (1904) the building was dismantled and moved to a new site about three-fourths of a mile distant. When the veranda was taken down the wren's nest came out and fell to the ground. The men employed noticed the birds at that time but do not remember whether they were about afterwards or not.

July 18, a pair of Wrens, presumably the same, were still using the crevice at the new location.

My chain of evidence I know is weak in the above, and I write this more to draw out the experience of other members of the club than for its scientific value.—FRANK BRUEN, Bristol, Conn., December, 1904.

BIRD BUILDERS AT FAULT.—Apropos to Mr. Bruen's "Transplanting a Robin." John Burroughs in *Bird Lore*, page 85, 1901, under the title of "A Bewildered Phœbe," gives an instance of a builder at fault. The bird confronted by new conditions, blunders, but through its great industry is not altogether baffled by the multiplicity of building sites. I have frequently observed the same trouble when the Robin or Pewee selected a place midway on a beam or plate partitioned by rafters into short stretches exactly alike, particularly where there was no resting perch in full view of the nesting site. Naturally, when a corner of the building is chosen in the beginning, there is no difficulty, for it may be easily distinguished. Less commonly the Bluebird, House Wren and Purple Martin err when building in a many-roomed box, also doubtless

the smaller Woodpeckers are sometimes bewildered when found digging indifferently in one of two or more cavities in the same stub. In fact the antiquated notion that a bird knows exactly what it is about at all times and places is no longer tenable.—FRANK L. BURNS, Berwyn, Penna.

SOME NEW JERSEY RECORDS.—I am indebted to Dr. W. R. Wharton the past season for some substantial additions to my collection. The folowing kinds, taken in Salem county, New Jersey, near the Delaware river, on the dates given, and handled in the flesh by the writer, seem worthy of special mention.

Shoveler or Spoon-billed Duck (*Spatula clypeata*). A female, September 23. One or two are said to have been met with every fall. Considered quite rare along the coast.

Florida Gallinule (*Gallinula galeata*). Two immature males, September 2. Local sportsmen shoot an occasional specimen in mistake for one of the larger Rails.

Willet (*Symphemia semipalmata*). A single male, October 8. This was in all probability a transient; although the species was at one time a summer resident.

Duck Hawk (*Falco peregrinus anatum*). An immature female, October 21. Another specimen was secured at the same place some time ago.—FRANK L. BURNS, Berwyn, Penna.

BOOK REVIEWS.

Taylor's Standard American Egg Catalogue, second edition, An Exchanger's Guide and Collector's Handbook. By Henry Reed Taylor. Price 25 cents. Alameda, California.

In eleven pages of preliminaries, some racy, some serious, the author and F. M. Dille discuss several matters of interest, among which the one on "Egg Valuations" will prove of general interest because of the previous varying values, the determining factor seeming to be the abundance of the species in the market, and the possessor's ability to secure prices in exchange. Purchase prices have always been far below exchange prices. It appears from this article that the prices herein quoted are the result of a sort of general agreement among those especially interested. The list proper is, of course, merely a check-list with prices attached where any can be determined. Thus something over 250 of the species and sub-species listed are not accompanied by prices. A careful perusal of the list sets one wondering if .100 for Black Tern and 5.00 for American Woodcock may not really stand for .10 and .50. Other evidences of rather careless work on the part of the printers leaves a degree of disappointment. But one cannot expect a list of this sort, especially at the price quoted, to be perfect. It is beyond question of great value to those who deal in eggs in any way. It suggests the possibility of a revival of the days now two decades

past when Oology was in the forefront. In our opinion Oology as such and such alone will never be able to stagger to its feet again. As a part of Ornithology in the broad sense it has its legitimate and necessary place, but as a distinct science it has and ever will prove a failure, for the very simple reason that it does not possess within itself the elements of a separate science. L. J.

The Birds of the Rockies. By Leander S. Keyser.

In this large octavo there is a sympathy of the author with his mountain surroundings and of the artist, Louis Agassiz Fuertes, with the conception of the book, that one rarely finds. No three-color-process colored pictures here, but finely executed colored engravings, full-page half-tone and sketchy marginal engravings. A good, clear and clean type makes easy reading of an entertainingly written account of a bird-lover's experiences in the Colorado mountains. If the author possesses any fault of execution it lies in giving himself too fully to the enamoring influences of the mountain environment. The price of the book, reduced to $1.50, will make it appeal to every lover of outdoors. L. J.

Annual Report of the Division of Zoology, p. 159–180, also Reports of the Ornithologist for 1902 and 1903, p. 227–232, and 611–616, by Prof. H. A. Surface, in the Pennsylvania Report of the Department of Agriculture, 1903.

I will quote the following as worthy of special mention: "During the summer the Turkey Vultures (*Cathartes aura*) were unusual in their northern flight. They have been seen as far north as Williamsport, and a few at State College, Center county. * * * During the fall a Cardinal (*Cardinalis cardinalis*) was seen by us in Center county. * * * I have evidence that the American Crossbill (*Loxia curvirostra minor*) nests in the central part of Pennsylvania, in the fact that during the middle of the summer of 1902 a pair of old birds accompanied by four of their young were seen for two days upon the State College campus, in Center county, feeding upon the mites of the cockscomb elm gall. During the winter, which was unusually long and severe, there was a decided immigration of northern birds into our State. Especially remarkable among these were the Pine Grosbeak (*Pinicola leucura*), the Crossbills (both specius of the genus *Loxia*), the Snowflake (*Passerina nivalis*), and the Pine Siskin (*Spinus pinus*). The severity of the winter resulted in killing most of the Quail (*Colinus virginianus*) of our State, many of the Wild Turkeys and some of the Ruffed Grouse. * * * * During the early portion of this spring (1904) there has been an unusual flight of the aquatic birds. In the vicinity of Harrisburg, gunners have shot several species of Wild Ducks, the Holbœll's Grebe, several specimens of the Whistling Swan (*Olor columbianus*), and one specimen of the Trumpeter Swan (*Olor buccinator*). We have been fortunate in securing speciments of these

very rare birds in this State, and hope to receive funds for having them permanently preserved in a State museum. There was also an unusual flight of Gulls along the Susquehanna river after the breaking up of the ice, and this was doubtless due to the masses of ice filling the bays and covering their feeding grounds near the mouth of the river." Modest quarterly and monthly bulletins with a mailing list of upward of twenty-four thousand names, have taken the place of the costly and highly colored picture books of a few years ago, and earnest efforts are being made to advance the knowledge of economic ornithology. For the first time in a number of years, continuous work is being done in the central part of the State, and while it might be said in criticism that more specific information would in no wise detract from the popular nature of the records of the occurrence of certain of the rarer species secured, an immense amount of work is being done. F. L. B.

PUBLICATIONS RECEIVED.

Amateur Sportsman, Vol. XXXII, No. 3.

American Ornithology, Vol. V, Nos. 1, 2.

Bird Lore, Vol. VII, No. 1.

Boys and Girls, Vol. III, No. 4; Vol. IV, Nos. 1, 2.

Bulletin of the Michigan Ornithological Club, Vol. V. No. 4.

Bulletin No. 18, U. S. Department of Agriculture, Biological Survey. Distribution and Migration of North American Warblers. By Wells W. Cooke.

Bulletin No. 19, U. S. Department of Agriculture, Biological survey. Hunting Licenses, Their History, Objects and Limitations. By T. S. Palmer.

Bulletin 222, Michigan State Agricultural College Experiment Station.

Bulletin 150, 156. Ohio Agriculture Experiment Station.

Bulletin No. 69. Pennsylvania State College Agriculture Experiment Station.

Bulletins of Pennsylvania Department of Agriculture, Vol. II, Nos. 2, 8, 9.

Cassinia, Vol. III.

Colorado College Studies, Science Series. Vol. XI, Nos. 30, 31, 32, 36, 37, 38.

Condor, The, Vol. VII, No. 1.

Journal of the Maine Ornithological Society, The, Vol. VI. No. 4.

Maine Sportman, Vol. XII, Nos. 136, 137.

Naturalistie Canadien, Le, Vol. XXXI, No. 12; Vol. XXXII. No. 1.

Ohio Naturalist, The, Vol. V, Nos. 2, 3.

Ornithologische Monatschrift, Vol. XXIX, Nos. 11, 12; Vol. XXX, No. 1.

Report of the Michigan State Board of Agriculture, 1904.

Summer Resident Birds of Brewster County, Texas. By Thos. H. Montgomery, Jr. From The Auk, Vol. XXII, No. 1.

ROLL OF MEMBERS.

Officers for 1905.

President—Lynds Jones, Oberlin, Ohio.
Vice-President—W. Leon Dawson, Seattle, Wash.
Secretary— John W. Daniel, Jr., Washington, D. C.
Treasurer—Frank L. Burns, Berwyn, Penn.
Executive Council—Harry C. Oberhalser, Washington, D. C.; John H. Sage, Portland, Conn.; Alexander W. Blain, Detroit, Mich.

Active Members.

Clinton G. Abbott, 153 West 73rd St., New York City.
Robert L. Baird, 148 Woodland Ave., Oberlin, Ohio.
Paul Bartsch, Smithsonian Institution, Washington, D. C.
A. C. Bent, Taunton, Mass.
Homer L. Bigelow, 511 Washington St., Boston, Mass.
Dr. Louis B. Bishop, 356 Orange St., New Haven, Conn.
Alexander W. Blain, Jr., 131 Elmwood Ave., Detroit, Mich.
Charles F. Brennan, Mt. Carmel, Ill.
William Brewster, 145 Brattle St., Cambridge, Mass.
Clarence Morrison Brooks, 105 West St., Keene, N. H.
W. H. Brownson, Advertiser office, Portland, Maine.
Frank Bruen, 34 Spring St., Bristol, Conn.
Henry R. Buck, City Hall, Hartford, Conn.
G. M. Burdick, Milton Junction, Wis.
Frank L. Burns, Berwyn, Penn.
Mrs. Agnes Chase, 59 Florida Ave., N. W., Washington, D. C.
Miss Esther Craigmile, Box 179, Hinsdale, Ill.
Maunsell S. Crosby, Rhinebeck, N. Y.
Frank S. Daggett, 341 Rialto Building, Chicago, Ill.
Edward M. Daniel, Lynchburg, Va.
John W. Daniel, Jr., 1340 Columbia Road, Washington, D. C.
Mrs. Elizabeth B. Davenport, 45 Green St., Brattleboro, Vt.
Rev. W. Leon Dawson, University Station, Seattle, Wash.
Walter Deane, 29 Brewster St., Cambridge, Mass.
Owen Durfee, Box 125, Fall River, Mass.
William Dutcher, 525 Manhattan Ave., New York City.
Dr. J. Dwight, Jr., 2 East 34th St., New York City.
George C. Embody, Bethel College, Russellville, Ky.
Louis J. Eppinger, 299 Gratiot Ave., Detroit, Mich.
Miss E. W. Fisher, 1502 Pine St., Philadelphia, Penn.
George L. Fordyce, 40 Lincoln Ave., Youngstown, Ohio.
Louis Agassiz Fuertes, 13 East Ave., Ithaca, N. Y.
Benj. T. Gault, Glen Ellyn, Ill.
N. C. Gilbert, Bethel College, Russellville, Ky.
Mrs. Stephen Cobb Goss, 5475 Ridgewood Court, Hyde Park, Ill.
Hon. Junius Henderson, Boulder, Boulder Co., Colo.
Carl Fitz Henning, Boone, Iowa.

Rev. W. F. Henninger, 206 Jefferson St., Tiffin, Ohio.
George E. Hix, 114 W. 90th St., New York City.
Ned Hollister, Delavan, Wis.
La Rue K. Holmes, Pine Grove Ave., Summit, N. J.
Frederick C. Hubel, 112 Alexandrine Ave., Detroit, Mich.
Chreswell J. Hunt, 1306 N. 53rd St., Philadelphia, Penn.
J. Warren Jacobs, Waynesburg, Penn.
Lynds Jones, 160 North Professor St., Oberlin, Ohio.
J. W. Kay, 62 Sheldon Ave., Detroit, Mich.
W. G. Kay, 62 Sheldon Ave., Detroit, Mich.
J. Eugene Law, Hollywood, Calif.
W. P. Lyons, 41 East Fourth St., Cincinnati, Ohio.
Miss Rebecca Moorhead Leete, North East, Penn.
Herbert H. Marble, Box 71, Fall River, Mass.
Harry B. McConnell, Cadiz, Ohio.
Frederick B. McKechnie, Ponkapog, Mass.
Mrs. Lida B. McMurry, Garden City, Kas.
Mrs. Olive Thorne Miller, 827 De Kalb Ave., Brooklyn, N. Y.
Dr. Walter I. Mitchell, Metropolitan Hospital, Blackwell's Island, New York City.
C. H. Morris, McConnelsville, Ohio.
Harry C. Oberholser, 1454 Sheridan Ave., Washington, D. C
O. H. Pease, Hope, Kansas.
Clark J. Peck, 6728 Leeds St., Philadelphia, Penn.
Harold E. Porter, 101 West 85th St., New York City.
Charles H. Rogers, 211 South Aberdeen Ave., Wayne, Penn.
Dr. Willet E. Rotzell, Narberth, Penn.
John H. Sage, Portland, Conn.
William E. Saunders, 352 Clarence St., London, Ont.
Orpheus M. Schantz, Morton Park, Ill.
George H. Selleck, 110 High St., Exeter, N. H.
Dr. A. R. Shearer, Mont Belvieu, Texas.
Prof. P. M. Silloway, Lewiston, Mont.
H. H. Skinner, 286 Commonwealth Ave., Cleveland, Ohio.
Clarence J. Stone, Branchport, N. Y.
Prof. H. A. Surface, Economic Zoologist, Harrisburg, Penn.
Bradshaw H. Swales, 46 Moffat Building, Detroit, Mich.
P. A. Taverner, 577 Warren Ave., Detroit, Mich.
John E. Thayer, Box 98, Lancaster, Mass.
Miss Adelaide Utter, Clerk U. S. Circuit Court, Kansas City, Mo.
Warner M. VanNorden, 751 Fifth Ave., New York City.
Walter F. Webb, First National Nurseries, Rochester, N. Y.
Mark L. C. Wilde, 35 N. 5th St., Camden, N. J.
Burtis H. Wilson, 1102 Seventeenth St., Rock Island, Ill.
Sidney S. Wilson, Attorney-at-Law, St. Joseph, Mo.

Honorary Members.

Prof. H. H. Ballard, Pittsfield, Mass.
Rev. J. H. Langille, Kensington, Md.

Prof. H. Nehrling, Palm Cottage, Gotha, Fla.
Mr. Otto Widmann, St. Louis, Mo.

Associate Members.

Harold Bowditch, "Sunnyside," Jamaica Plain, Mass.
H. W. Carriger, 486 Eddy St., San Francisco, Calif.
Mrs. T. D. Dershimer, Square Top, Wyoming Co., Penn.
Irving A. Field, 61 Oxford St., Cambridge, Mass.
J. L. Floyd, Ervine Block, Canton, Ohio.
Harry S. Hathaway, Box 498, Providence, R. I.
Clarence E. Hefflefinger, R. F. D. No. 2, Plimpton, Ohio.
Karl J. Heilmann, 401 East Market St., Tiffin, Ohio.
Thomas D. Keim, 405 Redcliffe St., Bristol, Penn.
Miss Caroline Matthews, 90 Main St., Waterville, Me.
C. L. Metcalf, Plimpton, Ohio.
William J. Mills, Box 16, East Point, Ga.
G. F. Richardson, 116 South Main St., Mt. Pleasant, Mich.
Miss Isabella Sand, Ardsley-on-Hudson, N. Y.
Dr. Charles E. Slocum, Defiance, Ohio.
Miss Marion E. Sparks, Urbana, Ill.
Miss Jane L. Spencer, 239 East First St., Orange, N. Y.
Millard Van Wagner, 47 Leonard St., New York City.
Dr. Charles Vetter, 152 Second St., New York City.
Edward R. Warren, 20 West Caramillo St., Colorado Springs, Colo.
Rev. George F. Weld, Lincoln St., Hingham, Mass.
Miss Clara S. Wheatley, Medina, Ohio.
Norman A. Wood, 1216 S. University Ave., Ann Arbor, Mich.

ELECTION OF MEMBERS.

The following are proposed for Active Membership:—

Clinton G. Abbott, New York City.
William Brewster, Cambridge, Mass.
Paul Bartsch, Washington, D. C.
William Dutcher, New York City.
Dr. J. Dwight, Jr., New York City.
George P. Ells, Norwalk, Conn.
Louis Agassiz Fuertes, Ithaca, N. Y.
George E. Hix, New York City.
J. W. Kay, 62 Sheldon avenue, Detroit, Mich.
W. G. Kay, 62 Sheldon avenue, Detroit Mich.
Clark J. Peck, Philadelphia, Pa.
Prof. H. A. Surface, Harrisburg, Pa.
Mark L. C. Wilde, Camden, N. J.

Mr. Karl J. Heilmann, Tiffin, O., and Miss Marion E. Sparks, Urbana, Ill., are proposed for Associate Membership.

THE WARBLER

JOHN LEWIS CHILDS, EDITOR

WITH 1905 **The Warbler** begins a new series which will contain many superb Colored Plates of rare eggs such as Kirtland and Olive Warbler, Carolina Paroquet, Clark's Crow, Ipswich and Rufous Crowned Sparrow, Yellow and Black Rail, Calaveras Warbler, etc. Also splendid illustrations of Birds and Nests, and leading articles by well known authorities.

Published Quarterly, 32 Pages & Cover

SUBSCRIPTION PRICE
$1.00 PER YEAR. SINGLE COPIES 30c

Eggs of Kirtland Warbler will be figured in first issue (Jan. or Feb.) of the new series.

ADDRESS THE WARBLER
FLORAL PARK, N. Y.

Bird Papers For Sale

Complete volumes as well as odd numbers of Birds and Nature, Osprey, and many others.

Write for list and prices.

LYNDS JONES,
Oberlin, O.

DAWSON'S BIRDS OF OHIO.

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones, 720 quarto pages. 322 birds accurately described, their habits interestingly given. Combines ready identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations to WILSON BULLETIN, Oberlin, Ohio.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES.

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds of Ohio are treated. It does not attempt a description of the plumage. Price is 50 cents. Address, LYNDS JONES, Oberlin, Ohio.

Veterinary Elements.

What the Animal Husbandry Expert, Mr. Geo. M. Rommel, B. S. A., of the U. S. Dep't. of Agriculture, has to say regarding "Veterinary Elements":

"I do not feel that I can add anything to that which has been already said in commendation . . . . but from my own personal experience I can say that if I could have had the training in veterinary work that the little book embodies, my course in animal husbandry would have been all the more valuable to me.

"You have pointed out the way for a splendid line of work, and I sincerely hope that the leaders in agricultural education will not be slow in following it."

The author is A. G. Hopkins, B. Agr., D. V. M.; Farmer and Agricultural Journalist; late instructor in Stock Judging and Veterinary Science, Wisconsin Agricultural College.

The above work is clearly and plainly written and easily understood by any one. It can be secured, postpaid, for $1.50, from

THE WILSON BULLETIN, Oberlin, Ohio.

New and Revised Edition Just Issued.

Principles of Plant Culture.

BY E. S. GOFF,

Professor of Horticulture in the University of Wisconsin.

The text-book used in the classes in Plant Life and Horticulture, in the popular Short course in Agriculture, of the University of Wisconsin.

A Few of the Many Things that this Book Teaches.

How to select, test, plant store, and stratify seeds. How to promote flowering and fruiting. How to prevent sun-scald on fruit and other trees. How to promote wood maturity in autumn. How to water and care for greenhouse plants.

The Book Also Teaches

How to maintain the fertility of the soil. How to prevent damage from mice, gophers, rabbits, woodchucks, and birds. How to spray plants to prevent insect and fungus ravages. How to prevent damage from chinch bugs, army worms and grasshoppers. How to treat oats to prevent smut. How to make and apply the Bordeaux mixture. How to grow plants from cuttings, layers and stolons. How to propagate plants by grafting and budding. How to make and take care of hot-beds and cold-frames. How to transplant trees, shrubs and herbaceus plants. How to prune fruit and ornamental trees. How to improve plants by selection and cross pollination, and many other things equally valuable.

While the work is written in plain and concise language, the subjects are treated in a thoroughly scientific manner and embody the results of the most recent experiment and research.

The book contains 285 printed pages with 179 illustrations. It is well printed on excellent paper and is neatly and substantially bound in cloth.

This book will prove a most valuable addition to the library of every school of agriculture, horticulture, forestry, and of every rural high or common school in this country, as well as to the library of every farmer and fruit-grower.

Price, $1.00 per copy, postage paid, when ordered from

THE WILSON BULLETIN, Oberlin, Ohio.

University
JUL
21
1905

Old Series Vol. XVII. New Series Vol. XII, No. 2.

The Wilson Bulletin

No. 52.

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

June, 1905.

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.



CONTENTS.

Common Tern. Frontispiece Illustration.
A Year with the Birds in New York City. GEORGE E. HIX. 35
Birds vs. Street Cars. MARION E. SPARKS. 44
Autumn Birds of the Les Cheneaux Islands. WALTER C. WOOD. 48
The Spotted Sandpiper. CHRESWELL J. HUNT. 51
Bird Horizons from Russelville, Ky. G. C. EMBODY. 52
Common Tern. Illustration. 55
Winter Notes on Yellow-bellied Sapsucker (*Sphyrapicus varius*). C. H. MORRIS. 56
Nesting of the Yellow-bellied Sapsucker. J. CLAIRE WOOD. 57
All Day with the Birds at Durmid, Va. W. F. HENNINGER. 58
Some Further Suggestion for Bird Study. LYNDS JONES. 61
Two Goose Records from Ohio. W. F. HENNINGER. 63
Additions to the Birds of Ohio. LYNDS JONES. 64
Editorial. 65
General Notes. 66
Some Nesting Sites of the Hairy Woodpecker (*Dryobates villosus*), J. CLAIRE WOOD; Ruby-Crowned Kinglet in Winter, ZENO METCALF; The Dickcissel (*Spiza americana*) in Wayne Co., Mich., J. CLAIRE WOOD; Some Spring Records from the Vicinity of New York City, CARLETON SCHALLER.
Book Reviews. 67
Publications Received. 69

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, O., edited by Lynds Jones.

Subscription: Fifty cents a year, including postage, *strictly in advance*. Single numbers, 15 cents, unless they are "Special" numbers, when a special price is fixed. The BULLETIN, including all "Special" numbers, is free to all paid up members, either Active, Associate, or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for review, should be addressed to Lynds Jones, Oberlin, Ohio.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 15th of the month preceding publication.

COMMON TERN. (Sterna hirundo.)
Egg, Young Just Hatched, Young Two Days Old.
Woepecket Id., Mass.

THE
WILSON BULLETIN
NO. 51.
A QUARTERLY JOURNAL OF ORNITHOLOGY

VOL. XVII. JUNE, 1905. No. 2

A YEAR WITH THE BIRDS IN NEW YORK CITY.

BY GEORGE E. HIX.

The year 1904 was a very good one for birds in New York City, not only the common species being abundant, but the number of rare species observed being quite large also. The localities worked were Central, Bronx, and Van Cortlandt Parks, and St. Paul's church-yard in the old city, and Coney Island, Rugby, and Rockaway Beach in that portion of Long Island which is now part of the greater city.

Central Park, in the heart of Manhattan Island, is a good place in migrations, especially for Warblers. At the height of the migrations more than half the birds seen are Warblers. Excepting Robins comparatively few birds breed there, seventeen being the average number of species.

Bronx Park in the northern part of the city has been left mostly in the natural condition, and is the home of about sixty species. The Lawrence Warbler, which nested there the past year, drew more attention to that locality than had been given before. This is a good place in which to study Hawks, as they are not molested. As the larger part of the park is in charge of the New York Zoological Society, every protection and encouragement is given to the birds.

Although Coney Island is mostly taken as a pleasure resort, the extensive marshes back of the beaches are the homes of a number of marsh birds. Seaside and Sharp-tailed Sparrows are abundant summer residents there. During migrations shore and sea birds frequently are seen.

Rockaway Beach is a narrow strip of land separating Jamaica Bay from the ocean. On the "bay" side there are extensive marshes and mud flats. Myrtle bushes are plentiful and there is a small cedar-forest.

Rugby is open country, mostly stubble fields, and very few trees. Adjoining Rugby are the New Utrecht meadows, which are very extensive, and through which several creeks flow. This is an admirable locality for Snipe, Herons, and Rails. There is no protection given to birds there, and everything is shot at, especially by boys with "bean-shooters."

During migration I was in Central Park practically every morning and afternoon for at least half an hour. On the Saturday afternoons the other places were visited. Rockaway Beach excepted. The entire day is needed for that trip, and I was able to get there only twice, onMay 22 and November 20. More trips there would have greatly increased the number of species seen. The first trip to Rugby was on July 22. In September and October several trips were made there. On these trips I was accompanied by one or two friends.

The localities being briefly described, notes on the birds are given in the following list. All of the 160 species were seen in the limits mentioned above and during the past year. Where no localities are mentioned, it is understood the birds were generally distributed. In migrations they were often seen in trees in the city streets.

1. **Colymbus holbœlli.**—Holbœll Grebe. One seen at Coney Island on October 23.

2. **Colymbus auritus.**—Horned Grebe. Two seen at Rockaway Beach on November 20.

3. **Podilymbus podiceps.**—Pied-billed Grebe. Several seen in Central Park in the fall.

4. **Gavia imber.**—Loon. One seen at Coney Island on November 26.

5. **Stercorarius pomarinus.**—Pomarine Jaeger. Several Jaegers seen at both Rockaway Beach and Coney Island, and while most of them were undoubtedly Pomrine, there may have been some Parasitics also.

6. **Rissa tridactyla.**—Kittiwake Gull. A young bird came to within thirty feet of the board-walk at Manhattan (Coney Island) after a storm, November 26.

7. **Larus marinus.**— Great Black-backed Gull. Two adults were seen the same time as the above.

8. **Larus argentatus.**—Herring Gull. A very abundant winter resident everywhere.

9. **Larus delawarensis.**—Ring-billed Gull. Very common along the coast in October and November.

10. **Sterna hirundo.**—Common Tern. A flock of nine at Rockaway on May 22.

11. **Sula bassana.**—Gannet. An adult at Rockaway on November 20.

12. **Phalacrocorax diplophus.**—Double-crested Cormorant. One at Rockaway on November 20.

13. **Anas obscura.**—Black Duck. A flock of twenty-seven seen in the Hudson River on March 12. Common in Central Park in September.

14. **Clangula clangula americana.**—American Golden-eye. Four males flew over Bronx Park on October 16.

15. **Oidemia americana.**—American Scoter. One at Coney Island on March 5.

16. **Oidemia deglandi.**—White-winged Scoter. Common at Rockaway and Coney Island in November.

17. **Branta canadensis.**—Canada Goose. Several flocks were seen in the fall.

18. **Ardetta exilis.**—Least Bittern. A pair bred on Coney Island.

19. **Ardea herodias.**—Great Blue Heron. Two seen at Rugby on July 22.

20. **Butorides virescens.**—Green Heron. Common at Rugby in the fall and may breed.

21. **Nycticorax nycticorax nævius.**—Black-crowned Night Heron. Very common summer resident at most bodies of water.

22. **Rallus crepitans.**—Clapper Rail. One seen at Rugby on September 17.

23. **Porzana carolina.**—Sora. Several seen at Rugby in September.

24. **Actodromas maculata.**—Pectoral Sandpiper. A flock of seven at Rockaway on May 22, and a flock of six at Rugby on October 8.

25. **Actodromas minutilla.**—Least Sandpiper. Three at Rockaway, May 22.

26. **Totanus melanoleucus.**—Greater Yellow-legs. Two at Rugby, September 17.

27. **Helodromas solitarius.**—Solitary Sandpiper. A common migrant both on the coast and fresh water streams.

28. **Bartramia longicauda.**—Bartramian Sandpiper. Two at Rugby, September 17.

29. **Actitis macularia.**—Spotted Sandpiper. A common migrant wherever there is water.

30. **Ægialitis semipalmata.**—Semipalmated Plover. Very common at Rockaway on May 22. Several flocks were seen, which at one time united, forming one flock of at least 2000 birds.

31. **Ægialitis meloda.**—Piping Plover. One at Rockaway, May 22.

32. **Circus hudsonius.**—Marsh Hawk. Common at Rugby in the late summer and fall.

33. **Accipiter velox.**—Sharp-shinned Hawk. Common migrant in Central Park.

34. **Accipiter cooperii.**—Cooper Hawk. Central Park, one on October 3 and one on November 7.

35. **Buteo borealis.**—Red-tailed Hawk. A few were seen at Coney Island in November.

36. **Buteo lineatus.**—Red-shouldered Hawk. Common permanent resident in Bronx Park. One in Central Park, December 25.

37. **Buteo platypterus.**—Broad-winged Hawk. Two in Bronx Park, October 29. One in Central Park, October 30, and one on November 6.

38. **Falco peregrinus anatum.**—Duck Hawk. A fine adult male at Rugby, September 17.

39. **Falco columbarius.**—Pigeon Hawk. One at Rockaway, November 20.

40. **Falco sparverius.**—American Sparrow Hawk. Common at Rugby in September and October.

41. **Pandion haliaëtus carolinensis.**—American Osprey. One in Central Park, April 9.

42. **Asio accipitrinus.**—Short-eared Owl. Several seen along the coast in October and November.

43. **Coccyzus americanus.**—Yellow-billed Cuckoo. Common summer resident in the parks.

44. **Coccyzus erythrophthalmus.**—Black-billed Cuckoo. Uncommon summer resident in the parks.

45. **Ceryle alcyon.**—Belted Kingfisher. Common summer resident in Bronx Park. A few in Central Park in migrations.

46. **Dryobates villosus.**—Hairy Woodpecker. A pair spent the winter of 1903-4 in Central Park. A male is present this winter.

47. **Dryobates pubescens medianus.**—Downy Woodpecker. Common winter resident.

48. **Sphyrapicus varius.**—Yellow-bellied Sapsucker. Common migrant.

49. **Melanerpes erythrocephalus.**—Red-headed Woodpecker. Two in Central Park on September 18.

50. **Colaptes auratus luteus.**—Northern Flicker. Common summer resident. Was abundant in Central Park in fall.

51. **Chordeiles virginianus.**—Nighthawk. Common summer resident. Often seen flying over the house-tops at dusk.

52. **Chætura pelagica.**—Chimney Swift. Abundant summer resident.

53. **Trochilus colubris.**—Ruby-throated Hummingbird. Summer resident in Bronx and Van Cortlandt Parks. A migrant only elsewhere.

54. **Tyrannus tyrannus.**—Kingbird. Common summer resident.

55. **Myiarchus crinitus.**—Crested Flycatcher. Common summer resident in Bronx Park. Common migrant in Central Park.

56. **Sayornis phœbe.**—Phœbe. Common summer resident in Bronx Park. Common migrant elsewhere.

57. **Nuttalornis borealis.**—Olive-sided Flycatcher. One in Central Park on May 28, and one September 30.

58. **Empidonax flaviventris.**—Yellow-bellied Flycatcher. One in Central Park on September 8.

59. **Empidonax minimus.**—Least Flycatcher. Common migrant.

60. **Alauda arvensis.**—Skylark. There is a colony of 75–100 at Rugby. They may be seen any day in the summer, but after the song season are hard to find. They may spend the winter in another neighborhood.

61. **Otocoris alpestris.**—Horned Lark. Very common at Rockaway on November 20.

62. **Cyanocitta cristata.**—Blue Jay. Common permanent resident in Bronx Park. In Central Park, only one was seen in the spring, but it was common in October.

63. **Corvus brachyrhynchos.**—American Crow. Very common permanent resident.

64. **Corvus ossifragus.**—Fish Crow. Summer resident in Bronx Park.

65. **Sturnus vulgaris.**—Starling. Abundant permanent resident, chiefly around tall buildings.

66. **Dolichonyx oryzivorus.**—Bobolink. Common fall migrant at Rugby.

67. **Molothrus ater.**—Cowbird. Common summer resident in Bronx Park.

68. **Agelaius phœniceus.**—Red-winged Blackbird. Common summer resident in the marshes.

69. **Sturnella magna.**—Meadowlark. Common summer resident in Bronx and Van Cortlandt Parks. Permanent resident along the coast.

70. **Icterus spurius.**—Orchard Oriole. Several in Central Park in May.

71. **Icterus galbula.**—Baltimore Oriole. Common summer resident in the parks.

72. **Euphagus carolinus.**—Rusty Blackbird. Common migrant.

73. **Quiscalus quiscula.**—Purple Grackle. Abundant migrant. Local summer resident in Bronx and Central Parks.

74. **Quiscalus quiscula æneus.**—Bronzed Grackle. A fine typical male was seen in Central Park, March 13.

75. **Passer domesticus.** — English Sparrow. Abundant everywhere.

76. **Fringilla cœlebs.**—Chaffinch. There is a male in Central Park which has been there three years.

77. **Carpodacus purpureus.**—Purple Finch. Common fall migrant. in Bronx and Central Parks.

78. **Astragalinus tristis.**—American Goldfinch. Common migrant.

79. **Carduelis carduelis.**—European Goldfinch. A few were seen in Central Park in late winter and Spring. This species appears to be diminishing in numbers.

80. **Spinus pinus.**—Pine Siskin. A flock of 15 in Central Park on October 30. Three at Manhattan Beach, November 19.

81. **Pocecetes gramineus.**—Vesper Sparrow. Abundant summer resident at Rugby.

82. **Passerculus princeps.**—Ipswich Sparrow. One at Manhattan Beach, November 19.

83. **Passerculus sandwichensis savanna.**—Savanna Sparrow. Very abundant fall migrant at Rugby. Rather uncommon at other places.

84. **Coturniculus savannarum passerinus.**—Grasshopper Sparrow. Common summer resident at Rugby.

85. **Ammodramus caudacutus.**—Sharp-tailed Sparrow. Common summer resident at Rubgy and Coney Island.

86. **Zonotrichia leucophrys.**—White-crowned Sparrow. Several were seen in October in various places.

87. **Zonotrichia albicollis.**—White-throated Sparrow. Common winter resident.

88. **Spizella monticola.**—Tree Sparrow. Local winter resident. Very rare in Central Park.

89. **Spizella socialis.**—Chipping Sparrow. Common summer resident.

90. **Spizella pusilla.**—Field Sparrow. Common summer resident in Bronx and Van Cortlandt Parks.

91. **Junco hyemalis.**—Junco. Common winter resident. Most common in fall.

92. **Melospiza cinerea melodia.**—Song Sparrow. Common permanent resident.

93. **Melospiza lincolnii.**—Lincoln Sparrow. One was seen in Central Park, May 6.

94. **Melospiza georgiana.**—Swamp Sparrow.. Common summer resident in the marshes.

95. **Passerella iliaca.**—Fox Sparrow. Common migrant. A few present in Central Park this winter.

96. **Pipilo erythrophthalmus.**—Towhee. Common migrant. A few summer residents at Bronx Park.

97. **Cardinalis cardinalis.**—Cardinal. There are several pairs present all the year in Central Park.

98. **Zamelodia ludoviciana.**—Rose-breasted Grosbeak. A few in Central Park in May.

99. **Cyanospiza cyanea.**—Indigo Bunting. Summer resident in Bronx Park.

100. **Piranga erythromelas.**—Scarlet Tanager. Common spring migrant in Central Park. Summer resident in Bronx Park.

101. **Hirundo erythrogaster.**—Barn Swallow. Summer resident in the northern part of city; elsewhere a common migrant.

102. **Iridoprocne bicolor.**—Tree Swallow. Common spring and abundant fall migrant. Every evening from early in August until late in October this species could be seen overhead in countless numbers.

103. **Riparia riparia.**—Bank Swallow. Common summer resident at Van Cortlandt Park. Elsewhere a migrant. Abundant along the coast in fall.

104. **Stelgidopteryx serripennis.**—Rough-winged Swallow. One pair bred in Bronx Park. Very likely nesting in the masonry of a new bridge.

105. **Ampelis cedrorum.**—Cedar Waxwing. Common summer resident in Bronx Park. In Central Park only one flock was seen in May, but it was very common in the fall.

106. **Lanius borealis.**—Northern Shrike. An immature bird in Central Park, November 2.

107. **Lanius ludovicianus migrans.**—Migrant Shrike. An individual was present at Rugby from about September 3 until October 22, if not later.

108. **Vireo olivaceus.**—Red-eyed Vireo. Abundant summer resident.

108. **Vireo gilvus.**—Warbling Vireo. A few were seen in Central Park in migrations.

110. **Vireo flavifrons.**—Yellow-throated Vireo. Uncommon summer resident in Central Park.

111. **Vireo solitarius.**—Blue-headed Vireo. Tolerably common migrant.

112. **Vireo noveboracensis.**—White-eyed Vireo. A pair bred in Bronx Park.

113. **Mniotilta varia.**—Black and White Warbler. Common migrant. Summer resident in Bronx Park.

114. **Helmitheros vermivorus.**—Worm-eating Warbler. One in Central Park, August 15.

115. **Helminthophila pinus.**—Blue-winged Warbler. Common summer resident in the northern part of city.

116. **Helminthophila lawrencii.**—Lawrence Warbler. A male mated with female Blue-winged Warbler in Bronx Park. It was described in various journals.

117. **Helminthophila chrysoptera.**—Golden-winged Warbler. A female in Central Park, May 7, and a male, August 15.

118. **Helminthophila rubricapilla.**—Nashville Warbler. One in Central Park, May 11, and one September 22.

119. **Helminthophila peregrina.**—Tennessee Warbler. A young bird in Central Park, September 8, and an adult male in St. Paul's church-yard, September 22.

120. **Compsothlypis americana usneæ.**—Northern Parula Warbler. Common migrant.

121. **Dendroica coronata.**—Myrtle Warbler. Abundant migrant.

122. **Dendroica pensylvanica.**—Chestnut-sided Warbler. Common spring migrant. In the fall a few were seen in Central Park only.

123. **Dendroica castanea.**—Bay-breasted Warbler. Several in Central Park in May.

124. **Dendroica striata.**—Black-poll Warbler. Abundant migrant. In the fall in Central Park, this species outnumbered all others put together. The last were seen October 19.

125. **Dendroica blacburniæ.**—Blackburnian Warbler. Several in Central Park in May.

126. **Dendroica virens.**—Black-throated Green Warbler. Common migrant in Central Park.

127. **Dendroica vigorsii.**—Pine Warbler. Tolerably common migrant in Central Park.

128. **Dendroica palmarum.**—Palm Warbler. A very typical specimen was seen in Central Park September 15.

129. **Dendroica palmarum hypochrisea.**—Yellow Palm Warbler. Common spring migrant. Uncommon fall migrant.

130. **Dendroica discolor.**—Prairie Warbler. Several in Central Park in May.

131. **Seiurus aurocapillus.**—Oven-bird. Summer resident in Bronx Park. A common migrant elsewhere.

132. **Seiurus noveboracensis.**—Water-Thrush. Common migrant.

133. **Seiurus motacilla.**—Louisiana Water-Thrush. Common migrant.

134. **Geothlypis trichas brachidactyla.**—Northern Yellow-throat. Summer resident in Bronx Park. Common migrant.

135. **Icteria virens.**—Yellow-breasted Chat. Summer resident in Bronx Park. One in Central Park, May 5.

136. **Wilsonia mitrata.**—Hooded Warbler. An adult male in Central Park, May 1.

137. **Wilsonia pusilla.**—Wilson Warbler. Common in Central Park in May. One seen October 4.

138. **Wilsonia canadensis.**—Canadian Warbler. Common spring, uncommon fall migrant.

139. **Setophaga ruticilla.**—American Redstart. Common summer resident in the parks.

140. **Anthus pensilvanicus.**—American Pipit. An abundant fall migrant along the coast.

141. **Galeoscoptes carolinensis.**—Catbird. Common summer resident.

142. **Toxostoma rufum.**—Brown Thrasher. Common migrant. Summer resident in Bronx Park.

143. **Thryothorus ludovicianus.**—Carolina Wren. Two in Central Park, September 22.

144. **Troglodytes aëdon.**—House Wren. A few in Central Park in May.

145. **Olbiorchilus hiemalis.**—Winter Wren. Several in Central Park in the fall.

146. **Cistothorus stellaris.**—Short-billed Marsh Wren. One was seen with other migrants at Rugby October 8. It may be a summer resident there.

147. **Telmatodytes palustris.**—Long-billed Marsh Wren. Summer resident in the salt marshes.

148. **Certhia familiaris americana.**—Brown Creeper. Uncommon winter resident.

149. **Sitta carolinensis.**—White-breasted Nuthatch. There were a few in Bronx and Central Parks last winter (1903–4), but there are none this winter.

150. **Sitta canadensis.**—Red-breasted Nuthatch. One spent the winter of 1903–4 in Central Park. In the fall one was seen in Central Park, October 15, and one in Bronx Park, October 16. The year before they were abundant in the fall from August to November.

151. **Parus atricapillus.**—Chickadee. Common winter resident.

152. **Regulus satrapa.**—Golden-crowned Kinglet. Winter resident in Bronx Park. Elsewhere a common migrant.

153. **Regulus calendula.**—Ruby-crowned Kinglet. Very common migrant.

154. **Hylocichla mustelina.**—Wood Thrush. Uncommon summer resident in Bronx and Central Parks.

155. **Hylocichla fuscescens.**—Wilson Thrush. Common migrant. Summer resident in Bronx Park.

156. **Hylocichla aliciæ.**—Gray-cheeked Thrush. Tolerably common migrant in Central Park. In early October a few, apparently Bicknell Thrushes (H. a. bicknelli), were seen in Central Park and St. Paul's Church-yard.

157. **Hylocichla swainsoni.**—Olive-backed Thrush. Abundant migrant.

158. **Hylocichla guttata pallasii.**—Hermit Thrush. Abundant migrant. There is one in Central Park this winter.

159. **Merula migratoria.**—American Robin. Abundant summer resident. A few are present in Central Park this winter.

160. **Sialia sialis.**—Bluebird. Summer resident in Bronx and Van Cortlandt Parks. In Central Park seen only in fall migration when it was quite common.

161. **Ammodramus maritimus.**—Seaside Sparrow. Abundant summer resident at Rugby and Coney Island.

BIRDS VS. STREET CARS.

BY MARION E. SPARKS.

The advance of the interurban electric lines, makes excursions to country places more easy, but are the birds near it undisturbed by the shrill whistle? Of course the birds may become used to it, for they pay little attention to railroad trains; but the frequency of the cars may make a difference.

The following notes give the birds seen without careful search, in a space of two lots; along the south edge is a cutting and beyond that a garden equal to a city block in size. During the time the notes were taken, interurban, local and construction cars passed to and fro in the cutting from 4 a. m. till 12 p. m. daily, at intervals of from thirty to three minutes.

The birds were only passing and the ones that remained all summer were noticeably more numerous than they had been in previous years.

Lack of close watching, no doubt missed many. Absence of cats, dogs, chickens, and children may account for the number in part, while the presence of a small stream in the space beyond the garden probably had some influence.

The yard is almost crowded with shrubs, while a dozen shade trees, and twice as many fruit trees helped to make it easy for the birds to escape too close notice.

Time, April 5–May 13, 1904.

April 5, 1904.—Ruby-crowned Kinglet, on a vine at the open window, later in a bush; fearless, almost tame, i.e. allowed observer to approach nearer than ten feet.

April 6-13.— A few, 3 or 4 Chickadees, one White-breasted Nuthatch, Hairy Woodpecker, Flicker, Robin, Wood Thrush, Bluebird, White-throated Sparrow, Veery, Towhee (male only), Brown Thrasher, Brown Creeper, Junco, Bronzed Grackle, Blue Jay.

Cooler April 14–17.

April 17.—The first House Wren of the season; two weeks later than last year.

April 25.—Black and White Warbler seen; only one specimen. The Blackbirds have not been seen since April 20. No

Towhees here since the 22d of April. Robins are plenty now, and two pairs of Brown Thrashers are nest-building.

The Wood Thrushes are fewer in number; the flocks of White-throated Sparrows are gone, too. The Veery (?) and Song Sparrow are often heard.

Cold and rainy, but the Cardinal came to-day, and the first Oven-bird. Male Cardinal was very busy in a brush-heap, feeding the female, who disdained even looking for food for herself. The Cardinals do not notice the interurban car, even when it whistles not twenty feet from them.

April 27.—The earliest Catbird came; the Bluebirds are here again, tame as ever. They were very curious about a bonfire, hardly waiting for the flames to die down before investigating it. The Indigo Bunting was here, taking a drink from a pan of water about thirty feet from the kitchen door.

The Black and White Warbler came again; he "fished up" an angleworm somewhere and was seen pounding it vigorously on a small branch, before he finally ate it. Did he pull it out of the ground as Robins do?

April 30.—Male and female Rusty Blackbird appeared; later they made a nest in a pear tree about ten feet from the house.

May 2.—Hummingbird in the tulip bed.

May 5.—Oven-birds still here, but the flock, or number, is smaller. The Ruby-crowned Kinglet is here yet; flocks of White-throated Sparrows are here morning and evening. A Sora (Carolina Rail), was in the yard this morning. It seemed confused, and attempts to get a closer view resulted in driving it away. Wilson Warbler here to-day.

May 6, 6:00 p. m.—Saw and heard the Rose-breasted Grosbeak.

May 8.—The Least Flycatcher was observed in an apple-tree; shy, and very busy. The Wrens are nest-making. A female Towhee was here; the others earlier, March 9–April 22, were all males, and have been gone more than ten days; they were fearless, but this one is very shy.

The Wood Thrushes are gone, so are the White-throated Sparrows. A Warbler, the Cerulean, almost surely, was here to-day; shy.

May 9.—The Chestnut-sided Warbler; not shy; not ten feet away from the observer and the open house door.

May 11.—Canadian Warbler flashed into sight in the lower boughs of a soft maple, for a few minutes, just long enough to make sure of his identity.

May 13.—The Chimney Swifts are here; and another different warbler, Magnolia? Probably not.

May 18.—Male and female Redstart in the peach tree, "for this day only."

The Baltimore Oriole heard once or twice; was not seen till June; so that he can hardly be counted.

The first Towhees came March 9; Blue Jays are here all winter; Robins too, but not in large numbers. About March 20, the Golden-crowned Kinglet was seen, and a flock of Lark Sparrows, Downy Woodpecker, and Sapsucker. The Robins are on record in my notes as carrying nest material, March 31.

Of those in the first list the following had nests in the yard, or the garden beyond it:

Robin.
Blue Jay.
Flicker.
Brown Thrasher.
House Wren.
Oven-bird.
Catbird.
Bronzed Grackle.
Hummingbird.
Rose-breasted Grosbeak (apparently).
Baltimore Oriole.
Chimney Swift.

The last in a chimney of the house, where they had nested for more than ten years.

As for the numbers of the various sorts the largest numbers seen at any one time were:

Robin, 10 or 12.
Blue Jay, 10 or 12.
Brown Creeper, 2 or 3.
Wilson Thrush, 5 or 6.
Downy Woodpecker, 2.
Towhee (male), 30.
Towhee (female), 1.
Flicker, 4.
Lark Sparrow, 15.
Golden-crowned Kinglet, 1.
Bronzed Grackle, 7 or 10.
Wood Thrush, 10 or 12.
Hairy Woodpecker, 2.
Bluebird, 4.
Yellow-bellied Sapsucker, 1.
Ruby-crowned Kinglet, 2.
Chickadee, 4 or 5.
Brown Thrasher, 4.
House Wren, 4.
White-throated Sparrow, 20.
Black and White Warbler, 1.
Cardinal, 4.
Oven-bird, 10 to 16.
Catbird, 6.
Indigo Bunting, 1.
Rusty Blackbird, 2.

Sora, 1.
Ruby-throated Hummingbird, 1.
Rose-breasted Grosbeak (all males), 6.
Baltimore Oriole, 1.
Chimney Swift, 4.
American Redstart, 2.
Least Flycatcher, 1.
Wilson Warbler, 1.
Chestnut-sided Warbler, 1.
Canadian Warbler, 1.
Cerulean Warbler, 1.
Magnolia, 1.

Total, at least 160, probably 175 individuals.

For the same period of time, and place, the previous year, 1903, when no cars ran up the cutting at all, the birds noticed were:

Blue Jay.
Robin.
Bronzed Grackle.
Brown Thrasher.
Wood Thrush.
Golden-crowned Kinglet.
Cardinal.
Rose-breasted Grosbeak.
Towhee.
House Wren.
Chipping Sparrow.
Baltimore Oriole.
Junco.
Lark Sparrow.
Wilson Thrush (?).
Chimney Swift.

A total of 16, as compared with 37 in 1904.

As a check on this, the birds seen in a piece of woodland a mile away from the car track, July 13, 5 to 10 a. m., may be of interest.

Indigo Bunting, 1.
Field Sparrow, 3.
Barn Swallow, 10.
Phœbe, 4.
Song Sparrow, 2.
Migrant Shrike, 1.
Bluebirds (some young ones), 12.
Yellow-billed Cuckoo, 1.
Chipping Sparrow, 4.
Meadowlark, 3.
American Goldfinch, 5.
Kingbird, 10.
American Crow, 7.
Blue Jay, 7.
Robin, 10.
Bronzed Grackle, 20.

Total, 100.

Urbana, Ill.

AUTUMN BIRDS OF THE LES CHENEAUX ISLANDS.

BY WALTER C. WOOD.

To the bird observer there is a deep interest and pleasant anticipation in exploring a new section of country, especially where a picturesque wilderness lends an added charm. The exquisite beauty of the wild rice marshes, the deep solitude of the thick cedar woods, the rugged magnificence of the windfall tangles and the clear waters reflecting sky and bordering forest form a picture of wild natural beauty never to be forgotten, and such a country is the Les Cheneaux Islands and adjacent mainland in T. 42. N. R. I. W., Macinac county, Michigan.

There is but little cultivated land in the vicinity of these islands. Some tamarack, balsam, pine, and birch trees are to be found, but spruce, hemlock and poplar are more common, while the cedar is the predominating tree and forms extensive swamps.

It is to be regretted that I could not devote my entire time to bird observation, but every opportunity was embraced, and probably very few species escaped notice. Of those observed, however, eight species could not be identified with certainty and are consequently not listed. It might also be mentioned that while no owls were met with one specimen of the Great Horned was shot a few days before my arrival.

Captain John Pollock of the Les Cheneaux Club stated that during his fifteen years residence in the vicinity of these islands he has never observed the Woodcock or Scarlet Tanager, birds he was familiar with in southern Michigan; also a small flock of Passenger Pigeons are observed each year. Of course, as none of these pigeons have been secured their identity is not beyond question. He showed me a nest of the Osprey, from which a gentleman from Chicago took three young and secured the parent birds; and also mentioned one of the Northern Ravens in a pine which had been occupied for many years up to 1900.

The following list represent the birds observed between October 15th and November 15th, 1903:

1. **Colymbus auritus.** Horned Grebe.

2. **Podilymbus podiceps.** Pied-billed Grebe.—These two species could be seen on the bays at all times, the former considered good eating by the residents.

3. **Gavia imber.** Loon.—Several observed.

4. **Larus argentatus.** Herring Gull.—Many seen. Captain Pollock showed me a nest on a small island within 300 yards of the club house, where a brood had been successfully reared during the summer. The pair have nested there five or six years, but no others breed in the vicinity, although the birds may be seen all summer in more or less numbers.

5. **Larus delawarensis.** Ring-billed Gull.—Six were observed one morning on piles in front of club house.

6. **Larus philadelphia.** Bonaparte Gull.—One seen with the last species.

7. **Merganser americanus.** American Merganser.—Common, breeds. Captain Pollock stated that many pairs with young were on the bay during the summer and became very tame, one brood coming to the landing to be fed.

8. **Merganser serrator.** Red-breasted Merganser.—A male dropped into my decoys, but dove at the flash and reappeared out of range. This was the only specimen I could positively identify.

9. **Lophodytes cucullatus.** Hooded Merganser.—Rather common and considered good eating. Pollock says, "a common breeder."

10. **Anas boschas.** Mallard.

11. **Anas obscura.** Black Duck.

12. **Aythya americana.** Redhead.—The various indentations of the bay were covered with wild rice and visited by flocks of the three above species ranging from ten to fifty birds.

13. **Aythya affinis.** Lesser Scaup Duck.—Common.

14. **Clangula clangula americana.** American Golden-eye.—A few small flocks.

15. **Charitonetta albeola.** Buffle-head.—Most abundant of all the ducks. Flocks of 500 or more often seen.

16. **Oidemia americana.** American Scoter.—One bird secured from a flock of five. No others seen.

17. **Branta canadensis.** Canada Goose.—Several small flocks.

18. **Totanus melanoleucus.** Greater Yellow-legs.

19. **Charadrius dominicus.** American Golden Plover.—Flocks of five to twenty of this plover and yellow-legs seen daily and many secured.

20. **Bonasa umbellus.** Ruffed Grouse.—An abundant resident. Its greatest enemies are the foxes and half-starved Indian dogs.

21. **Buteo borealis.** Red-tailed Hawk.—One seen.

22. **Buteo lineatus.** Red-shouldered Hawk.—One seen.

23. **Haliæetus leucocephalus.** Bald Eagle.—One, adult specimen, came daily to the outer bay.

24. **Falco sparverius.** American Sparrow Hawk.—Several seen.

25. **Dryobates villosus.**. Hairy Woodpecker.—Common.

26. **Dryobates pubescens medianus.** Downy Woodpecker.—Common. Most abundant of the woodpeckers.

27. **Picoides arcticus.** Arctic Three-toed Woodpecker.—Most abundant of the woodpeckers.

28. **Ceophlœus pileatus abieticola.** Northern Pileated Woodpecker.—Common. About 30 seen and one secured.

29. **Colaptes auratus luteus.** Northern Flicker.—Several observed.

30. **Cyanocitta cristata.** Blue Jay.—Common.

31. **Perisoreus canadensis.** Canada Jay.—A rather common and conspicuous species found mainly along the Indian trails and deer runs.

32. **Corvus brachyrhynchos.** American Crow.—Several seen.

33. **Pinicola enucleator leucura.** Northern Pine Grosbeak.—Many large flocks noted and a very small percentage in the red plumage.

34. **Loxia curvirostra minor.** American Crossbill.—Many small flocks observed.

35. **Acanthis linaria.** Red-poll. One small flock.

36. **Spinus pinus.** Pine Siskin.—Very abundant.

37. **Passerina nivalis.** Snowflake.—Common. Always in large flocks.

38. **Spizella monticola.** Tree Sparrow.—Common.

39. **Junco hyemalis.** Slate-colored Junco.—Common.

40. **Melospiza cinerea melodia.** Song Sparrow.—Three seen.

41. **Lanius borealis.** Northern Shrike.—The only specimen seen was secured.

42. **Lanius ludovicianus migrans.** Migrant Shrike. Several observed.

43. **Sitta carolinensis.** White-breasted Nuthatch.—Common.

44. **Sitta canadensis.** Red-breasted Nuthatch.—Not so abundant as the preceding.

45. **Parus atricapillus.** Chickadee.—This energetic little bird was distributed throughout the country visited.

46. **Regulus calendula.** Ruby-crowned Kinglet.—One brought me for identification.

46. **Merula migratoria.** American Robin.—Several individuals were noted at various times.

48. **Sialia sialia.** Bluebird.—Several small flocks seen. None observed after November 1st.

Detroit, Mich.

THE SPOTTED SANDPIPER.

CHRESWELL J. HUNT.

Surely every bird student knows the Spotted Sandpiper (*Actitis macularia*). No matter how small the body of water we visit, whether river, creek, or mill-pond, we are almost certain to find at least one pair of these little fellows at home. Visit a rapid stream dashing over rocks and gliding here and there into deep pools such as the trout love and you will hear the shrill "Peet-weet, peet-weet" of the Sandpiper as he takes wing from some point along the water's edge at your approach; or you may come upon him "teetering" upon some rock. Or visit a tide creek or river and you will find him just as much at home upon the sandy beach or upon the mud flats left by the receding tide. He also finds the quiet mill-pond much to his liking and the stream must be small indeed which may not be honored by his presence.

Although so well known generally a few observations made during the past year or two may not be amiss.

Common enough along the streams I frequent, and though one of the first birds whose acquaintance I made when I began the study of ornithology, it is still more or less of a stranger to me. We are on speaking terms and that is all. Never as yet have I been fortunate enough to discover the nest or even to receive a hint from the birds as to the nest location.

It was a hot June afternoon. I had been lying upon a log at the edge of a mill-pond watching a Kingfisher. The mid-day hush in the bird chorus was in progress and 'ere long I fell asleep. When I awakened, without moving I looked out over the water and there within only a few yards of me stood two Spotted Sandpipers. They were standing in about three inches of water and were apparently fast asleep. Each stood upon one leg. The head was turned back and the bill was hidden in the feathers of the back. They were perfectly motionless. I wondered if it was customary for these birds to sleep in this manner. Do they always sleep thus? Or were these birds simply taking a siesta?

Mr. C. J. Peck and myself spent most of the nights of May 28 and 29, 1904, in an open boat upon a New Jersey tide creek. During the early night and even at midnight we heard a strange whistle which some bird made as it flew across the marsh. It was of course too dark to see the bird, but on the following evening we heard a Spotted Sandpiper give the same note. Is this nocturnal flight common with this species? Has it anything to do with the breeding season? Is this bird astir all night? I have found them sleeping during the day and flying about at night. If they sleep at night do they sleep standing in the water or where do they roost? Can not some other bird student throw some more light upon this subject?

We always associate the Spotted Sandpiper with the vicinity of water. Whenever I think of them I imagine I can hear water rushing about rocks or see before me an expanse of mud flats. What was my surprise, therefore, when one July morning I found one of them walking the railroad tracks. To be sure it was near a trestle that spanned the creek, but here was the bird walking back and forth upon the steel rail, occasionally stopping and "teetering" in its customary way. It was a young bird of the year, lacking the conspicuous spots upon the breast. As it walked the rail the toes were turned in and it was "pigeon-toed" to perfection. I soon found, however, that there was an object in its madness as I saw it several times drop down to one of the ties and capture an unsuspecting spider, which it devoured with apparent relish.

Philadelphia, Pa., May, 1905.

BIRD HORIZONS FROM RUSSELLVILLE, KY.

BY G. C. EMBODY.

March 19, 1904.

Time, 2:30–5:30 P. M.; temp., 70°; clear; light south wind.

Wilson Snipe, 3; Bob-white, 15; Turkey Vulture, 10; Red-bellied Woodpecker, 2; Flicker, 10; Bronzed Grackle, 50; Vesper Sparrow, 20; Savanna Sparrow, 5; Leconte Sparrow, 1; White-crowned Sparrow, 2; White-throated Sparrow, 10; Field Sparrow, 8; Slate-colored Junco, 50; Song Sparrow, 12; Swamp Sparrow, 1; Cardinal,

10; Myrtle Warbler, 6; Tufted Titmouse, 5; Carolina Chickadee, 10; Golden-crowned Kinglet, 4.

Total: Species, 20; individuals, 234.

April 27, 1904.

Time, 2–6 P. M.; temp., 75°; cloudy.

Bob-white, 1; Turkey Vulture, 6; Downy Woodpecker, 1; Pileated Woodpecker, 1; Chimney Swift (estimate), 100; Crested Flycatcher, 5; Wood Pewee, 2; Phœbe, 1; Blue Jay, 5; Crow, 10; Purple Finch (est.), 100; Goldfinch, 10; White-throated Sparrow, 5; Chipping Sparrow, 10; Field Sparrow, 30; Towhee, 2; Cardinal, 5; Scarlet Tanager, 4; Summer Tanager, 5; Cliff Swallow, 1; Red-eyed Vireo, 3; Warbling Vireo, 1; White-eyed Vireo, 6; Black and White Warbler, 3; Worm-eating Warbler, 1; Cape May Warbler, 1; Prairie Warbler, 3; Myrtle Warbler, 10; Oven-bird, 3; Black-poll Warbler, 2; Mockingbird, 5; Catbird, 6; Brown Thrasher, 3; Carolina Wren, 1; Tufted Titmouse, 5; Carolina Chickadee, 3; Blue-gray Gnatcatcher, 10; Wilson Thrush, 1; Robin, 10; Bluebird, 4.

Total: Species, 40; individuals, 385.

April 30, 1904.

Time, 10 A. M.–5 P. M.; temp., 80; clear; light east wind.

Bob-white, 1; Mourning Dove, 4; Turkey Vulture, 10; Broad-winged Hawk, 1; Downy Woodpecker, 1; Flicker, 4; Chimney Swift, 40; Hummingbird, 1; Kingbird, 6; Crested Flycatcher, 10; Blue Jay, 15; Crow, 10; Meadowlark, 2; Bronzed Grackle, 5; Orchard Oriole, 1; Baltimore Oriole, 3; Purple Finch, 40; Goldfinch, 10; White-throated Sparrow, 10; Chipping Sparrow, 10; Field Sparrow, 20; Savanna Sparrow, 2; Swamp Sparrow, 1; Lincoln Sparrow, 2; Bachman Sparrow, 1; Rose-breasted Grosbeak, 4; Indigo Bunting, 5; Cardinal, 8; Towhee, 20; Summer Tanager, 3; Scarlet Tanager, 2; Bank Swallow, 2; Rough-winged Swallow, 1; Red-eyed Vireo, 10; White-eyed Vireo, 10; Yellow-throated Vireo, 5; Black and White Warbler, 6; Prairie Warbler, 6; Myrtle Warbler, 2; Black-throated Green Warbler, 3; Cerulean Warbler, 2; Black-poll Warbler, 6; Yellow Palm Warbler, 1; Maryland Yellow-throat, 10; Yellow-breasted Chat, 15; Redstart, 2; Ovenbird, 6; Mockingbird, 5; Catbird, 20; Brown Thrasher, 15; Carolina Wren, 6; Red-breasted Nuthatch, 1; Tufted Titmouse, 10; Carolina Chickadee, 8; Ruby-crowned Kinglet, 6; Blue-gray Gnatcatcher, 20; Robin, 18; Bluebird, 12.

Total: Species, 58; individuals, 459.

May 14, 1904.

Time, 9 A. M.–5 P. M.; temp., 75°; cloudy; light south wind.

Bob-white, 20; Mourning Dove, 5; Turkey Vulture, 10; Yellow-billed Cuckoo, 3; Downy Woodpecker, 1; Pileated Woodpecker, 1; Red-headed Woodpecker, 3; Kingbird, 20; Wood Pewee, 2; Prairie

Horned Lark, 2; Crow, 30; Meadowlark, 10; Bronzed Grackle, 30: Orchard Oriole, 10; Baltimore Oriole, 5; White-throated Sparrow, 1; White-crowned Sparrow, 1; Chipping Sparrow, 7; Field Sparrow, 12; Grasshopper Sparrow, 25; Lincoln's Sparrow, 1; Indigo Bunting, 10; Dickcissel, 3; Cardinal, 8; Red-eyed Vireo, 10: White-eyed Vireo, 10; Yellow-throated Vireo, 1; Warbling Vireo, 1; Black and White Warbler, 4; Prairie Warbler, 2; Magnolia Warbler, 1; Yellow Warbler, 2; Black-poll Warbler, 2; Oven-bird, 8; Yellow-breasted Chat, 15; Mockingbird, 6; Catbird, 20; Brown Thrasher, 5; Carolina Wren, 6; Bewick Wren, 2; Tufted Titmouse, 5; Carolina Chickadee, 4; Wilson Thrush, 2; Wood Thrush, 2; Robin, 28; Bluebird, 15.

Total: Species, 46; individuals, 371.

August 25, 1904.

Time, 2–5 P. M.; temp., 95°; clear.

Solitary Sandpiper, 1; Mourning Dove, 15; Turkey Vulture, 3; Red-headed Woodpecker, 5; Kingbird, 10; Crested Flycatcher, 2; Wood Pewee, 1; Crow, 5; Field Sparrow, 8; Grasshopper Sparrow, 3; Bachman Sparrow, 10; Cardinal, 3; Red-eyed Vireo, 1; Yellow-throated Vireo, 1; Berwick Wren, 1; Tufted Titmouse. 5; Carolina Chickadee, 4; Bluebird, 20.

Total: Species, 20; individuals, 101.

September 17, 1904.

Time, 10 A. M.–5 P. M., temp., 90°; clear.

Solitary Sandpiper, 1; Bob-white, 1; Mourning Dove, 8; Turkey Vulture, 20; Black-billed Cuckoo, 1; Downy Woodpecker, 1; Red-headed Woodpecker, 5; Red-bellied Woodpecker, 5; Pileated Woodpecker, ; Flicker, 6; Nighthawk, 12; Chimney Swift, 20; Phœbe, 1; Acadian Flycatcher, 1; Wood Pewee, 5; Yellow-bellied Flycatcher, 2; Crow, 5; Blue Jay, 5; Goldfinch, 1; Field Sparrow, 8; Grasshopper Sparrow, 4; Summer Tanager, 2; Cardinal, 20; Red-eyed Vireo, 2; White-eyed Vireo, 2; Ovenbird, 1; Chestnut-sided Warbler, 1; Black-throated Green Warbler, 5; Magnolia Warbler, 1; Mockingbird, 6; Bewick Wren, 2; Tufted Titmouse, 10; Carolina Chickadee, 6; Robin, 10; Bluebird, 12.

Total: Species, 35; individuals, 193.

December 28. 1904.

Time, 2–5 P. M.; temp., 20°; north west wind: clear.

Bob-white, 30; Mourning Dove, 5; Turkey Vulture, 3; Downy Woodpecker, 2; Hairy Woodpecker, 1; Flicker, 20; Blue Jay, 3: Crow, 10; Meadowlark, 25; Leconte Sparrow, 1; Tree Sparrow,, 12; Song Sparrow, 3; Junco, 100; Cardinal, 4; Towhee, 1; Cedar Waxwing, 10; Myrtle Warbler, 10; Mockingbird, 5; Brown Creeper, 1; Tufted Titmouse, 10; Carolina Chickadee, 5; Hermit Thrush, 1; Bluebird, 8.

Total: Species, 23; individuals, 270.

January 21, 1905.

Time, 10 A. M.–5 P. M.; temp., 32°–40°; north west wind; clear.

Turkey Vulture, 15; Downey Woodpecker, 3; Yellow-bellied Sapsucker, 1; Flicker, 2; Prairie Horned Lark (est.), 90; Crow, 20; Blue Jay, 1; Meadowlark, 2; Purple Finch, 12; Goldfinch, 4; Tree Sparrow, 6; Field Sparrow, 20; Junco (est.), 100; Song Sparrow, 4; Towhee, 1; Cardinal, 10 Cedar Waxwing, 10; Loggerhead Shrike, 1; Myrtle Warbler, 30; Mockingbird, 4; Carolina Wren, 3; Bewick Wren, 1; Tufted Titmouse, 5; Carolina Chickadee, 20; Golden-crowned Kinglet, 1; Robin (flock), 100; Bluebird, 5.

Total: Species, 27; individuals, 471.

COMMON TERN. (Sterna hirundo.)
Nest with Egg and Two Just Hatched Young, and Two Half Grown Young.
Woepecket Id., Mass.

WINTER NOTES ON YELLOW-BELLIED SAPSUCKER (*Sphyrapicus varius*).

BY C. H. MORRIS.

On January 15, 1905, on one of the coldest days of the winter, Yellow-bellied Sapsucker—a fine male—dropped in near my lunch counter, and from that time until now, February 15, he has been about nearly every day. The first time I saw him he was sticking close to the ground on the southeastern side of a maple catching the first level rays of the sun. With shoulders rounded and feathers ruffled he looked to be as cold as was the weather. On the 23d, the temperature having risen to thawing point, he drilled his row of holes in the maple, where first I saw him, and during the forenoon never once left his post. The sap ran sluggishly and his appearance matched it. I had noticed him flying about a couple of plum trees upon which the dried fruit hung in abundance—Dawson plums they are—and on the 29th, while watching his scarlet-throated lordship, I saw him pluck a plum. He did not insert it in a crack as his family does with nuts, but ate it partially, dropping the stone. His favorite method is taking small bits of the fruit as it hangs. On the 27th, the sap from the holes had frozen into icicles some five inches long, and Yellow-belly sat below, and nipping off the points leisurely, ate the ice. Yesterday, February 14, the coldest day we have had, with the temperature 8°, the bird flew on the lunch-box in which I had just placed cracked hickorynuts and walnuts. Close by him was suet in great plenty, which the Downy and Hairy feed from almost exclusively, seeming to have no taste for the nuts. But Yellow-bellied ate greedily of the nuts, thus following in the footsteps of his brother, Red-bellied Woodpecker, who scarcely ever will touch the meat or suet.

Mr. Dawson, in his charming work, "The Birds of Ohio," gives this latitude as about the northern winter range of this bird. February 11, another one came, this time an immature specimen, but he only remained for two or three days.

The birds are in the habit of frequenting a spruce and pine tree close by, and several times they have been observed eat-

ing the small nodules of the pitch that has, during the preceding year, exuded from the tree.

Morgan Co., O.

NESTING OF THE YELLOW-BELLIED SAPSUCKER.

BY J. CLAIRE WOOD.

I know of but two sets of eggs of this species taken in the county. I was collecting birds in a thick woods on P. C. 49, Ecorse township, on May 10, 1891, when I met with a party of young egg collectors. Soon after I heard one calling that he had found a Downy Woodpecker's nest. On being told the birds were Yellow-bellied Sapsuckers, he was greatly surprised. The cavity contained three fresh eggs, which he left for a larger set, but there is a humorous sequel. It seems that after I left a consultation was held and it was decided to take the eggs for fear I might return and do a little collecting myself. When the collector arrived home his father, a well known oologist, decided to return at once and secure the bird in order to discover, by dissection, the number of eggs that would have been laid. This was done, and when he cut open the bird next day, Great Cæsar! it was the male.

This is the most abundant and noisy woodpecker of the woodlands here during April. Before the expiration of the month the majority are already mated and looking for nesting sites, but after drilling numerous test holes, in one case devoting three days to a single excavation, they apparently feel a renewal of the migratory impulse and fade away toward the north; so when, on April 21, 1903, in Section 3, Van Buren township, I saw a female drilling into a dead stub in the thickest portion of a low swampy woods, I merely watched her awhile and passed on without further consideration. Chancing to pass the stub on May 19 I tapped it and out popped madame's head—the surprise was mutual. As I ascended and paused at her level she remained motionless, in fact, did not move until my finger was extended, when she retreated before it, but paused just out of reach and proceeded to drill a hole into the end. As it was withdrawn she followed closely until

her head protruded as before. This was repeated many times, she never forgetting to pound the end with a good will. A puff of smoke, blown into the cavity, finally induced her to vacate, but so small was the entrance that she actually stuck tight for about ten seconds before she could squeeze through. She eyed me a moment from the nearest tree trunk, then returned and clung to the stub only a few feet above my head. The excavation was twelve feet above the ground and the stub about twenty high. The entrance went straight in for three inches and was eighteen in depth, widening out to six at the bottom and occupying the core of the tree. It contained four fresh eggs, best described as exactly like average bluebird's in size and shape, but, of course, pure white. This was probably an incomplete set, as both Dr. P. E. Moody and myself have taken sets of six eggs in Oakland county, where the birds exhibited none of the courage of this individual.

Wayne County, Mich.

ALL DAY WITH THE BIRDS AT DURMID, VA.

BY W. F. HENNINGER.

This year I had to make my "All Day with the Birds" in a new territory. Having arrived at Lynchburg, April 25th, I immediately went to work to study the bird world. For that reason I preferred staying with friends at Durmid, as the mountains are only two miles distant. In some respects the bird world is similar to that of middle southern Ohio, for the climate is practically the same. Bewick Wren and Bachmann Sparrow greeted me, together with the Prairie Warbler the very first day.

The hills are crowded with woods, both decidious trees and pines; little streams gushing down from the hillsides; stones are abundantly sown over the shining brick-red soil, and bird-life lacks the vigor and freshness of the North.

Hawks seemed scarce and the great numbers of Turkey Vultures could not atone for this. Owls I did not see at all. Woodpeckers were persent, but only in very few individuals. The Robin was quite rare, the Meadowlark was heard but

twice, the Baltimore Oriole I found at Rivermount Park only, while the Grackles were confined almost entirely to Spring Hill cemetery. It took me almost a week to find a good corner for the study of the Warblers, but here they were thick and I have never made a better study of the Blackburnian, Chestnut-sided and Parula Warblers' songs than at this place. In the intervals could be heard the low notes of a pair of Worm-eating Warblers, the lazy weird song of the Prairie Warbler, and above them all the strong chant of the Carolina Wren: while as soon as you stepped a little farther out of the forest, Catbird, Brown Thrasher and Mockingbird were trying to outdo one another.

After having made a careful survey of this territory and also of Rivermount Park, another splendid place for the birds, where they seemed to have lost all shyness, and gaudy Summer and Scarlet Tanagers, Orioles and Wood Thrushes would sit only a few feet away from you. I decided to make May 5th an All Day with the Birds, but I became ill and for three days had a severe fever. As soon as it left me, however, I decided I had to go at once, and May 8th found me at work. My previous careful study enabled me to come out of the woods at 10 a. m. with 79 species seen, then I went to the James River, but work there was a sore disappointment. No doubt our secretary, Mr. John W. Daniel, Jr., whose home is at Lynchburg, could have done far better than I did, as he no doubt would know the haunts of the birds there. Suffice it to say the James River only netted me eight new species in three hours of the hardest kind of work, but it included a species I had not yet seen here—the Warbling Vireo. Rivermount Park furnished the greatest surprise of the day—a pair of Philadelphia Vireos,—a late date for this bird as also for the Hermit Thrush, but in Lynchburg, as everywhere, the past winter had been a severe one and no doubt this kept back many a migrant.

By 5:30 p. m. I began to feel the effects of the three days of fever and was compelled to go home and remain quiet, but still I was able to add a few more, till at 9 p. m. the doleful note of the Whippoorwill gave me my 105th species. Taking into consideration my physical condition and the fact that

I worked in a new territory, may account for the smaller list, but at the same time it also added a new charm to bird study to see what could be done for an "All Day" record under such circumstances.

Lynchburg, Pa., May 8, 1905.—Weather warm, about 85°, sun bright from 6 till 12, then sky overcast with clouds, strong wind in evening. Leaving Durmid 6 a. m., two miles south through woods and hills, half way up the Candler Mountains till 10 a. m. (species seen, 79); from 11 a. m. from Durmid to 12th street depot (N. & W.) Lynchburg, new species seen, 3; car to Union depot, walking on left side of James River east four miles, returning by way of Island till 3 p. m. (new species added, 8); car to Rivermount Park, where I staid till 5:30 p. m. (new species added, 11); returning to Durmid, where I added four species, the last one at 9 p. m. Total species seen, 105; total individuals, 604.

Bewick Wren, 15; Robin, 2; Mockingbird, 5; Chipping Sparrow, 8; Catbird, 20; Brown Thrasher, 7; Meadowlark, 1; Maryland Yellow-throat, 6; Field Sparrow, 10; Yellow Warbler, 10; Yellow-breasted Chat, 6; Tufted Titmouse, 2; Green-crested Flycatcher, 6; Mourning Dove, 2; Goldfinch, 20; Killdeer, 1; Kingbird, 5; Nighthawk, 2; Vesper Sparrow, 2; Bachmann's Sparrow, 12; Towhee, 15; Crested Flycatcher, 6; Black and White Warbler, 6; Redstart, 25; Traill Flycatcher, 2; Carolina Wren, 3; Black-throated Blue Warbler, 4; Prairie Warbler, 15; Tennessee Warbler, 2; Black-throated Green Warbler, 8; Black-poll Warbler, 9; Pine Warbler, 4; Cape May Warbler, 4; Chestnut-sided Warbler, 20; Blue-winged Warbler, 2; Worm-eating Warbler, 2; Bay-breasted Warbler, 2; Parula Warbler, 4; Nashville Warbler, 1; Golden-winged Warbler, 4; Magnolia Warbler, 1; Connecticut Warbler, 2; Myrtle Warbler, 1; Wilson Warbler, 2; Canadian Warbler, 4; Summer Tanager, 4; Red-headed Woodpecker, 4; Grasshopper Sparrow, 2; Carolina Chickadee, 3; Cardinal, 20; Indigo Bunting, 4; Phœbe, 1; Wood Pewee, 12; American Crow, 7; Yellow-throated Vireo, 3; Red-eyed Vireo, 40; Solitary Vireo, 2; White-eyed Vireo, 1; Warbling Vireo, 4; Oven-bird, 12; Blue-gray Gnatcatcher, 8; House Wren, 3; Wood Thrush, 18; Flicker, 3; Louisiana Water Thrush, 2; Wilson Thrush, 2; White-breasted Nuthatch, 2; Turkey Vulture, 50; Bob-white, 2; Chimney Swift, 9; Bluebird, 6; Red-winged Blackbird, 2; Purple Martin, 2; Downy Woodpecker, 1; Loggerhead Shrike, 1; Sharp-shinned Hawk, 1; Orchard Oriole, 3; Sparrow Hawk, 1; Yellow-bellied Flycatcher, 1; Song Sparrow, 6; Bank Swallow, 2; Blue Grosbeak, 4; Least Flycatcher, 2; Rough-winged Swallow, 8; Spot-

ted Sandpiper, 1; Green Heron, 1; Yellow-legs, 1; Kingfisher, 1; Baltimore Oriole, 3; Hairy Woodpecker, 1; White-throated Sparrow, 1; White-crowned Sparrow, 1; Hermit Thrush, 2; Ruby-throated Hummer, 1; Blue Jay, 3; Barn Swallow, 1; Broad-winged Hawk, 1; Cowbird, 4; Purple Grackle, 2; Whippoorwill, 2; Blackburnian Warbler, 11; Cerulean Warbler, 2; Scarlet Tanager, 10.

SOME FURTHER SUGGESTIONS FOR BIRD STUDY.

BY LYNDS JONES.

We may keep plainly in view two main objects of "All Day" studies, such as Christmas, New Year, May, and July. The first object, as it lies in the writer's mind, is to stimulate interest in outdoor studies. The second is to determine, as far as possible, what birds inhabit the given region at the time of the study. The first object is certainly a worthy one, and the second is the beginning of the carefully prepared local list. It is the oft repeated and firm conviction of the writer that a conservative local list, whether it cover but a small area surrounding some village, or more ambitious, covers a whole county, is one of the most valuable helps for the beginner in bird study. It reduces the liability to gross mistakes in the almost universal expectation, shall I say, of the marvelous in bird life. If the local list is prepared by some person who knows what birds should be found in that locality a list of the species which have not been found up to the preparation of the list, but which should be found there may well be appended. Such a list is certain to stimulate interest in the local study. Who shall be first to add one and another of these species to the real list? These points have all along been given more emphasis than any others because it has been assumed that most of the members of The Wilson Ornithological Club have not yet attained to the degree of knowledge where special studies can profitably be undertaken. There are, however, not a few members who are capable and ready to undertake special work, and it is to such persons that I wish to speak further.

There is a great deal of work that needs to be done before we know anywhere near all about the birds, some of it out of reach of the average person who knows birds well, some of it within the reach of every person. Mention has many times

been made of the need for more extensive and more exact knowledge of the breeding habits of the birds. The nest location, nest material (of what sort and where obtained), how the nest is made, by both or only one parent; the possibility that two pairs of birds unite in one nest, particularly among the more gregarious species; the relation of the nest construction to season and weather; the time in building; the real office it performs in the act of breeding for parents, eggs and young.

There is a whole chapter in the "story of the birds" in the eggs. Who will write that chapter? Some of us may be strongly inclined to the oological side of bird study. If so here is a nearly untrodden field. It is for you, if you feel this tendency, to bring forward a new oology.

A little aside from the subject of oology proper is the embryology of nearly all of our native birds. The concern here is rather with the late than with the earlier stages of incubation. At what time do the first suggestions of the future feathers appears as little papillæ on the skin surface? The 5th day of incubation will probably be found the earliest date. How rapidly do these papillæ grow? What is their condition at hatching? When do the true feathers begin to appear beneath the skin and at the lower end of the down? How are these feathers arranged on the body? What is their rate of growth? How does their rate of growth correspond to the growth of the young bird? Why is down first followed by feathers? What is the relation of the color pattern of the nestling or downy young to the nest and other environment? What is the relation of the color pattern of the downy young to the first feather plumage, and of the first feather plumage to succeeding ones? Just here lies the secret to the development of the color patterns because we have given almost no attention to the downy young and to the earliest stages of the first feather plumage. It is a very enticing field for the original worker.

Coming to the adult birds, we don't begin to know yet all about their geographical distribution and migrations. The winter distribution and the spring migrations are comparatively easy and fairly well known, because at these times peo-

ple feel the inspiration to get out for actual field study. We are also coming to know something about the autumn migrations, but even yet far too little in a particular way. The fall migrations are not so easy to study, and the inducements to it are less enticing, and the difficulties greatly increased for most of us. One must become familiar with the least conspicuous plumages to know what birds he finds, and that is certainly not an easy task, but it can be accomplished. The time of year which has been the most neglected is the summer season. There are two main objects for the summer study, each worthy and each equally important. The first and most evident one is to determine exactly the breeding area of that species. The second reason is to determine the time when the southward migration actually begins. We need to know this as accurately as we know the time of the beginning of the northward migration before we can assume that we know much about it. There are a great many questions about the southward migration that need a conclusive answer, but the data must first be gathered.

Let no one suppose that he can do nothing with these suggestions because he cannot attempt something with all of them. One minor point under one of the minor heads is worth earnest effort and will repay anyone who has the courage and enterprise to undertake the study. Let those of us who can undertake some of these more particular studies, while those who cannot, for whatever reason, do what they can in their own localities.

TWO GOOSE RECORDS FROM OHIO.

W. F. HENNINGER.

On the morning of March 24, 1905, one of the proprietors of the Empire Hotel here at Tiffin shot two wild geese on the Sandusky River, one and one-half miles south of the city. Thanks to the persistent efforts and tireless energy of Mr. Karl J. Heilmann, I was able to secure the skins on condition of returning the meat to the hunter, to which I gladly consented. One of them was a fine old male of the Blue Goose (*Chen cærulescens*). Measurements: Extent, 168 ctm.; bill, 6 ctm.;

wing, 48 ctm.; tail, 20 ctm.; tarsus, 8 ctm.; length without bill, 89 ctm. Weight, 11½ lbs.

The books say the bill of this species is entirely purplish red in life, but the bill of this specimen was of a straw yellow, the purplish red being confined to two stripes parallel to the culmen. The other species was a male of the Hutchins Goose (*Branta canadensis hutchinsii*): Measurements: Extent, 153 ctm.; bill, 5 ctm.; tarsus, 7½ ctm.; wing, 43 ctm.; tail, 16½ ctm. (14 tail-feathers); length, without bill, 78 ctm. Weight, 5¾lbs. Both specimens are now in my collection. The Hutchins Goose is to my knowledge the first Ohio specimen preserved in one of the collections of the state. It is indeed a streak of good luck to secure two such rareties in one day.

ADDITIONS TO THE BIRDS OF OHIO.

LYNDS JONES.

Anhinga anhinga.—Anhinga. Mr. C. H. Morris informs me that a specimen of this southern species was taken in the Muskingum river at Lowell, Ohio, in Morgan county, in November, 1885, by a Mr. Davis. The specimen was mounted by Mr. Davis and is now in his private collection. Mr. Morris photographed the specimen and sent a print to me. At the time of writing this note particulars are wanting, but they will be ascertained and announced later. This species should be added to the list of Accidental Birds of Ohio.

Milvulus forficatus.—Scissor-tailed Flycatcher. Robert McCrory, Esq., reports a specimen of this southwestern species at Marysville, Ohio, the last of May, 1903. A careful description and sketch sent with the announcement seem to leave no doubt of the authenticity of this identification. Pariculars are lacking at this writing, but will be forthcoming for the next issue of the Bulletin. This, if corroborated, is the second occurrence for this bird in Ohio.

THE WILSON BULLETIN.

A Quarterly Magazine Devoted to the Study of Living Birds.
Official Organ of the Wilson Ornithological Club.

Edited by LYNDS JONES.

PUBLISHED BY THE CLUB, AT OBERLIN, OHIO.

Price in the United States, Canada and Mexico, 50 cents a year, 15 cents a number, postpaid
Price in all countries in the International Postal Union, 65 cents a year, 20 cents a number.
Subscriptions may be sent to Lynds Jones, Oberlin, Ohio, or to Mr. Frank L. Burns, Berwyn, Penn., or to Mr. John W. Daniel, Jr., Washington, D. C.

EDITORIAL.

More and more local lists of birds are being called for by teachers and others who are either trying to interest others in bird study, or who are beginning that study themselves. The Wilson Bulletin always welcomes local lists which have the marks of careful preparation. If authors of such lists desire separates of their lists, arrangements for such separates should be arranged for before the type is thrown down.

For a summer "All Day" we suggest Saturday, July 1st. Such a day will be well within the summer season when no southward movement will likely have begun anywhere within the United States, and will avoid the objection which has been felt to the 4th, when some people wish to celebrate in the good old fashion. It will also give a glimpse of the summer population of your region. All such lists which are sent to the editor will be put into tabular arrangement for the September Bulletin.

The half tones which appear in this number are from photographs taken in the vicinity of Woods Hole, Mass.

To the many kindly inquiries regarding the line of the editor's study while he was at the University of Chicago he wishes to say that it was specifically a careful study of the development of the first down and the relation of that down to the first feather, a feather always following a down and pushing the down out upon its tip. Some may be interested to know that the recall from Chicago to take up the work of Professor Albert A. Wright, who died suddenly on April 3, has not made necessary the complete cessation of the University work. The editor has reasonable hope of being able to complete the work begun before the summer.

GENERAL NOTES.

SOME NESTING SITES OF THE HAIRY WOODPECKER (*Dryobates villosus*.—According to my experience the Hairy Woodpecker is not a common breeder in this county, the majority of birds noted during the period of nidification being transitory, as many a time I have followed a bird to the woodland margin and seen it launch forth, bound for some distant piece of timber. The four nesting sites I have noted were in thick woods and discovered when least expected.

Site 1. P. C. 618, Grosse Pointe Twp., June 11, 1899.—Excavation in trunk of very large barkless dead elm about 50 feet above ground and 20 below first limb. My attention was directed to it by watching the birds conveying food. The clamor of the young could be distinctly heard.

Site 2. Section 31, Nankin Twp., May 6, 1900.—Excavation in trunk of dead beech 55 feet up and just under large limb. Contained four slightly incubated eggs. Depth of cavity, eight inches. Was located April 22, by watching the birds.

Site 3. Section 3, Van Buren Twp., May 17, 1903.—Excavation in poplar stub about fifteen feet above ground in thick second growth containing many dead trees and stubs. Young responded in joyous chorus to knock upon stub base.

Site 4. Section 11, Brownstown Twp., May 23, 1903.—Total height of beech about 40 feet and only the upper three feet of it dead. Excavation near center of dead portion. Young noisy and could be heard for a considerable distance whenever parents approached with food. Frequently protruded their heads from cavity and were apparently about ready to leave nest. Parents very solicitous and fearless. J. CLAIRE WOOD.

Wayne Co., Mich.

RUBY-CROWNED KINGLET IN WINTER.—On February 16, on the Ohio State University campus, I observed a Ruby-crowned Kinglet (*Regulus calendula*) in a flock of about one hundred Golden-crowned Kinglets (*Regulus satrapa*). While the specimen was not secured I am convinced that the bird was *calendula*. I am fairly familiar with this brid as an April and October migrant; and further, I was able to contrast it directly. ZENO METCALF.

THE DICKCISSEL (*Spiza americana*) IN WAYNE CO., MICH.—The Dickcissel is a rare summer resident in Wayne county. This assertion, as to its rarity, is based upon the fact that while business and pleasure takes me into various sections of the county every summer, I have met with it during two years only, and furthermore, no one else has observed it here so far as I know.

The first bird was singing from the top of an elm sapling on the margin of a dense bush land divided by a swale of marsh grass

and rushes in Private Claim 50, Ecorse Twp. This was in May, 1899, and the bird was seen on several subsequent visits. On July 30th, of the same year, I discovered a colony of about 15 pairs on P. C. 405, Village of Delray, and about a mile from the other locality. They had taken possession of some ten acres of neglected land, covered mostly with thick weeds, but there was a mud flat in one portion fringed with rushes. This field is surrounded by factory buildings, railroad tracks and the River Rouge. I secured a male, but, despite protest, my companion collected four males and two females, saying the birds were rare and we might not get another chance. I found an empty nest, probably of this species, fastened to the weeds about a foot above the ground. While examining it I noticed a motion of the weed tops a few feet away and investigation discovered a young Dickcissel not yet able to fly. No birds have since been seen in the two above localities.

The next bird was perched upon a wire fence between two meadows in Section 19,Monguagon Twp., about the middle of May, 1904, and the last seen was a male and female on July 20th, in the bushes along a fence between a corn and clover field on P. C. 26, Village of Fairview. J. CLAIRE WOOD.

SOME SPRING RECORDS FROM THE VICINITY OF NEW YORK CITY.—Central Park, March 5.—150 Snowflakes remained about a week.

March 7.—3 American Crossbills.

March 26.—1 Mourning Dove.

May 8.—13 Bay-breasted Warblers.

May 10.—1 female Blue-gray Gnatcatcher.

Leonia, N. J., March 25.—1 Baldpate.

May 6.—2 Tufted Titmice.

Vicinity of Englewood, N. J., May 13.—1 Brewster's Warbler (*Helminthophila leucobronchialis*) observed in an elm tree by the road eating the fruit. CARLETON SCHALLER.

New York City.

BOOK REVIEWS.

Gleanings No. IV. Some Notes on the Summer Birds of Monongahela Co., W. Va. By J. Warren Jacobs.

We are always pleased to receive these Gleanings from time to time, not alone because they possess intrinsic worth, but because they illustrate in an unmistakable way one persons lively interest in what the birds about him are doing. The observations made on two short trips introduce us to some of the features of a region all too little known. We trust that Mr. Jacobs may yet find time to thoroughly work this territory. L. J.

Bird Study in the Rural School. Normal School Bulletin, No. 12, April 1, 1905. By Thomas L. Hankinson, B.S.

In ten pages Professor Hankinson has discussed in a pleasant manner the possibilties of bird study without either museum or collection or teacher, and has clearly shown that there are no insurmountable difficulties. The writer has a very tender memory of the rural school, for it was there that his early education was wholly obtained. There, too, all of his early experiences in bird study were cast. If he still sighs for the old times and scenes who shall say him nay! We cannot but envy the rural teacher and scholar their unrivalled opportunities for bird study at first hand. L. J.

The Mammal and Bird Fauna of Beaver County, Pennsylvania. By W. E. Clyde Todd. (Reprinted from History of Beaver County, Pennsylvania, at its Centennial Celebration.)

Some pertinent remarks upon the past fauna and the influence of settlement upon that fauna, and remarks relating to the zoogeographic position of Beaver County, are followed by an annotated list of 30 mammals and 178 birds. We welcome it as a further contribution to faunal literature. L. J.

The Origin of the Kirtland's Warbler. By P. A. Taverner. From the Ontario Natural Science Bulletin, No. 1, April 15, 1905.

The conclusion of this interesting paper may be given in the author's own words. "It seems probable, then, from our present knowledge, that Kirtland's Warbler is what remains of a once far more widely distributed species wintering along the Gulf States and spread from thence to the Bahama Islands. . . . The breeding grounds then likely covered the greater part of the Alleghanian Transition Zone." L. J.

The Ontario Natural Science Bulletin comes to us as an entirely new 48 page claimant for attention, under the able editorship of Mr. A. B. Klug. The first number contains about 28 pages of interest to bird students, and in all respects gives promise of being a valuable addition to the literature of natural science. The editor's list of 197 Wellington County birds, and his migration tables are worthy of special notice. We wish this new venture into the field of scientific literature every success. L. J.

The Warbler, in its new form and cover, steps into the ranks of bird magazines with the apparent purpose of presenting a series of accurate colored pictures of rare eggs and photographs of nests and eggs and nesting sites, and if the first two numbers are an indication of the future policy, to giving fuller accounts of birds in particular localities than most magazines seem to be willing to

give room for. It is well printed and its arrangement is good. Mr. John Lewis Childs, the editor, is to be congratulated upon the change from the first series. L. J.

With the first number of Volume VII., The Journal of the Maine Ornithological Society makes larger claims for our attention, in a new cover design, better paper, better print, and an increased number of pages. We are pleased to note that the Maine ornithologists are making good progress in bringing the state bird fauna upon a firm basis. It is inevitable that a state so old and so much written about should have amassed a great deal of questionable authenticity, particularly among the earlier writers. It is far more difficult to remove a species from a state list than to place one upon it. L. J.

It will be a source of pleasure to all bird lovers to know that Mr. Frank Chapman is preparing a complete revision of his admirable Handbook of the Birds of Eastern North America. It would have been ready ere this but for the delay of the committee having in charge another revision of the birds of North America. Changes have been so rapid and so radical these last few years, that an author always runs the risk of finding his work obsolete so far as the nomenclature is concerned, almost before the ink is dry. We hope for a fairly stable nomenclature. L. J.

PUBLICATIONS RECEIVED.

Amateur Sportsman, Vol. XXXII, Nos. 4, 5, 6.

American Ornithology, Vol. V, Nos. 3, 4, 5.

Bird-Lore, Vol. VII, No. 2.

Boys and Girls, Vol. IV, Nos. 3, 4, 5.

Bulletin No. 20, U. S. Dep't Agri. Biol. Survey. Coyotes in their Economic Relations. By David E. Lantz.

Bulletins 119, 120 120, 121, 122, University of Wisconsin Agricultural Experiment Station.

Bulletin 70, Pennsylvania State College Agriculture Experiment Station.

Colorado College Studies, Science Series. Vol. XI, Nos. 39–41.

Condor, The, Vol. VII, Nos. 2, 3.

Journal of the Maine Ornithological Society, The, Vol. VII, No. I.

Maine Sportsman, Vol. XII, Nos. 138, 139, 140.

Mitteilungen uber die Vogelwelt, Vol. IV, No. 7.

Naturaliste Canadian, Le, Vol. XXXII, Nos. 2, 3, 4.

Ohio Naturalist, The Vol. V, Nos. 4, 5, 6.

Ontario Natural Science Bulletin, The, Vol. I, No. 1.

Ornithologische Monatschrift, Vol. XXX, Nos. 2, 3.

Warbler, The, Vol. I. Nos. 1, 2.

Wilson Chapter Publications

Consists of two Series : Old and New

The Old Series comprise the following issues :

Ornithologists' and Oologists' Semi-Annual, Vols. I and II, two numbers each, Vol. III, one number. (No. 1 of both Vols. I. and II are out of print.) 25 cents a number.

Wilson Quarterly, Vol. IV, two numbers. 25 cents a number.

The Journal, two numbers. 10 cents a number.

The whole series (available numbers) $1.00.

The New Series comprise the

Wilson Bulletins, from No. 1 to 51 inclusive. (Nos. 1, 2, 3, 4, 5, 6, 7, 8, 24, are out of print.)

Of the Wilson Bulletins, No. 15, is a study of "The Oberlin Grackle Roost," 18 pages, by LYNDS JONES. Price 15 cents. No. 30, "Warbler Songs," 56 pages, by LYNDS JONES; in which all North American Warblers are discussed, the songs of nearly all described, and a field key to the adult males given. Price 50 cents. No. 31, "A Monograph of the Flicker," 82 pages, by FRANK L. BURNS. Price 50 cents. No. 33, "A Summer Reconnoissance in the West," by LYNDS JONES and W. L. DAWSON, being a study of the birds in fourteen states during a Journey of 7000 miles. Price 20 cents. No. 37, "A Sectional Bird Census," by FRANK L. BURNS. Price 25 cents.

The other numbers consist of "General Notes." Price 15 cents each. The whole available New Series for $5.00.

Address all communications to

LYNDS JONES, Oberlin, Ohio.

A History of North American Land Birds

BY

S. F. BAIRD, T. M. BREWER, *and* R. RIDGWAY

With more than 1000 Illustrations, including Colored Frontispieces; 64 full-page plates containing 545 heads of birds, printed in colors, and 593 wood engravings.

THIS important work on American ornithology, giving a larger amount of information than any other work upon the subject, has hitherto been published without any illustrations in colors at $24. It is now issued in volumes which are much handier in size, and at a price which is extremely low, taking into consideration its wealth of illustration (including many in colors) and immense amount of information.

An especially important feature of the work, one which greatly illuminates the subject, is the portion written by Dr. Brewer, containing the most graphic and accurate accounts of the habits of the birds which have been penned since Audubon.

Other valuable features are the statements given of the exact rank of each bird, the analytical and synoptical tables simplifying identification, accounts of the geographical distribution in America and other parts of the world, descriptions of varieties not in other works, etc.

3 Volumes. Large Octavo. Over 1800 pages. Cloth. Gilt Top, in Box. Price, $10. Express charges prepaid. Address, The Wilson Bulletin, Oberlin, Ohio.

IF YOU

desire to keep in touch with the Ornithology of Ontario you should read

The Ontario Natural Science Bulletin

An Annual devoted to the Fauna and Flora of Ontario.

Ontario's Best Ornithologists are its contributors.

No. 1. Issued April 15, 1905.

Price 25 cents per copy.

ORDER AT ONCE, as over 1200 copies are already sold.

THE BUSINESS MANAGER, "Ont. Nat. Sc. Bull.," Box 668, GUELPH, ONT.

THE WARBLER

JOHN LEWIS CHILDS, Editor

WITH 1905 **The Warbler** begins a new series which will contain many superb Colored Plates of rare eggs such as Kirtland and Olive Warbler, Carolina Parroquet, Clark's Crow, Ipswich and Rufous Crowned Sparrow, Yellow and Black Rail, Calaveras Warbler, etc. Also splendid illustrations of Birds and Nests, and leading articles by well known authorities.

Published Quarterly, 32 Pages & Cover

SUBSCRIPTION PRICE

$1.00 PER YEAR. SINGLE COPIES 30c

Eggs of Kirtland Warbler will be figured in first issue (Jan. or Feb.) of the new series.

ADDRESS THE WARBLER
FLORAL PARK, N. Y.

Bird Papers For Sale

Complete volumes as well as odd numbers of Birds and Nature, Osprey, and many others.

Write for list and prices.

LYNDS JONES,
Oberlin, O.

DAWSON'S BIRDS OF OHIO.

Profusely illustrated, 80 full-page colored pictures, 200 original half-tones, 720 quarto pages. 322 birds accurately described, their habits interestingly given. Combines ready identification keys with adequate descriptions of habits as well as plumage and form. Far and away the best bird book of its kind. Send to-day for descriptive and sample pages and illustrations to WILSON BULLETIN, Oberlin, Ohio.

A REVISED CATALOGUE OF THE BIRDS OF OHIO.

BY LYNDS JONES.

This annotated catalogue of 241 pages is without illustrations, except a map of Ohio. In it the feeding habits, migrations and distribution of the birds of Ohio are treated. It does not attempt a description of the plumage. Price is 50 cents. Address, LYNDS JONES, Oberlin, Ohio.

Veterinary Elements.

What the Animal Husbandry Expert, Mr. Geo. M. Rommel, B. S. A., of the U. S. Dep't. of Agriculture, has to say regarding "Veterinary Elements":

"I do not feel that I can add anything to that which has been already said in commendation . . . . but from my own personal experience I can say that if I could have had the training in veterinary work that the little book embodies, my course in animal husbandry would have been all the more valuable to me.

"You have pointed out the way for a splendid line of work, and I sincerely hope that the leaders in agricultural education will not be slow in following it."

The author is A. G. Hopkins, B. Agr., D. V. M.; Farmer and Agricultural Journalist; late instructor in Stock Judging and Veterinary Science, Wisconsin Agricultural College.

The above work is clearly and plainly written and easily understood by any one. It can be secured, postpaid, for $1.50, from

THE WILSON BULLETIN, Oberlin, Ohio.

New and Revised Edition Just Issued.

Principles of Plant Culture.

BY E. S. GOFF,

Professor of Horticulture in the University of Wisconsin.

The text-book used in the classes in Plant Life and Horticulture, in the popular Short course in Agriculture, of the University of Wisconsin.

A Few of the Many Things that this Book Teaches.

How to select, test, plant store, and stratify seeds. How to promote flowering and fruiting. How to prevent sun-scald on fruit and other trees. How to promote wood maturity in autumn. How to water and care for greenhouse plants.

The Book Also Teaches

How to maintain the fertility of the soil. How to prevent damage from mice, gophers, rabbits, woodchucks, and birds. How to spray plants to prevent insect and fungus ravages. How to prevent damage from chinch bugs, army worms and grasshoppers. How to treat oats to prevent smut. How to make and apply the Bordeaux mixture. How to grow plants from cuttings, layers and stolons. How to propagate plants by grafting and budding. How to make and take care of hot-beds and cold-frames. How to transplant trees, shrubs and herbaceus plants. How to prune fruit and ornamental trees. How to improve plants by selection and cross pollination, and many other things equally valuable.

While the work is written in plain and concise language, the subjects are treated in a thoroughly scientific manner and embody the results of the most recent experiment and research.

The book contains 285 printed pages with 179 illustrations. It is well printed on excellent paper and is neatly and substantially bound in cloth.

This book will prove a most valuable addition to the library of every school of agriculture, horticulture, forestry, and of every rural high or common school in this country, as well as to the library of every farmer and fruit-grower.

Price, $1.00 per copy, postage paid, when ordered from

THE WILSON BULLETIN, Oberlin, Ohio.

Stanford University Library
OCT 19 1905

Old Series Vol. XVII. New Series Vol. XII, No. 3.

The Wilson Bulletin

No. 52.

Published Quarterly by the

WILSON ORNITHOLOGICAL CLUB

Edited by LYNDS JONES

September, 1905.

OBERLIN, OHIO.

Entered as second-class mail matter at the Postoffice at Oberlin, Ohio.



CONTENTS.

Laughing Gull—Frontispiece Illustration.

Bird Migration at Oberlin, O. ROB'T L. BAIRD. 75

Two Bird Days Near Washington, D. C. HARRY C. OBERHOLSER. 84

Further Notes on the Birds of Middle Southern Ohio.
W. F. HENNINGER. 89

The Drumming of the Ruffed Grouse. LYNDS JONES. 94

Two Weeks with the Birds in Kent, Conn. CARLETON SCHALLER. 97

July Horizons. 99

Information Wanted. 101

General Notes. 102

Widow Hummer. FRANK BRUEN.

Louisiana Water-Thrush in Philadelphia in Summer.
CHAS. H. ROGERS.

Editorials 103

THE WILSON BULLETIN

is mailed on the fifteenth of March, June, September and December, by the Wilson Ornithological Club at Oberlin, O., edited by Lynds Jones.

Subscription: Fifty cents a year, including postage, *strictly in advance.* Single numbers, 15 cents, unless they are "Special" numbers, when a special price is fixed. The BULLETIN, including all "Special" numbers, is free to all paid up members, either Active, Associate, or Honorary, after their election.

Subscriptions may be addressed to the editor, to Mr. Frank L. Burns, Berwyn, Pa., or to Mr. John W. Daniel, Washington, D. C.

Advertisements should be addressed to THE WILSON BULLETIN, Oberlin, Ohio. Terms will be made known upon application.

All articles and communications intended for publication, and all publications and books for review, should be addressed to Lynds Jones, Oberlin, Ohio.

Articles of general interest relating to bird life are solicited. They should be in the hands of the editor not later than the 15th of the month preceding publication.

LAUGHING GULL. (Larus atricilla.)
Typical Nest and Eggs.
Muskeget Id., Mass., July, 1904.

THE
WILSON BULLETIN
NO. 52.
A QUARTERLY JOURNAL OF ORNITHOLOGY
VOL. XVII. SEPTEMBER, 1905. No. 3

BIRD MIGRATION AT OBERLIN, OHIO.

BY ROB'T L. BAIRD.

In The Wilson Bulletin No. 40, September, 1902, an article upon "The Spring Migration of 1901, With an Average Table for Lorain County, Ohio," occurs under the writer's name. The presnt article has to do with the southward as well as with the northward migration, attempting to give the whole annual movement of the migratory birds. The former table of migrations was based upon a relatively few records, while the present one has for its basis all of the records made in the region of Oberlin for the past ten years.

For the greatest number of records I am indebted to Professor Lynds Jones, whose field studies have been so extensive about Oberlin. For many records I wish to express my thanks to Messrs. S. D. Morrill, D. F. Nye, and H. H. Skinner, who have shown great interest in the work and frequently accompanied us on trips. A large number of records are from my own note-books. Practically all of the records are from were made by myself, in the absence of Professor Jones.

It must be borne in mind that the dates given are for migraing birds. Several species which are listed among the migrants are represented all the year by a few individuals, while some which are given as passing north of this region to breed, and returning south of it to spend the winter, are represented all winter by a few individuals. Species belonging to both classes will be found mentioned in the appropriate places. Those persons who take the trouble to compare this series of

lists with that published in 1902 will discover some differences in the times of arrival from the south of several species. This difference is accounted for by the much greater number of records from which this list is made.

SPRING MIGRATION.

ARRIVAL		DEPARTURE
Feb. 15-Mar. 1	American Crow	November 1
	American Robin	November 15
	Bluebird	November 15
March 1-10	Black Duck	April 1
	Canada Goose	March 25
	Killdeer	November 15
	Broad-winged Hawk	
	Sparrow Hawk	Oct. 15-Nov. 15
	Northern Flicker	Oct. 15-Nov. 15
	Red-winged Blackbird	November 1
	Meadowlark	November 1
	Bronzed Grackle	November 1
	Song Sparrow	Oct. 10-Oct. 21
March 10-20	Red-breasted Merganser	April 25-May 10
	Baldpate	April 25
	Pintail	April 5
	American Golden-eye	March 28
	American Woodcock	
	Mourning Dove	October 25
	Marsh Hawk	
	Cowbird	November 1
	Rusty Blackbird	May 1-10
	Field Sparrow	October 15-25
	Fox Sparrow	April 25
	Towhee	October 25
	Migrant Shrike	October 31
March 20-31	American Merganser	May 1
	Mallard	March 28
	Greater Scaup Duck	May 20
	Lesser Scaup Duck	May 20
	American Coot	May 5
	Wilson Snipe	May 15
	Turkey Vulture	October 11
	Cooper Hawk	
	Belted Kingfisher	November 1
	Phœbe	October 15
	Chipping Sparrow	October 1-10
	Swamp Sparrow	October 1-10
	Brown Creeper	May 7